

Full wwPDB X-ray Structure Validation Report (i)

Sep 21, 2021 – 05:09 am BST

PDB ID : 7PN0

Title: Crystal structure of the Phosphorybosylpyrophosphate synthetase II from

Thermus thermophilus at R32 space group

Authors: Timofeev, V.I.; Abramchik, Y.A.; Kostromina, M.A.; Esipov, R.S.; Kuranova,

I.P.

Deposited on : 2021-09-04

Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as 541 be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.23.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.23.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	316	84%	12%	
1	В	316	79%	16%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



\mathbf{M}	ol Ty	рe	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	A]	DΡ	A	401	X	_	_	_



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ribose-phosphate pyrophosphokinase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	312	Total 2395	C 1523	N 420	O 445	S 7	0	2	0
1	В	308	Total 2356	C 1497	N 412	O 440	S 7	0	2	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	311	LEU	=	expression tag	UNP A0A7G5B4V6
A	312	GLU	-	expression tag	UNP A0A7G5B4V6
A	313	HIS	_	expression tag	UNP A0A7G5B4V6
A	314	HIS	_	expression tag	UNP A0A7G5B4V6
A	315	HIS	_	expression tag	UNP A0A7G5B4V6
A	316	HIS	-	expression tag	UNP A0A7G5B4V6
A	317	HIS	-	expression tag	UNP A0A7G5B4V6
A	318	HIS	_	expression tag	UNP A0A7G5B4V6
В	311	LEU	-	expression tag	UNP A0A7G5B4V6
В	312	GLU	_	expression tag	UNP A0A7G5B4V6
В	313	HIS	_	expression tag	UNP A0A7G5B4V6
В	314	HIS	-	expression tag	UNP A0A7G5B4V6
В	315	HIS	_	expression tag	UNP A0A7G5B4V6
В	316	HIS	=	expression tag	UNP A0A7G5B4V6
В	317	HIS	-	expression tag	UNP A0A7G5B4V6
В	318	HIS	-	expression tag	UNP A0A7G5B4V6


• Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
9	Λ	1	Total	С	N	О	Р	0	0	
2	Α	1	27	10	5	10	2	0	0	
9	D	1	Total	С	N	О	Р	0	0	
2	Б	1	27	10	5	10	2	0	0	

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0

• Molecule 4 is water.


\mathbf{N}	/Iol	Chain	Residues	Atoms	ZeroOcc	AltConf
	4	A	138	Total O 138 138	0	0
	4	В	114	Total O 114 114	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ribose-phosphate pyrophosphokinase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	106.29Å 106.29Å 333.29Å	Danagitan
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.77 - 1.85	Depositor
Resolution (A)	19.77 - 1.85	EDS
% Data completeness	99.8 (19.77-1.85)	Depositor
(in resolution range)	99.9 (19.77-1.85)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.33 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
D D.	0.186 , 0.221	Depositor
R, R_{free}	0.195 , 0.228	DCC
R_{free} test set	3111 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.40 , 41.4	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o , F_c correlation	0.95	EDS
Total number of atoms	5097	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.86	$1/2440 \ (0.0\%)$	0.89	2/3318 (0.1%)	
1	В	0.78	$1/2399 \ (0.0\%)$	0.88	1/3262 (0.0%)	
All	All	0.82	$2/4839 \ (0.0\%)$	0.89	3/6580 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(\operatorname{\AA})$
1	A	310	THR	C-N	19.13	1.78	1.34
1	В	51	GLU	CD-OE1	6.88	1.33	1.25

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	310	THR	CA-C-N	-5.85	104.33	117.20
1	В	35	ARG	CG-CD-NE	-5.64	99.95	111.80
1	A	313	HIS	CA-CB-CG	5.45	122.86	113.60

There are no chirality outliers.

All (2) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	В	307	SER	Peptide
1	В	308	LEU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2395	0	2446	31	2
1	В	2356	0	2410	33	0
2	A	27	0	12	0	0
2	В	27	0	12	3	0
3	A	20	0	0	0	0
3	В	20	0	0	0	0
4	A	138	0	0	1	0
4	В	114	0	0	1	0
All	All	5097	0	4880	65	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:A:310:THR:C	1:A:311:LEU:N	1.78	1.36
1:A:118:GLN:HE22	1:A:140:LYS:H	1.14	0.89
1:A:32:THR:CG2	1:A:46:GLU:OE2	2.29	0.80
1:A:32:THR:HG23	1:A:46:GLU:OE2	1.82	0.80
1:A:7[B]:ILE:CD1	1:A:24:LEU:CD1	2.64	0.74
1:B:118:GLN:HE22	1:B:140:LYS:H	1.35	0.72
1:A:7[B]:ILE:HD11	1:A:24:LEU:CD1	2.21	0.70
1:A:310:THR:CA	1:A:311:LEU:N	2.61	0.62
1:B:170:PRO:O	1:B:194:LYS:HD2	1.99	0.62
1:A:310:THR:C	1:A:311:LEU:CA	2.67	0.61
1:A:7[B]:ILE:CD1	1:A:24:LEU:HD12	2.30	0.61
1:A:7[B]:ILE:HD12	1:A:24:LEU:HD12	1.83	0.60
1:A:118:GLN:NE2	1:A:140:LYS:H	1.95	0.59
1:B:61:VAL:O	1:B:67:HIS:HD2	1.86	0.58



Continued from previous page...

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:53:ASP:OD2	1:A:300:HIS:HD2	1.88	0.57
1:A:155:TYR:CE1	1:A:159:ARG:HG3	2.40	0.56
1:A:118:GLN:HE21	1:A:141:ILE:HG12	1.70	0.56
1:B:170:PRO:O	1:B:194:LYS:CD	2.53	0.56
1:A:136:HIS:HD2	4:A:592:HOH:O	1.88	0.56
2:B:402:ADP:O2B	2:B:402:ADP:O2A	2.24	0.55
1:B:195:GLU:OE2	1:B:203:ARG:NH1	2.40	0.54
1:B:298:ARG:HD3	1:B:304:SER:O	2.07	0.54
1:A:272:THR:HA	1:A:286:LEU:O	2.07	0.54
1:B:136:HIS:HD2	4:B:562:HOH:O	1.92	0.53
1:B:309:PHE:CD1	1:B:310:THR:HB	2.44	0.52
1:B:147:SER:HB2	1:B:310:THR:HA	1.91	0.52
1:A:49:LEU:C	1:A:49:LEU:HD12	2.31	0.51
1:A:61:VAL:O	1:A:67:HIS:HD2	1.93	0.50
1:B:258:LEU:H	1:B:258:LEU:HD22	1.75	0.50
1:B:297:TRP:CE2	1:B:301:ARG:HG3	2.47	0.49
1:A:248:ALA:O	1:A:271:ALA:HA	2.13	0.49
1:B:118:GLN:HE21	1:B:141:ILE:HG12	1.77	0.49
1:A:149:GLU:HB2	1:A:150:PRO:HD3	1.95	0.48
1:B:258:LEU:HD12	1:B:283:LEU:HD11	1.96	0.48
1:B:44:ARG:NH1	1:B:46:GLU:OE2	2.47	0.47
1:A:169:ALA:HB3	1:A:175:LEU:HD13	1.97	0.47
1:B:131:HIS:NE2	2:B:402:ADP:PB	2.87	0.47
1:B:14:ARG:N	1:B:15:PRO:CD	2.78	0.47
1:A:169:ALA:CB	1:A:175:LEU:HD13	2.46	0.46
1:A:14:ARG:N	1:A:15:PRO:CD	2.79	0.46
1:B:131:HIS:NE2	2:B:402:ADP:O2B	2.49	0.46
1:B:35:ARG:NH2	1:B:66:ASP:OD2	2.48	0.46
1:A:7[B]:ILE:HD12	1:A:24:LEU:CD1	2.41	0.45
1:A:46:GLU:HA	1:A:46:GLU:OE1	2.17	0.45
1:A:247:ALA:HA	1:A:270:ALA:O	2.16	0.45
1:A:7[B]:ILE:HD11	1:A:24:LEU:HD13	1.98	0.45
1:A:220:ASP:O	1:A:248:ALA:HA	2.17	0.45
1:B:247:ALA:HA	1:B:270:ALA:O	2.17	0.45
1:A:9:SER:HB2	1:A:17:ALA:CB	2.48	0.43
1:B:87:THR:OG1	1:B:124:ARG:HG3	2.19	0.43
1:B:149:GLU:N	1:B:150:PRO:CD	2.82	0.42
1:B:215:THR:HB	1:B:243:GLU:HG3	2.01	0.42
1:B:147:SER:HB2	1:B:310:THR:CA	2.50	0.42
1:B:298:ARG:NH1	1:B:308:LEU:HD22	2.34	0.42
1:B:162:LEU:HD12	1:B:162:LEU:HA	1.91	0.41



$\alpha \cdots$	· ·	•	
Continued	trom	nromanne	naae
-	110111	picolous	payc

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	overlap (Å)
1:B:220:ASP:O	1:B:248:ALA:HA	2.21	0.41
1:B:297:TRP:CZ2	1:B:301:ARG:HG3	2.55	0.41
1:B:248:ALA:O	1:B:271:ALA:HA	2.20	0.41
1:A:95:TYR:O	1:A:110:ALA:HB2	2.21	0.41
1:B:35:ARG:NH1	1:B:66:ASP:OD2	2.52	0.41
1:B:272:THR:HA	1:B:286:LEU:O	2.20	0.41
1:B:53:ASP:OD2	1:B:300:HIS:HD2	2.04	0.41
1:B:222:GLU:OE1	1:B:224[A]:SER:OG	2.39	0.41
1:B:65:GLN:NE2	1:B:95:TYR:OH	2.54	0.40
1:A:118:GLN:HE22	1:A:140:LYS:N	1.98	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
1:A:313:HIS:CE1	1:A:313:HIS:CE1[18_444]	1.63	0.57
1:A:313:HIS:CE1	1:A:313:HIS:NE2[18_444]	2.05	0.15

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	312/316 (99%)	304 (97%)	7 (2%)	1 (0%)	41	26
1	В	308/316 (98%)	293 (95%)	14 (4%)	1 (0%)	41	26
All	All	620/632 (98%)	597 (96%)	21 (3%)	2(0%)	41	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	200	THR
1	В	195	GLU



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	es
1	A	254/256~(99%)	240 (94%)	14 (6%)	21 7	
1	В	250/256~(98%)	234 (94%)	16 (6%)	17 4	
All	All	504/512 (98%)	474 (94%)	30 (6%)	19 5	

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	32	THR
1	A	41	LEU
1	A	49	LEU
1	A	101	LYS
1	A	159	ARG
1	A	175	LEU
1	A	187	LEU
1	A	194	LYS
1	A	200	THR
1	A	208	VAL
1	A	210	GLU
1	A	279	GLU
1	A	284	ARG
1	В	5	LEU
1	В	49	LEU
1	В	162	LEU
1	В	167	VAL
1	В	176	LYS
1	В	194	LYS
1	В	203	ARG
1	В	205	ARG
1	В	206	MET
1	В	215	THR
1	В	231	GLU
1	В	279	GLU
1	В	284	ARG



Continued from previous page...

Mol	Chain	Res	Type
1	В	308	LEU
1	В	309	PHE
1	В	310	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	GLN
1	A	67	HIS
1	A	118	GLN
1	A	136	HIS
1	A	164	ASN
1	A	238	GLN
1	A	300	HIS
1	В	65	GLN
1	В	67	HIS
1	В	118	GLN
1	В	136	HIS
1	В	164	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les				
MIOI	Type	Chain	res	rtes	res	nes	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	405	-	4,4,4	0.51	0	6,6,6	0.24	0				
3	SO4	A	403	-	4,4,4	0.64	0	6,6,6	0.25	0				
3	SO4	В	401	-	4,4,4	0.46	0	6,6,6	0.22	0				
3	SO4	A	402	-	4,4,4	0.27	0	6,6,6	0.30	0				
3	SO4	В	405	-	4,4,4	0.17	0	6,6,6	0.30	0				
3	SO4	В	403	-	4,4,4	0.30	0	6,6,6	0.35	0				
3	SO4	В	404	-	4,4,4	0.33	0	6,6,6	0.14	0				
3	SO4	A	404	-	4,4,4	0.35	0	6,6,6	0.21	0				
2	ADP	В	402	-	24,29,29	0.65	0	29,45,45	1.01	2 (6%)				
2	ADP	A	401	-	24,29,29	0.68	0	29,45,45	1.21	2 (6%)				

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	В	402	-	-	5/12/32/32	0/3/3/3
2	ADP	A	401	-	1/1/6/6	6/12/32/32	0/3/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	401	ADP	PA-O3A-PB	-3.92	119.39	132.83
2	В	402	ADP	PA-O3A-PB	-3.24	121.72	132.83
2	A	401	ADP	O3'-C3'-C2'	-2.50	103.75	111.82
2	В	402	ADP	O2A-PA-O1A	2.29	123.54	112.24

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	ADP	C3'

All (11) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	401	ADP	C5'-O5'-PA-O1A
2	A	401	ADP	C5'-O5'-PA-O2A
2	В	402	ADP	PA-O3A-PB-O2B
2	В	402	ADP	C5'-O5'-PA-O1A
2	В	402	ADP	C5'-O5'-PA-O3A
2	A	401	ADP	O4'-C4'-C5'-O5'
2	A	401	ADP	C3'-C4'-C5'-O5'
2	A	401	ADP	PB-O3A-PA-O5'
2	A	401	ADP	C5'-O5'-PA-O3A
2	В	402	ADP	C5'-O5'-PA-O2A
2	В	402	ADP	PA-O3A-PB-O1B


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	402	ADP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	310:THR	С	311:LEU	N	1.78



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	312/316 (98%)	-0.00	12 (3%) 40 38	10, 20, 45, 94	0
1	В	308/316 (97%)	0.09	18 (5%) 23 22	12, 21, 51, 81	0
All	All	$620/632 \ (98\%)$	0.04	30 (4%) 30 29	10, 20, 50, 94	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	198	SER	8.7
1	В	309	PHE	8.3
1	A	197	VAL	6.4
1	В	310	THR	6.1
1	A	199	ASP	5.0
1	A	200	THR	4.9
1	В	308	LEU	4.9
1	A	311	LEU	4.0
1	В	203	ARG	3.9
1	A	205	ARG	3.3
1	A	201	GLU	3.2
1	A	203	ARG	3.1
1	В	238	GLN	3.1
1	В	202	VAL	3.1
1	В	173	GLY	3.0
1	В	197	VAL	2.9
1	В	161	ASP	2.8
1	В	183	ARG	2.7
1	A	202	VAL	2.6
1	A	163	GLU	2.5
1	В	163	GLU	2.5
1	В	195	GLU	2.5
1	A	279	GLU	2.4
1	В	217	LEU	2.3



Continued from previous page...

Mol	Chain	Res	Type	RSRZ	
1	A	210	GLU	2.3	
1	В	196	ARG	2.2	
1	В	176	LYS	2.1	
1	В	279	GLU	2.1	
1	В	186	GLY	2.1	
1	В	200	THR	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	ADP	A	401	27/27	0.82	0.17	16,30,72,75	0
3	SO4	В	405	5/5	0.84	0.36	48,50,54,55	0
2	ADP	В	402	27/27	0.87	0.15	14,26,70,87	0
3	SO4	В	404	5/5	0.97	0.12	27,32,35,39	0
3	SO4	A	405	5/5	0.97	0.24	40,42,47,48	0
3	SO4	В	403	5/5	0.98	0.08	20,22,26,26	0
3	SO4	В	401	5/5	0.99	0.05	16,17,18,20	0
3	SO4	A	403	5/5	0.99	0.05	14,15,16,17	0
3	SO4	A	404	5/5	0.99	0.07	18,21,23,24	0
3	SO4	A	402	5/5	0.99	0.06	22,22,25,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

