



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2020 – 10:41 AM EDT

PDB ID : 4PO7
Title : Structure of the Sortilin:neurotensin complex at excess neurotensin concentration
Authors : Quistgaard, E.M.; Groftehaug, M.K.; Thirup, S.S.
Deposited on : 2014-02-25
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	685	6% (Poor fit) 82% (0 outliers), 13% (1 outlier), 1% (2 outliers), 0% (≥ 3 outliers)
2	N	13	15% (Poor fit) 38% (0 outliers), 31% (1 outlier), 15% (2 outliers), 16% (≥ 3 outliers)
2	P	13	38% (0 outliers), 8% (1 outlier), 54% (2 outliers)
3	B	3	100% (0 outliers)
3	C	3	67% (0 outliers), 33% (1 outlier)

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 5518 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sortilin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	658	5173	3269	870	1004	30	0	0	0


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	617	MET	VAL	variant	UNP Q99523
A	724	HIS	-	expression tag	UNP Q99523
A	725	HIS	-	expression tag	UNP Q99523
A	726	HIS	-	expression tag	UNP Q99523
A	727	HIS	-	expression tag	UNP Q99523
A	728	HIS	-	expression tag	UNP Q99523
A	729	HIS	-	expression tag	UNP Q99523

- Molecule 2 is a protein called Neurotensin/neuromedin N.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	N	9	81	55	10	16	0	0	0
2	P	6	51	32	12	7	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	B	3	39	22	2	15	0	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	3	39	22	2	15	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0

- Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	3	0
			13	8	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	104	Total	O	0	0
			104	104		
6	N	3	Total	O	0	0
			3	3		
6	P	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sortilin



- Molecule 2: Neurotensin/neuromedin N



- Molecule 2: Neurotensin/neuromedin N



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  67% 33%

MAG1
MAG2
BMA3

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.01Å 78.65Å 110.98Å 90.00° 126.62° 90.00°	Depositor
Resolution (Å)	20.97 – 2.66 20.97 – 2.66	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.97-2.66) 99.2 (20.97-2.66)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.67Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.3_1479)	Depositor
R, R_{free}	0.166 , 0.212 0.170 , 0.212	Depositor DCC
R_{free} test set	1621 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	68.4	Xtrriage
Anisotropy	0.491	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 59.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5518	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, BMA, PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/5294	0.43	0/7170
2	N	0.22	0/75	0.53	0/100
2	P	0.22	0/52	0.63	0/69
All	All	0.25	0/5421	0.44	0/7339

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5173	0	4968	56	0
2	N	81	0	76	1	0
2	P	51	0	45	1	0
3	B	39	0	34	0	0
3	C	39	0	34	2	0
4	A	14	0	13	1	0
5	A	13	0	18	1	0
6	A	104	0	0	2	0
6	N	3	0	0	0	0
6	P	1	0	0	0	0
All	All	5518	0	5188	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:680:ASN:HB3	1:A:698:ARG:H	1.40	0.87
1:A:515:LYS:HG2	1:A:526:THR:HG22	1.79	0.65
1:A:548:MET:HB2	3:C:1:NAG:H82	1.83	0.61
1:A:502:VAL:HG11	1:A:550:ILE:HD13	1.84	0.59
1:A:130:ASN:O	2:P:8:ARG:HG3	2.04	0.57
1:A:176:GLN:HG3	4:A:807:NAG:HN2	1.70	0.57
1:A:451:ALA:HB1	1:A:454:ILE:HD12	1.87	0.56
1:A:450:ASN:N	1:A:450:ASN:OD1	2.34	0.55
1:A:698:ARG:HG3	1:A:700:VAL:HG13	1.89	0.55
1:A:249:GLY:HA3	1:A:285:MET:HG3	1.89	0.54
1:A:471:ASP:H	5:A:808:PG4:H11	1.74	0.52
1:A:514:ILE:HD12	1:A:552:ILE:HD11	1.90	0.51
1:A:148:LYS:NZ	1:A:194:SER:OG	2.41	0.51
1:A:142:GLY:HA3	1:A:148:LYS:HB2	1.93	0.50
1:A:558:SER:HB3	1:A:560:LEU:HG	1.94	0.50
1:A:416:ASP:HB3	1:A:418:THR:H	1.77	0.50
1:A:220:HIS:CE1	1:A:238:THR:HG21	2.48	0.48
1:A:252:GLU:HG3	1:A:268:VAL:HG12	1.95	0.48
1:A:450:ASN:ND2	1:A:544:GLY:O	2.32	0.47
1:A:338:THR:HB	1:A:340:PHE:CE2	2.49	0.47
1:A:673:ARG:HB2	1:A:676:HIS:CD2	2.50	0.47
1:A:647:TYR:CD2	1:A:658:GLN:HG2	2.50	0.47
1:A:103:ILE:HG22	1:A:105:THR:H	1.80	0.46
1:A:704:LYS:HG3	1:A:705:LYS:N	2.30	0.46
1:A:699:GLU:OE1	1:A:699:GLU:N	2.49	0.45
1:A:682:TYR:CD2	1:A:696:PRO:HB2	2.51	0.45
1:A:306:MET:HE3	1:A:668:PHE:HD1	1.81	0.45
1:A:346:SER:HB3	1:A:353:TYR:CE1	2.52	0.45
1:A:386:LEU:HD11	1:A:390:ASN:HA	1.99	0.45
1:A:641:LEU:HG	1:A:685:ILE:HG12	1.99	0.44
1:A:360:HIS:NE2	1:A:383:THR:OG1	2.47	0.44
1:A:326:ASP:O	1:A:348:ASP:HA	2.18	0.44
1:A:520:GLU:OE2	1:A:614:LYS:HE2	2.18	0.44
1:A:179:LEU:HA	1:A:180:PRO:HD3	1.88	0.44
1:A:670:LEU:HD21	1:A:711:PHE:CE2	2.53	0.44
1:A:590:HIS:CG	1:A:602:ILE:HG13	2.54	0.43
1:A:258:ASP:OD1	1:A:258:ASP:N	2.44	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:606:LYS:O	1:A:630:PRO:HA	2.19	0.42
1:A:378:ARG:NH2	6:A:917:HOH:O	2.33	0.42
1:A:75:LEU:HB3	1:A:564:TRP:HD1	1.84	0.42
1:A:647:TYR:HD2	1:A:658:GLN:HG2	1.83	0.42
1:A:536:PHE:HZ	1:A:539:LEU:HG	1.85	0.42
1:A:318:TYR:HB2	2:N:13:LEU:HB2	2.02	0.42
1:A:704:LYS:HG3	1:A:705:LYS:H	1.84	0.42
1:A:312:VAL:HB	1:A:316:GLN:CB	2.50	0.42
1:A:702:ASP:OD2	1:A:704:LYS:HE2	2.20	0.41
1:A:90:VAL:HA	1:A:114:SER:O	2.21	0.41
1:A:680:ASN:N	1:A:680:ASN:OD1	2.53	0.41
1:A:548:MET:N	3:C:1:NAG:H82	2.35	0.41
1:A:713:SER:HA	1:A:714:PRO:HD3	1.89	0.41
1:A:387:SER:HB3	1:A:393:GLN:OE1	2.20	0.41
1:A:539:LEU:HD22	1:A:550:ILE:HD11	2.03	0.41
1:A:164:ILE:HG23	1:A:179:LEU:HD11	2.03	0.41
1:A:634:LEU:HA	1:A:691:GLN:O	2.22	0.40
1:A:646:TYR:CD2	1:A:681:GLY:HA2	2.56	0.40
1:A:68:HIS:HE1	6:A:975:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	652/685 (95%)	623 (96%)	27 (4%)	2 (0%)	41	56
2	N	5/13 (38%)	5 (100%)	0	0	100	100
2	P	4/13 (31%)	3 (75%)	1 (25%)	0	100	100
All	All	661/711 (93%)	631 (96%)	28 (4%)	2 (0%)	41	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	700	VAL
1	A	698	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	574/599 (96%)	561 (98%)	13 (2%)	50 68
2	N	8/12 (67%)	6 (75%)	2 (25%)	0 0
2	P	4/12 (33%)	4 (100%)	0	100 100
All	All	586/623 (94%)	571 (97%)	15 (3%)	46 64

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	ASN
1	A	163	ARG
1	A	278	ARG
1	A	362	TYR
1	A	365	THR
1	A	450	ASN
1	A	532	ASP
1	A	547	SER
1	A	548	MET
1	A	585	THR
1	A	680	ASN
1	A	698	ARG
1	A	700	VAL
2	N	2	LEU
2	N	11	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	N	1	2	7,8,9	1.82	1 (14%)	9,10,12	2.20	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	N	1	2	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	1	PCA	CD-N	4.70	1.47	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	N	1	PCA	CB-CA-C	-3.26	108.21	112.70
2	N	1	PCA	OE-CD-CG	-2.94	121.64	126.76
2	N	1	PCA	CA-N-CD	-2.82	103.94	113.58
2	N	1	PCA	CB-CA-N	2.50	110.48	103.30
2	N	1	PCA	CG-CD-N	2.35	114.48	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	1	1,3	14,14,15	0.27	0	17,19,21	0.56	0
3	NAG	B	2	3	14,14,15	0.41	0	17,19,21	0.44	0
3	BMA	B	3	3	11,11,12	0.76	0	15,15,17	0.96	0
3	NAG	C	1	1,3	14,14,15	0.41	0	17,19,21	0.70	1 (5%)
3	NAG	C	2	3	14,14,15	0.33	0	17,19,21	0.47	0
3	BMA	C	3	3	11,11,12	0.70	0	15,15,17	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	B	2	3	-	2/6/23/26	0/1/1/1
3	BMA	B	3	3	-	1/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	2.08	115.01	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	3	BMA	C4-C5-C6-O6
3	B	2	NAG	C8-C7-N2-C2
3	B	2	NAG	O7-C7-N2-C2
3	C	3	BMA	O5-C5-C6-O6
3	B	3	BMA	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	2	0

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PG4	A	808	-	12,12,12	0.59	0	11,11,11	0.22	0
4	NAG	A	807	1	14,14,15	0.27	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PG4	A	808	-	-	4/10/10/10	-
4	NAG	A	807	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	808	PG4	O3-C5-C6-O4
4	A	807	NAG	C3-C2-N2-C7
4	A	807	NAG	C4-C5-C6-O6
4	A	807	NAG	O5-C5-C6-O6
5	A	808	PG4	C3-C4-O3-C5
5	A	808	PG4	C1-C2-O2-C3
5	A	808	PG4	C8-C7-O4-C6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	808	PG4	1	0
4	A	807	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	658/685 (96%)	0.08	38 (5%) 23 19	46, 83, 126, 159	0
2	N	8/13 (61%)	0.48	2 (25%) 0 0	75, 101, 131, 131	0
2	P	6/13 (46%)	0.76	0 100 100	124, 129, 145, 148	0
All	All	672/711 (94%)	0.09	40 (5%) 21 18	46, 83, 129, 159	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	MET	7.1
1	A	102	VAL	5.2
1	A	55	ARG	4.9
1	A	246	ALA	4.4
1	A	103	ILE	4.1
1	A	423	ASN	4.0
1	A	53	CYS	3.8
1	A	560	LEU	3.7
1	A	106	PHE	3.7
1	A	242	GLY	3.7
1	A	531	ARG	3.7
1	A	105	THR	3.5
1	A	288	LYS	3.4
1	A	74	ASP	3.3
1	A	649	PRO	3.2
1	A	500	ILE	3.2
1	A	675	GLU	3.1
1	A	192	GLN	3.1
1	A	76	ARG	3.1
1	A	160	ARG	3.1
1	A	648	ARG	3.0
1	A	241	ASN	3.0
1	A	660	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	499	GLY	2.8
1	A	243	SER	2.7
2	N	5	ASN	2.7
1	A	152	THR	2.6
1	A	582	LYS	2.5
1	A	647	TYR	2.5
1	A	101	LEU	2.5
1	A	673	ARG	2.5
1	A	107	GLY	2.4
1	A	422	LYS	2.3
1	A	429	ILE	2.3
1	A	700	VAL	2.3
1	A	705	LYS	2.2
1	A	65	ASN	2.1
1	A	594	PRO	2.1
2	N	3	TYR	2.0
1	A	697	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	N	1	8/9	0.90	0.12	119,125,132,134	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	B	3	11/12	0.77	0.39	109,118,124,132	0
3	BMA	C	3	11/12	0.80	0.36	124,131,135,136	0
3	NAG	B	2	14/15	0.88	0.18	79,89,111,115	0
3	NAG	C	2	14/15	0.91	0.20	97,110,120,127	0
3	NAG	C	1	14/15	0.94	0.15	75,81,90,92	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	B	1	14/15	0.97	0.10	46,58,66,67	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	807	14/15	0.86	0.47	123,133,145,146	0
5	PG4	A	808	13/13	0.87	0.20	72,79,99,104	3

6.5 Other polymers [i](#)

There are no such residues in this entry.