



# wwPDB EM Validation Summary Report ⓘ

Apr 2, 2024 – 10:36 pm BST

PDB ID : 8PQV  
EMDB ID : EMD-17825  
Title : Cytoplasmic dynein-1 motor domain in post-powerstroke state  
Authors : Singh, K.; Lau, C.K.; Manigrasso, G.; Gassmann, R.; Carter, A.P.  
Deposited on : 2023-07-12  
Resolution : 4.00 Å (reported)  
Based on initial model : 7Z8G

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	4646	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 24122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytoplasmic dynein 1 heavy chain 1.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	2996	24122	15366	4163	4472	121	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1567	GLU	ARG	engineered mutation	UNP Q14204
A	1610	GLU	LYS	engineered mutation	UNP Q14204



PHE	ASN	PHE	GLN	GLY	LEU	VAL	ASP	ASP	LEU	LEU	ILE	ILE	GLY	GLU	PHE	ASN	VAL	
ASN	LYS	LEU	ASP	MET	GLN	THR	ASN	GLN	VAL	GLY	ILE	ARG	GLY	VAL	ASN	GLY	HIS	
GLU	LEU	ARG	ILE	THR	ASN	THR	VAL	GLN	PRO	ASP	GLY	ASP	GLY	PRO	ARG	GLY	ARG	
ALA	ALA	LEU	THR	THR	ARG	THR	PRO	GLY	ASP	GLY	ALA	GLY	THR	THR	VAL	GLN	GLN	
ALA	LEU	LEU	VAL	ILE	ILE	ARG	ARG	GLY	ALA	ASP	THR	THR	THR	THR	THR	THR	THR	
GLU	PHE	HIS	THR	GLN	PRO	GLN	PRO	SER	LYS	SER	LYS	THR	THR	THR	THR	THR	THR	
ARG	PHE	GLN	PHE	PRO	PRO	PRO	TRP	SER	TRP	TRP	TRP	TRP	TRP	TRP	TRP	TRP	TRP	
TRP	GLU	THR	THR	PRO	VAL	VAL	THR	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	GLY	
VAL	ALA	ALA	GLU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	LEU	
Q1327	D1328	E1335	E1341	D1344	Q1345	E1348	W1351	P1356	D1364	K1371	S1372	F1373	L1377	R1378	A1381	R1388	M1394	
I1395	M1397	M1398	V1400	I1401	R1411	H1412	W1413	K1414	Q1415	L1416	M1417	K1418	R1419	H1421	W1424	L1429		
W1435	L1439	Q1440	K1441	K1447	E1456	L1459	E1460	F1461	L1463	K1464	R1467	E1468	V1469	W1470	N1471	Q1481	C1484	
R1485	I1496	V1497	K1498	E1499	V1504	M1507	K1514	V1515	F1516	E1524	A1532	D1539	V1540	R1543	L1547	E1548	G1549	
I1550	F1551	T1552	G1553	K1558	E1567	I1571	K1581	Q1598	L1601	E1602	R1603	L1604	L1607	I1611	E1620	P1627	R1628	
F1628	N1671	L1672	L1673	L1674	L1675	L1676	L1677	S1678	R1679	E1680	E1683	I1698	W1701	L1702	V1721	V1724	I1739	
L1752	G1771	L1792	Q1800	R1804	D1831	E1871	Y1872	P1883	T1891	F1905	G1906	K1912	T1913	R1925	C1932	F1936	M1941	
I1944	F1945	V1946	G1947	L1948	V1951	G1952	W1954	D1958	R1966	H1967	L1968	S1972	Q1979	M1987	P1988	M1989	T1990	
D1991	A1995	P1996	L2002	N2003	K2004	G2005	V2006	K2007	D2011	M2012	T2017	P2020	G2021	ALA	GLY	ARG	S2026	
L2035	L2039	A2066	I2069	Q2079	L2080	Q2083	S2084	H2085	W2275	T2276	D2277	G2278	T2281	R2285	I2288	D2289	S2290	
V2291	R2292	L2295	R2298	G2299	W2300	D2304	G2305	D2306	V2307	D2320	P2336	F2337	M2338	W2442	T2445	Q2424	G2431	
L2432	V2433	T2434	K2435	A2436	L2437	R2451	L2452	R2453	C2454	L2455	C2466	P2460	R2488	L2499	D2505	S2506	R2507	
D2516	D2536	E2537	E2538	G2543	K2551	Q2552	P2553	A2563	V2569	P2570	T2571	L2572	R2576	L2581	V2592	F2606	G2619	
F2622	S2623	T2644	G2647	F2662	I2666	P2669	R2678	F2682										





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	67795	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	53	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.055	Depositor
Minimum map value	-0.020	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.013	Depositor
Map size (Å)	296.52002, 296.52002, 296.52002	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.059, 1.059, 1.059	Depositor



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/24633	0.58	12/33378 (0.0%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2002	LEU	CA-CB-CG	9.37	136.86	115.30
1	A	3683	ASP	CB-CG-OD2	9.34	126.71	118.30
1	A	3833	LEU	CA-CB-CG	7.26	132.00	115.30
1	A	3690	PRO	CA-N-CD	-6.97	101.75	111.50
1	A	4418	LYS	CA-CB-CG	6.58	127.89	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	24122	0	24177	278	0
All	All	24122	0	24177	278	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1416:LEU:O	1:A:1420:LEU:HB2	1.81	0.80
1:A:3502:THR:O	1:A:3504:ALA:N	2.22	0.73
1:A:1351:TRP:HB3	1:A:1429:LEU:HB3	1.74	0.70
1:A:3503:ILE:HG12	1:A:3503:ILE:O	1.93	0.67
1:A:2320:ASP:HB3	1:A:2358:ARG:HE	1.62	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	2984/4646 (64%)	2892 (97%)	89 (3%)	3 (0%)	51 84

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1996	PRO
1	A	3503	ILE
1	A	1421	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	2666/4125 (65%)	2609 (98%)	57 (2%)	53 72

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2897	LEU
1	A	4399	LYS
1	A	3124	ASP
1	A	4390	LEU
1	A	3923	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2964	HIS
1	A	3214	GLN
1	A	4386	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17825. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 140



Y Index: 140




Z Index: 140


### 6.2.2 Raw map



X Index: 140



Y Index: 140




Z Index: 140


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 114




Y Index: 170




Z Index: 150


### 6.3.2 Raw map



X Index: 114



Y Index: 170




Z Index: 150


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.013. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_17825\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 203 nm<sup>3</sup>; this corresponds to an approximate mass of 184 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i




\*Reported resolution corresponds to spatial frequency of 0.250 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.250 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.00	-	-
Author-provided FSC curve	3.93	5.93	4.00
Unmasked-calculated*	4.47	8.16	6.35

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.47 differs from the reported value 4.0 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-17825 and PDB model 8PQV. Per-residue inclusion information can be found in section 3 on page 4.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.013 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.013).







## 9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.013) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7940	 0.4160
A	 0.7930	 0.4160

