



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 7, 2023 – 04:14 PM EDT

PDB ID : 4Q0G  
Title : Crystal structure of beta subunit of acyl-CoA carboxylase AccD1 from Mycobacterium tuberculosis  
Authors : Bie, H.Y.; Yin, J.; James, M.N.G.; TB Structural Genomics Consortium (TB-SGC)  
Deposited on : 2014-04-02  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	529	 2% 94% 5%
1	B	529	 3% 91% 5%
1	C	529	 2% 94% 5%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 12306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable acetyl-/propionyl-CoA carboxylase (Beta subunit) AccD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	519	Total 3928	C 2479	N 699	O 731	S 19	0	0	0
1	B	505	Total 3846	C 2427	N 691	O 709	S 19	0	1	0
1	C	520	Total 3935	C 2484	N 700	O 732	S 19	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	221	Total	O	0	0
			221	221		
3	B	160	Total	O	0	0
			160	160		
3	C	192	Total	O	0	0
			192	192		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable acetyl-/propionyl-CoA carboxylase (Beta subunit) AccD1



- Molecule 1: Probable acetyl-/propionyl-CoA carboxylase (Beta subunit) AccD1



- Molecule 1: Probable acetyl-/propionyl-CoA carboxylase (Beta subunit) AccD1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	142.57Å 221.25Å 167.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.81 – 2.31 48.77 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.81-2.31) 99.4 (48.77-2.31)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.29 (at 2.32Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.189 , 0.213 0.181 , 0.201	Depositor DCC
$R_{free}$ test set	5807 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtrriage
Anisotropy	0.625	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 37.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12306	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/4017	0.53	0/5453
1	B	0.30	0/3930	0.53	0/5328
1	C	0.29	0/4025	0.53	0/5464
All	All	0.29	0/11972	0.53	0/16245

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3928	0	3884	12	0
1	B	3846	0	3819	8	0
1	C	3935	0	3892	8	0
2	A	6	0	8	0	0
2	C	18	0	24	0	0
3	A	221	0	0	0	0
3	B	160	0	0	1	0
3	C	192	0	0	0	0
All	All	12306	0	11627	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:ARG:HG2	1:A:41:ARG:HH11	1.17	1.02
1:A:41:ARG:HH11	1:A:41:ARG:CG	1.87	0.86
1:C:107:LYS:O	1:C:110:THR:HG23	1.79	0.82
1:A:76:TYR:O	1:A:79:GLU:HG2	1.85	0.77
1:A:41:ARG:HG2	1:A:41:ARG:NH1	1.87	0.74
1:A:268:TRP:HE1	1:A:331:HIS:HD2	1.45	0.63
1:B:107:LYS:O	1:B:110:THR:HG23	2.00	0.61
1:C:268:TRP:HE1	1:C:331:HIS:HD2	1.48	0.60
1:C:461:THR:O	1:C:461:THR:CG2	2.49	0.60
1:B:296:ASP:OD1	1:B:298:HIS:HD2	1.85	0.60
1:C:461:THR:O	1:C:461:THR:HG22	2.05	0.57
1:A:296:ASP:OD1	1:A:298:HIS:HD2	1.96	0.49
1:B:164:GLN:HE22	1:B:189:VAL:HA	1.78	0.49
1:A:406:VAL:HB	1:A:430:MET:CG	2.44	0.48
1:A:406:VAL:HB	1:A:430:MET:HG2	1.96	0.47
1:B:96:GLN:HE21	1:B:96:GLN:HA	1.80	0.47
1:B:505:THR:HG23	3:B:626:HOH:O	2.15	0.46
1:A:232:HIS:HA	1:A:236:SER:OG	2.14	0.46
1:A:200:ARG:NH1	1:A:244:ASP:OD1	2.49	0.46
1:B:296:ASP:OD1	1:B:298:HIS:CD2	2.66	0.45
1:A:473:ALA:HB3	1:A:474:PRO:HD3	1.99	0.45
1:B:200:ARG:NH1	1:B:244:ASP:OD1	2.51	0.43
1:B:192:MET:HG2	1:C:389:ALA:HB1	2.00	0.43
1:C:184:ALA:HA	1:C:207:LEU:O	2.19	0.43
1:C:35:ASN:HD22	1:C:35:ASN:C	2.23	0.42
1:A:41:ARG:HH11	1:A:41:ARG:CB	2.34	0.41
1:C:57:LEU:HD21	1:C:134:ILE:HG21	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/529 (98%)	502 (97%)	15 (3%)	0	100	100
1	B	503/529 (95%)	489 (97%)	13 (3%)	1 (0%)	47	58
1	C	518/529 (98%)	501 (97%)	15 (3%)	2 (0%)	34	41
All	All	1538/1587 (97%)	1492 (97%)	43 (3%)	3 (0%)	47	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	202	GLN
1	B	202	GLN
1	C	151	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/410 (98%)	395 (98%)	7 (2%)	60	75
1	B	393/410 (96%)	381 (97%)	12 (3%)	40	55
1	C	403/410 (98%)	389 (96%)	14 (4%)	36	49
All	All	1198/1230 (97%)	1165 (97%)	33 (3%)	43	59

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	ARG
1	A	51	ARG
1	A	206	PHE
1	A	271	ARG
1	A	399	ARG
1	A	463	TRP
1	A	513	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	32	LEU
1	B	35	ASN
1	B	43	VAL
1	B	96	GLN
1	B	206	PHE
1	B	220	ILE
1	B	272	ARG
1	B	275	GLU
1	B	399	ARG
1	B	481	ASP
1	B	489	THR
1	B	505	THR
1	C	32	LEU
1	C	35	ASN
1	C	41	ARG
1	C	51	ARG
1	C	56	ARG
1	C	62	SER
1	C	206	PHE
1	C	358	ASP
1	C	377	ARG
1	C	399	ARG
1	C	461	THR
1	C	463	TRP
1	C	489	THR
1	C	513	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	HIS
1	A	163	ASN
1	A	267	GLN
1	A	298	HIS
1	A	331	HIS
1	A	368	GLN
1	B	35	ASN
1	B	96	GLN
1	B	163	ASN
1	B	164	GLN
1	B	267	GLN
1	B	298	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	415	ASN
1	B	516	HIS
1	C	25	ASN
1	C	35	ASN
1	C	331	HIS
1	C	368	GLN
1	C	415	ASN
1	C	433	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	C	601	-	5,5,5	0.39	0	5,5,5	0.37	0
2	GOL	C	603	-	5,5,5	0.31	0	5,5,5	0.29	0
2	GOL	A	601	-	5,5,5	0.38	0	5,5,5	0.23	0
2	GOL	C	602	-	5,5,5	0.35	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	601	-	-	0/4/4/4	-
2	GOL	C	603	-	-	0/4/4/4	-
2	GOL	A	601	-	-	2/4/4/4	-
2	GOL	C	602	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	C1-C2-C3-O3
2	A	601	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	519/529 (98%)	-0.33	9 (1%) 70 76	14, 23, 43, 56	0
1	B	505/529 (95%)	-0.22	18 (3%) 42 49	17, 27, 52, 84	0
1	C	520/529 (98%)	-0.32	11 (2%) 63 70	16, 26, 45, 69	0
All	All	1544/1587 (97%)	-0.29	38 (2%) 57 64	14, 26, 45, 84	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	471	PHE	5.0
1	B	445	ALA	4.8
1	B	453	GLY	4.2
1	B	475	ILE	3.8
1	B	449	ALA	3.8
1	C	220	ILE	3.6
1	B	451	VAL	3.6
1	A	220	ILE	3.5
1	B	452	ARG	3.5
1	B	450	THR	3.2
1	A	371	ALA	3.2
1	B	473	ALA	3.1
1	A	465	PRO	3.1
1	A	219	GLU	3.0
1	A	466	ASP	2.9
1	C	218	GLY	2.9
1	C	466	ASP	2.8
1	B	448	LEU	2.8
1	C	10	PRO	2.7
1	C	41	ARG	2.6
1	B	292	ARG	2.6
1	B	474	PRO	2.6
1	B	470	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	472	LYS	2.5
1	A	45	ARG	2.5
1	A	272	ARG	2.5
1	B	272	ARG	2.3
1	C	36	GLU	2.3
1	C	37	ARG	2.2
1	C	219	GLU	2.2
1	C	371	ALA	2.2
1	C	465	PRO	2.2
1	C	217	THR	2.2
1	A	37	ARG	2.1
1	B	219	GLU	2.1
1	B	454	GLU	2.0
1	A	457	SER	2.0
1	B	441	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
2	GOL	C	601	6/6	0.61	0.23	61,63,63,63	0
2	GOL	C	603	6/6	0.65	0.22	54,56,56,56	0
2	GOL	C	602	6/6	0.74	0.30	51,51,52,52	0
2	GOL	A	601	6/6	0.79	0.21	49,49,50,50	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.