



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 21, 2020 – 04:55 PM BST

PDB ID : 1Q2E  
Title : CELLOBIOHYDROLASE CEL7A WITH LOOP DELETION 245-252 AND BOUND NON-HYDROLYSABLE CELLOTETRAOSE  
Authors : Stahlberg, J.; Harris, M.; Jones, T.A.  
Deposited on : 2003-07-24  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.6  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	
1	B	426	
2	C	4	
2	D	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	C	3	X	-	-	-
2	GLC	C	4	X	-	-	-
2	GLC	D	3	X	-	-	-
2	GLC	D	4	X	-	-	-
4	CA	B	499	-	-	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7037 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EXOCELLOBIOHYDROLASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	426	3156	1936	521	673	26	0	0	0
1	B	426	3156	1936	521	673	26	0	0	0


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	94	ASP	GLY	cloning artifact	UNP P00725
A	?	-	GLY	deletion	UNP P00725
A	?	-	THR	deletion	UNP P00725
A	?	-	TYR	deletion	UNP P00725
A	?	-	SER	deletion	UNP P00725
A	?	-	ASP	deletion	UNP P00725
A	?	-	ASN	deletion	UNP P00725
A	?	-	ARG	deletion	UNP P00725
A	?	-	TYR	deletion	UNP P00725
B	94	ASP	GLY	cloning artifact	UNP P00725
B	?	-	GLY	deletion	UNP P00725
B	?	-	THR	deletion	UNP P00725
B	?	-	TYR	deletion	UNP P00725
B	?	-	SER	deletion	UNP P00725
B	?	-	ASP	deletion	UNP P00725
B	?	-	ASN	deletion	UNP P00725
B	?	-	ARG	deletion	UNP P00725
B	?	-	TYR	deletion	UNP P00725

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-4-thio-beta-D-glucopyranose-(1-4)-methyl beta-D-glucopyranoside.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	4	Total	C	O	S	0	0	0
			46	25	20	1			
2	D	4	Total	C	O	S	0	0	0
			46	25	20	1			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	14	0
			14	8	1	5		
3	B	1	Total	C	N	O	14	0
			14	8	1	5		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	271	Total	O	0	0
			271	271		
5	B	333	Total	O	0	0
			333	333		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: EXOCELLOBIOHYDROLASE I




- Molecule 1: EXOCELLOBIOHYDROLASE I



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-4-thio-beta-D-glucopyranose-(1-4)-methyl beta-D-glucopyranoside



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-4-thio-beta-D-glucopyranose-(1-4)-methyl beta-D-glucopyranoside



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.34Å 84.31Å 110.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.75 31.25 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.75) 100.0 (31.25-1.75)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.82 (at 1.75Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.226 , 0.239 0.217 , 0.232	Depositor DCC
$R_{free}$ test set	2322 reflections (2.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.6	Xtrriage
Anisotropy	0.518	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7037	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.05 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0487e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA, GLC, MGL, SGC, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/3222	0.67	0/4393
1	B	0.32	0/3222	0.68	0/4393
All	All	0.32	0/6444	0.67	0/8786

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3156	0	2867	22	0
1	B	3156	0	2867	22	0
2	C	46	0	41	0	0
2	D	46	0	41	0	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
4	B	1	0	0	0	0
5	A	271	0	0	4	0
5	B	333	0	0	3	0
All	All	7037	0	5842	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:LEU:HD12	1:B:360:MET:HE1	1.60	0.84
1:B:356:THR:HA	1:B:360:MET:HE2	1.61	0.80
1:A:69:LYS:HE2	1:A:69:LYS:HA	1.62	0.79
1:A:313:GLN:HE21	1:A:314:PRO:HD2	1.49	0.77
1:A:314:PRO:HG2	5:A:1150:HOH:O	1.86	0.75
1:B:314:PRO:HG2	5:B:1248:HOH:O	1.91	0.70
1:B:313:GLN:HE21	1:B:314:PRO:HD2	1.56	0.69
1:A:218:ALA:HB2	1:A:223:GLU:HA	1.78	0.65
1:B:122:LEU:HD12	1:B:360:MET:CE	2.28	0.62
1:A:186:GLN:HG2	5:A:1449:HOH:O	2.05	0.56
1:B:218:ALA:HB2	1:B:223:GLU:HA	1.88	0.55
1:A:209:CYS:O	1:A:210:CYS:HB3	2.09	0.53
1:A:69:LYS:HG2	5:A:1412:HOH:O	2.09	0.53
1:A:213:MET:HB2	1:A:352:PHE:HZ	1.73	0.52
1:B:186:GLN:HG2	5:B:1315:HOH:O	2.09	0.52
1:B:213:MET:HB2	1:B:352:PHE:HZ	1.74	0.51
1:A:257:ASP:OD2	1:A:260:GLY:HA2	2.10	0.51
1:A:227:PRO:HD2	1:A:261:CYS:O	2.12	0.50
1:B:381:TYR:HA	1:B:382:PRO:C	2.31	0.49
1:A:76:ALA:HB1	1:A:82:TYR:CE2	2.48	0.49
1:A:381:TYR:HA	1:A:382:PRO:C	2.33	0.48
1:B:227:PRO:HD2	1:B:261:CYS:O	2.15	0.47
1:B:76:ALA:HB1	1:B:82:TYR:CE2	2.49	0.47
1:B:209:CYS:O	1:B:210:CYS:HB3	2.16	0.46
1:A:35:ASP:OD2	1:A:37:ASN:HB2	2.14	0.46
1:A:347:GLY:HA3	1:A:351:GLN:HB2	1.98	0.46
1:A:34:ILE:HA	1:A:108:LEU:HD23	1.99	0.45
1:B:360:MET:HE2	5:B:1023:HOH:O	2.17	0.45
1:A:137:PRO:HD2	1:A:140:LEU:HD12	1.98	0.45
1:B:347:GLY:HA3	1:B:351:GLN:HB2	1.99	0.45
1:A:334:GLU:HG2	5:A:1556:HOH:O	2.17	0.44
1:B:257:ASP:OD2	1:B:260:GLY:HA2	2.18	0.44
1:A:315:ASN:ND2	1:A:324:ASN:HD22	2.16	0.44
1:B:34:ILE:HA	1:B:108:LEU:HD23	2.00	0.43
1:B:95:PHE:CZ	1:B:140:LEU:HD11	2.54	0.43
1:B:35:ASP:OD2	1:B:37:ASN:HB2	2.18	0.43
1:B:95:PHE:CE1	1:B:140:LEU:HD11	2.54	0.42
1:A:206:HIS:CD2	1:A:237:ILE:HD11	2.54	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:TYR:HA	1:B:159:PRO:HD2	1.86	0.42
1:A:315:ASN:HD21	1:A:324:ASN:HD22	1.67	0.42
1:A:158:TYR:HA	1:A:159:PRO:HD2	1.85	0.41
1:A:95:PHE:CZ	1:A:140:LEU:HD21	2.56	0.41
1:B:99:SER:OG	1:B:100:ALA:N	2.54	0.41
1:B:318:LEU:HD21	1:B:334:GLU:HA	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/426 (100%)	414 (98%)	10 (2%)	0	100	100
1	B	424/426 (100%)	418 (99%)	6 (1%)	0	100	100
All	All	848/852 (100%)	832 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/349 (100%)	347 (99%)	2 (1%)	86	79
1	B	349/349 (100%)	345 (99%)	4 (1%)	73	60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	698/698 (100%)	692 (99%)	6 (1%)	78 67

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	223	GLU
1	B	37	ASN
1	B	223	GLU
1	B	241	ASP
1	B	307	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	GLN
1	A	186	GLN
1	A	307	ASN
1	A	313	GLN
1	A	315	ASN
1	A	373	ASN
1	A	413	ASN
1	B	98	GLN
1	B	101	GLN
1	B	186	GLN
1	B	307	ASN
1	B	313	GLN
1	B	373	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PCA	A	1	1	7,8,9	2.22	1 (14%)	9,10,12	1.37	2 (22%)
1	PCA	B	1	1	7,8,9	2.24	1 (14%)	9,10,12	1.33	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
1	PCA	B	1	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	PCA	CD-N	5.37	1.48	1.34
1	A	1	PCA	CD-N	5.32	1.48	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CA-N-CD	-2.39	105.41	113.58
1	A	1	PCA	CB-CA-C	-2.37	109.44	112.70
1	B	1	PCA	CB-CA-C	-2.36	109.46	112.70
1	B	1	PCA	CA-N-CD	-2.26	105.85	113.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MGL	C	1	2	13,13,13	0.93	1 (7%)	18,18,18	0.80	1 (5%)
2	SGC	C	2	2	10,11,12	0.63	0	12,15,17	0.91	0
2	GLC	C	3	2	11,11,12	0.52	0	15,15,17	0.54	0
2	GLC	C	4	2	11,11,12	0.56	0	15,15,17	0.47	0
2	MGL	D	1	2	13,13,13	0.92	1 (7%)	18,18,18	0.75	1 (5%)
2	SGC	D	2	2	10,11,12	0.54	0	12,15,17	1.06	0
2	GLC	D	3	2	11,11,12	0.53	0	15,15,17	0.57	0
2	GLC	D	4	2	11,11,12	0.64	0	15,15,17	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MGL	C	1	2	-	0/4/24/24	0/1/1/1
2	SGC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	3	2	1/1/4/5	0/2/19/22	0/1/1/1
2	GLC	C	4	2	1/1/4/5	0/2/19/22	0/1/1/1
2	MGL	D	1	2	-	2/4/24/24	0/1/1/1
2	SGC	D	2	2	-	2/2/19/22	0/1/1/1
2	GLC	D	3	2	1/1/4/5	0/2/19/22	0/1/1/1
2	GLC	D	4	2	1/1/4/5	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	MGL	O1-C1	2.72	1.44	1.40
2	D	1	MGL	O1-C1	2.70	1.44	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	C	1	MGL	O1-C1-C2	2.83	111.46	108.15
2	D	1	MGL	O1-C1-C2	2.60	111.19	108.15

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	D	4	GLC	C1
2	C	3	GLC	C1
2	D	3	GLC	C1
2	C	4	GLC	C1


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1	MGL	C4-C5-C6-O6
2	D	1	MGL	O5-C5-C6-O6
2	D	2	SGC	C4-C5-C6-O6
2	D	2	SGC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	B	436	1	14,14,15	0.59	0	17,19,21	0.94	1 (5%)
3	NAG	A	435	1	14,14,15	0.46	0	17,19,21	0.83	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	436	1	-	0/6/23/26	0/1/1/1
3	NAG	A	435	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	436	NAG	C2-N2-C7	-2.47	119.38	122.90
3	A	435	NAG	C2-N2-C7	-2.01	120.05	122.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/426 (99%)	0.54	19 (4%) 33 39	7, 11, 18, 26	0
1	B	425/426 (99%)	0.43	9 (2%) 63 71	5, 8, 16, 21	0
All	All	850/852 (99%)	0.48	28 (3%) 46 53	5, 10, 17, 26	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	100	ALA	6.5
1	A	101	GLN	5.0
1	A	387	SER	4.6
1	A	100	ALA	4.6
1	A	388	SER	4.0
1	A	45	ASN	3.5
1	B	45	ASN	3.4
1	B	101	GLN	3.0
1	A	384	ASN	3.0
1	B	434	GLY	2.7
1	A	371	TYR	2.7
1	A	195	SER	2.7
1	A	5	THR	2.6
1	A	114	ASP	2.6
1	A	196	SER	2.6
1	A	65	GLU	2.4
1	B	195	SER	2.4
1	A	193	GLU	2.3
1	B	193	GLU	2.3
1	B	5	THR	2.3
1	A	244	GLY	2.3
1	A	386	THR	2.3
1	A	241	ASP	2.3
1	A	215	ILE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	434	GLY	2.2
1	A	99	SER	2.1
1	B	388	SER	2.1
1	B	196	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PCA	A	1	8/9	0.94	0.11	12,13,13,13	0
1	PCA	B	1	8/9	0.97	0.08	9,9,9,9	0

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MGL	D	1	13/13	0.79	0.25	13,15,15,16	0
2	MGL	C	1	13/13	0.86	0.22	12,14,14,14	0
2	GLC	C	3	11/12	0.93	0.11	9,10,12,12	0
2	SGC	D	2	11/12	0.93	0.13	8,11,12,13	0
2	GLC	C	4	11/12	0.93	0.12	8,10,11,11	0
2	GLC	D	4	11/12	0.93	0.11	7,8,9,10	0
2	GLC	D	3	11/12	0.94	0.11	9,10,11,11	0
2	SGC	C	2	11/12	0.95	0.12	8,12,13,13	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	B	436	14/15	-	-	10,10,10,10	14
3	NAG	A	435	14/15	-	-	11,11,11,11	14
4	CA	B	499	1/1	0.35	1.75	54,54,54,54	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.