wwPDB NMR Structure Validation Summary Report (i)

Feb 19, 2022 – 01:19 PM EST

PDB ID	:	1Q60
Title	:	Solution Structure of RSGI RUH-004, a GTF2I domain in Mouse cDNA
Authors	:	Doi-Katayama, Y.; Hayashi, F.; Hirota, H.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2003-08-12

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	99	46% 37% • 14%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:7-A:91 (85)	0.27	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 13, 14, 15, 19
2	18, 20
3	12, 16
Single-model clusters	11; 17

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1558 atoms, of which 790 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called General transcription factor II-I.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	99	1558	497	790	128	141	2	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP Q9ESZ8
A	2	SER	-	cloning artifact	UNP Q9ESZ8
A	3	SER	-	cloning artifact	UNP Q9ESZ8
A	4	GLY	-	cloning artifact	UNP Q9ESZ8
A	5	SER	-	cloning artifact	UNP Q9ESZ8
A	6	SER	-	cloning artifact	UNP Q9ESZ8
A	7	GLY	-	cloning artifact	UNP Q9ESZ8
A	23	ALA	THR	SEE REMARK 999	UNP Q9ESZ8
A	94	SER	-	cloning artifact	UNP Q9ESZ8
A	95	GLY	-	cloning artifact	UNP Q9ESZ8
A	96	PRO	-	cloning artifact	UNP Q9ESZ8
A	97	SER	-	cloning artifact	UNP Q9ESZ8
A	98	SER	-	cloning artifact	UNP Q9ESZ8
A	99	GLY	-	cloning artifact	UNP Q9ESZ8

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: General transcription factor II-I

Chain A: 46% 37% • 14%

G1 S2 S3 G4 S5 S6 G7 L8 K9 Q10 K11 V12 E13 F16 N17 E18 K19 C20 G21 E22 V30 K31 V32 P33 L36 F37 E38 S39 D43 F44 Y45 V46 E47 E51 G52 V53 P54 F55 R56 T60 F61 G62 I63 P64 R65 I69 L70 K73 K77 F78 I79

K82 F86 E87 E92 S93 S94 G95 P96 S97 S98 G99

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 6. Colouring as in section 4.1 above.

• Molecule 1: General transcription factor II-I

Chain A: 41% 40% • • 14%

G1 S2 S3 G4 S5 S6 G7 L8 K9 Q10 K11 V12 E13 E18 K19 C20 G21 E22 A23 L24 V30 K31 V32 P33 F34 A35 L36 F37 E38 S39 F40 P41 E42 D43 F44 Y45 V46 E47 E51 G52 V53 P54 F55 T60 F61 G62 I63 P64 R65 L66 I69 L70 K77 F78

I79 I80 K81 K82 F86 E87 K91 E92 S93 S94 G95 P96 S97 S98 G99

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.7
CYANA	refinement	1.0.7

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	688	725	725	32±6
All	All	13760	14500	14500	642

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

5 of 188 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:36:LEU:HD23	1:A:44:PHE:CE2	1.08	1.84	6	7
1:A:20:CYS:SG	1:A:32:VAL:HG22	0.90	2.06	11	3
1:A:46:VAL:HG21	1:A:55:PHE:CE1	0.88	2.04	18	11
1:A:45:TYR:OH	1:A:79:ILE:HG21	0.83	1.72	7	18
1:A:45:TYR:CZ	1:A:79:ILE:HG21	0.83	2.09	11	18

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	85/99 (86%)	65±3 (76±4%)	17±4 (20±4%)	3±1 (3±2%)	6 35
All	All	1700/1980 (86%)	1296 (76%)	345 (20%)	59 (3%)	6 35

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	60	THR	16
1	A	43	ASP	10
1	A	61	PHE	10
1	A	31	LYS	5
1	A	64	PRO	5

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	74/84 (88%)	60±2 (81±3%)	14±2 (19±3%)	4 36
All	All	1480/1680 (88%)	1199 (81%)	281 (19%)	4 36

5 of 39 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	60	THR	20
1	A	18	GLU	19
1	A	22	GLU	19
1	A	87	GLU	17
1	A	10	GLN	15

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided