



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 02:14 am BST

PDB ID : 3Q6P  
Title : Salivary protein from *Lutzomyia longipalpis*. Selenomethionine derivative  
Authors : Andersen, J.F.; Xu, X.; Chang, B.W.; Collin, N.; Valenzuela, J.G.; Ribeiro, J.M.  
Deposited on : 2011-01-03  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 43.2 kDa salivary protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	381	Total 3056	C 1963	N 520	O 562	S 4	Se 7	0	0	0
1	B	381	Total 3056	C 1963	N 520	O 562	S 4	Se 7	0	0	0

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	B	1	Total 13	C 6	O 7	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	A	33	Total 33	O 33	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	32	Total	O	0	0
			32	32		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 43.2 kDa salivary protein



- Molecule 1: 43.2 kDa salivary protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.36Å 120.36Å 248.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.75 39.96 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.3 (40.00-2.75) 99.4 (39.96-2.75)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.58 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.197 , 0.261 0.196 , 0.261	Depositor DCC
$R_{free}$ test set	1425 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.1	Xtrriage
Anisotropy	0.078	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 31.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6190	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/3132	0.66	1/4227 (0.0%)
1	B	0.58	0/3132	0.71	3/4227 (0.1%)
All	All	0.57	0/6264	0.69	4/8454 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	46	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	B	46	ARG	NE-CZ-NH2	-5.93	117.34	120.30
1	A	268	LEU	N-CA-C	5.78	126.60	111.00
1	B	50	ARG	N-CA-C	-5.24	96.86	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	113	ARG	Peptide
1	B	49	PRO	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3056	0	3014	30	0
1	B	3056	0	3014	31	0
2	B	13	0	5	0	0
3	A	33	0	0	0	0
3	B	32	0	0	0	0
All	All	6190	0	6033	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:ALA:CA	1:B:340:MSE:HE1	2.21	0.70
1:B:90:TYR:OH	1:B:109:GLU:OE2	2.12	0.67
1:A:6:TYR:OH	1:A:63:SER:HB3	1.96	0.64
1:A:200:THR:HG21	1:A:240:TYR:OH	1.97	0.64
1:B:7:LYS:HE2	1:B:361:ARG:NH2	2.12	0.64
1:B:30:ALA:HA	1:B:340:MSE:HE1	1.79	0.63
1:A:271:ARG:NH2	1:A:275:ASN:O	2.33	0.60
1:B:268:LEU:N	1:B:269:GLY:HA2	2.17	0.60
1:A:268:LEU:H	1:A:269:GLY:HA2	1.68	0.58
1:B:198:HIS:ND1	1:B:200:THR:HB	2.17	0.58
1:A:379:MSE:HE3	1:A:379:MSE:HA	1.86	0.58
1:A:36:GLU:HG3	1:A:98:ARG:HD2	1.83	0.58
1:A:268:LEU:N	1:A:269:GLY:HA2	2.18	0.58
1:B:97:CYS:SG	1:B:168:CYS:SG	3.00	0.57
1:B:46:ARG:NH2	1:B:86:LEU:O	2.38	0.57
1:A:306:LYS:HB3	1:A:313:ASN:OD1	2.05	0.56
1:B:268:LEU:H	1:B:269:GLY:HA2	1.69	0.56
1:B:297:LYS:HD3	1:B:321:SER:HA	1.88	0.55
1:B:46:ARG:HG2	1:B:51:VAL:HB	1.88	0.55
1:A:268:LEU:O	1:A:309:LEU:HB3	2.07	0.54
1:B:223:PHE:H	1:B:224:GLY:HA3	1.73	0.54
1:A:241:LEU:O	1:A:241:LEU:HD12	2.08	0.53

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:ALA:HA	1:B:340:MSE:CE	2.38	0.53
1:B:30:ALA:C	1:B:340:MSE:HE1	2.29	0.53
1:A:49:PRO:HG2	1:A:113:ARG:CZ	2.39	0.53
1:A:144:PRO:HG2	1:A:147:LEU:HD12	1.91	0.51
1:B:308:PRO:O	1:B:313:ASN:ND2	2.43	0.51
1:B:113:ARG:N	1:B:114:GLY:HA2	2.26	0.51
1:A:365:ILE:HG21	1:A:371:ALA:HB2	1.94	0.50
1:A:97:CYS:CB	1:A:168:CYS:HG	2.25	0.49
1:A:200:THR:CG2	1:A:240:TYR:OH	2.60	0.48
1:B:31:PHE:HD2	1:B:340:MSE:HE2	1.78	0.48
1:A:97:CYS:HB3	1:A:168:CYS:HG	1.78	0.48
1:B:167:ASP:O	1:B:168:CYS:HB2	2.14	0.47
1:A:372:ILE:HD12	1:A:378:ASP:HA	1.97	0.47
1:A:247:LYS:HA	1:A:269:GLY:O	2.14	0.47
1:A:223:PHE:O	1:A:223:PHE:HD1	1.98	0.46
1:A:247:LYS:HG2	1:A:267:LEU:HD11	1.97	0.46
1:B:205:ARG:O	1:B:244:SER:HB3	2.15	0.46
1:A:46:ARG:HG2	1:A:51:VAL:HB	1.98	0.46
1:B:7:LYS:HE3	1:B:363:MSE:HE2	1.97	0.45
1:A:205:ARG:NH2	1:A:276:ASP:OD1	2.49	0.45
1:A:161:VAL:O	1:A:230:ARG:NH1	2.49	0.45
1:A:205:ARG:HH12	1:A:246:ILE:HG23	1.81	0.44
1:B:200:THR:HG21	1:B:240:TYR:OH	2.17	0.44
1:B:247:LYS:HA	1:B:269:GLY:O	2.18	0.43
1:B:127:ALA:HB3	1:B:139:VAL:HB	2.01	0.43
1:A:112:SER:OG	1:A:113:ARG:N	2.51	0.43
1:B:311:MSE:HB3	1:B:311:MSE:HE2	1.99	0.43
1:B:331:VAL:HA	1:B:336:GLY:O	2.18	0.43
1:B:161:VAL:O	1:B:230:ARG:NH1	2.52	0.42
1:A:349:LYS:HA	1:A:349:LYS:HD3	1.78	0.42
1:B:224:GLY:O	1:B:240:TYR:HB2	2.19	0.42
1:A:96:ASP:N	1:A:96:ASP:OD1	2.51	0.41
1:B:53:TYR:CE1	1:B:76:LYS:HD3	2.56	0.41
1:B:348:ARG:O	1:B:349:LYS:HB2	2.20	0.41
1:A:322:ARG:O	1:A:361:ARG:HD3	2.20	0.41
1:A:35:ALA:HB1	1:A:334:LYS:HG3	2.03	0.41
1:A:137:GLU:CD	1:A:140:ARG:HE	2.24	0.41
1:B:184:PHE:CZ	1:B:193:SER:HB2	2.56	0.41
1:B:200:THR:HG22	1:B:240:TYR:HE1	1.85	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/381 (100%)	351 (93%)	25 (7%)	3 (1%)	19	34
1	B	379/381 (100%)	358 (94%)	18 (5%)	3 (1%)	19	34
All	All	758/762 (100%)	709 (94%)	43 (6%)	6 (1%)	19	34

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	ALA
1	A	325	PHE
1	B	64	LEU
1	B	224	GLY
1	A	269	GLY
1	B	269	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/319 (102%)	310 (95%)	16 (5%)	25	43
1	B	326/319 (102%)	305 (94%)	21 (6%)	17	31
All	All	652/638 (102%)	615 (94%)	37 (6%)	20	36

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	VAL
1	A	63	SER
1	A	71	SER
1	A	96	ASP
1	A	120	SER
1	A	152	THR
1	A	189	LYS
1	A	195	ASN
1	A	200	THR
1	A	205	ARG
1	A	223	PHE
1	A	232	SER
1	A	241	LEU
1	A	259	LYS
1	A	276	ASP
1	A	363	MSE
1	B	49	PRO
1	B	50	ARG
1	B	58	VAL
1	B	64	LEU
1	B	67	LYS
1	B	97	CYS
1	B	113	ARG
1	B	120	SER
1	B	134	ASN
1	B	152	THR
1	B	180	ARG
1	B	183	LEU
1	B	223	PHE
1	B	232	SER
1	B	241	LEU
1	B	268	LEU
1	B	276	ASP
1	B	312	LYS
1	B	363	MSE
1	B	367	ASP
1	B	380	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CIT	B	382	-	3,12,12	1.31	0	3,17,17	2.51	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	382	-	-	0/6/16/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	382	CIT	C3-C4-C5	-3.65	109.14	114.98
2	B	382	CIT	C3-C2-C1	-2.27	111.35	114.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/381 (98%)	-0.05	10 (2%) 54 63	27, 50, 76, 90	0
1	B	374/381 (98%)	-0.25	4 (1%) 80 86	31, 45, 64, 87	0
All	All	748/762 (98%)	-0.15	14 (1%) 66 75	27, 47, 71, 90	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	146	ARG	5.0
1	A	147	LEU	4.2
1	A	1	ALA	4.0
1	B	381	ALA	3.9
1	A	180	ARG	3.5
1	B	380	ASN	3.2
1	A	148	VAL	2.8
1	A	97	CYS	2.6
1	A	62	ASN	2.4
1	A	170	GLU	2.3
1	A	367	ASP	2.2
1	B	260	GLY	2.1
1	A	2	ASP	2.1
1	B	1	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CIT	B	382	13/13	0.80	0.23	80,81,85,86	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.