



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 9, 2022 – 03:29 pm GMT

PDB ID : 7Q6P  
Title : Crystal Structure of bacterial Prolyl Peptidyl Isomerase with 5,5'-difluoroleucines  
Authors : Tars, K.; Jaudzems, K.; Recacha, R.  
Deposited on : 2021-11-09  
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	90% (green), 9% (yellow), 1% (orange), 0% (red), 0% (grey)
1	B	170	92% (green), 7% (yellow), 1% (orange), 0% (red), 0% (grey)
1	C	170	3% (red), 70% (green), 15% (yellow), 1% (orange), 14% (grey)
1	D	170	4% (red), 76% (green), 15% (yellow), 1% (orange), 8% (grey)

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5549 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidyl-prolyl cis-trans isomerase B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	F	N	O	S			
1	A	169	1335	833	10	234	250	8	0	0	0
1	B	168	1330	830	10	231	251	8	0	1	0
1	C	147	1153	722	4	205	214	8	0	0	0
1	D	157	1227	768	8	210	233	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	165	HIS	-	expression tag	UNP P23869
A	166	HIS	-	expression tag	UNP P23869
A	167	HIS	-	expression tag	UNP P23869
A	168	HIS	-	expression tag	UNP P23869
A	169	HIS	-	expression tag	UNP P23869
A	170	HIS	-	expression tag	UNP P23869
B	165	HIS	-	expression tag	UNP P23869
B	166	HIS	-	expression tag	UNP P23869
B	167	HIS	-	expression tag	UNP P23869
B	168	HIS	-	expression tag	UNP P23869
B	169	HIS	-	expression tag	UNP P23869
B	170	HIS	-	expression tag	UNP P23869
C	165	HIS	-	expression tag	UNP P23869
C	166	HIS	-	expression tag	UNP P23869
C	167	HIS	-	expression tag	UNP P23869
C	168	HIS	-	expression tag	UNP P23869
C	169	HIS	-	expression tag	UNP P23869
C	170	HIS	-	expression tag	UNP P23869
D	165	HIS	-	expression tag	UNP P23869
D	166	HIS	-	expression tag	UNP P23869
D	167	HIS	-	expression tag	UNP P23869

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	168	HIS	-	expression tag	UNP P23869
D	169	HIS	-	expression tag	UNP P23869
D	170	HIS	-	expression tag	UNP P23869

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	171	Total O 171 171	0	0
3	B	161	Total O 161 161	0	0
3	C	79	Total O 79 79	0	0
3	D	78	Total O 78 78	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peptidyl-prolyl cis-trans isomerase B

Chain A: 




- Molecule 1: Peptidyl-prolyl cis-trans isomerase B

Chain B: 





- Molecule 1: Peptidyl-prolyl cis-trans isomerase B

Chain C: 



- Molecule 1: Peptidyl-prolyl cis-trans isomerase B

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.06Å 83.46Å 122.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.40 – 1.82 57.40 – 1.82	Depositor EDS
% Data completeness (in resolution range)	98.5 (57.40-1.82) 98.5 (57.40-1.82)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.42 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.239 , 0.279 0.239 , 0.278	Depositor DCC
$R_{free}$ test set	3596 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.3	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5549	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FFL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/1311	0.75	2/1761 (0.1%)
1	B	0.62	0/1308	0.72	0/1757
1	C	0.53	0/1155	0.68	0/1552
1	D	0.53	0/1206	0.66	0/1618
All	All	0.58	0/4980	0.71	2/6688 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	MET	CG-SD-CE	5.93	109.69	100.20
1	A	10	ASP	CB-CG-OD2	-5.15	113.66	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1335	0	1214	8	0
1	B	1330	0	1211	7	0
1	C	1153	0	1084	21	0
1	D	1227	0	1141	19	0
2	A	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	171	0	0	1	0
3	B	161	0	0	3	0
3	C	79	0	0	5	0
3	D	78	0	0	4	0
All	All	5549	0	4650	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:HIS:ND1	1:A:135:LYS:HE2	1.87	0.90
1:C:80:ARG:NH2	1:C:134:ASP:OD2	2.17	0.76
1:C:95:THR:HA	3:C:305:HOH:O	1.96	0.65
1:C:69:LYS:HA	1:C:92:HIS:HB3	1.79	0.65
1:C:79:THR:O	1:C:82:THR:HG22	1.98	0.64
1:C:84:ALA:HB3	1:C:99:PHE:CZ	2.34	0.62
1:C:67:PRO:HA	3:C:305:HOH:O	2.02	0.60
1:A:125:GLU:HB3	1:D:1:MET:HE1	1.84	0.58
1:D:72:ALA:HB1	1:D:110:PHE:HB2	1.85	0.58
1:C:80:ARG:HH22	1:C:134:ASP:CG	2.05	0.58
1:C:69:LYS:HD3	1:C:92:HIS:HB3	1.86	0.58
1:C:70:ASN:OD1	1:C:121:CYS:HB2	2.04	0.58
1:C:78:ASN:HA	1:C:82:THR:HG21	1.87	0.57
1:D:70:ASN:HB2	1:D:91:PRO:O	2.03	0.57
1:C:85:MET:O	1:C:99:PHE:HE1	1.88	0.57
1:D:13:ILE:CG2	1:D:124:ALA:HB1	2.35	0.56
1:B:53:GLY:N	3:B:303:HOH:O	2.29	0.55
1:B:143:ARG:NH2	3:B:304:HOH:O	2.38	0.54
1:D:19:LYS:HE2	1:D:76:FFL:F2	1.98	0.54
1:C:131:ASP:O	1:C:135:LYS:HG2	2.09	0.52
1:C:90:ALA:O	1:C:93:SER:OG	2.18	0.51
1:C:16:PHE:HB3	1:C:19:LYS:HG2	1.94	0.49
1:D:67:PRO:HB2	1:D:92:HIS:O	2.12	0.49
1:D:102:VAL:HG22	3:D:201:HOH:O	2.14	0.48
1:A:46:ASN:OD1	1:A:137:LYS:HE2	2.14	0.47
1:D:87:ARG:HG3	1:D:88:THR:O	2.14	0.46
1:D:19:LYS:HB3	1:D:19:LYS:HE3	1.58	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:19:LYS:HE3	1:D:74:ASN:HB2	1.98	0.46
1:C:167:HIS:HE1	3:C:319:HOH:O	1.99	0.45
1:C:17:ASP:HB3	3:C:314:HOH:O	2.16	0.45
1:A:84:ALA:HB3	1:A:99:PHE:CZ	2.52	0.45
1:D:102:VAL:O	3:D:201:HOH:O	2.21	0.44
1:B:116:GLN:HA	1:B:116:GLN:OE1	2.18	0.44
1:B:60:LYS:HE2	1:B:60:LYS:HB2	1.85	0.43
1:D:134:ASP:O	1:D:137:LYS:HG2	2.19	0.43
1:A:66:GLU:HG3	3:A:378:HOH:O	2.18	0.42
1:C:68:ILE:HD12	1:C:69:LYS:H	1.84	0.42
1:C:26:ASN:ND2	1:C:95:THR:O	2.50	0.42
1:C:69:LYS:HD3	1:C:92:HIS:CB	2.49	0.42
1:A:159:SER:CB	1:B:37:ASN:HD22	2.33	0.42
1:D:45:ILE:HG22	1:D:48:PHE:HB3	2.02	0.42
1:B:168:HIS:HD1	1:D:158:GLU:CD	2.22	0.41
1:D:19:LYS:CE	1:D:74:ASN:HB2	2.50	0.41
1:D:87:ARG:HD3	3:D:261:HOH:O	2.21	0.41
1:A:1:MET:HB2	1:A:163:SER:HB3	2.03	0.41
1:C:90:ALA:HB1	3:C:317:HOH:O	2.20	0.41
1:A:125:GLU:HB3	1:D:1:MET:CE	2.48	0.41
1:B:66:GLU:HB3	3:B:394:HOH:O	2.21	0.41
1:D:14:LYS:NZ	3:D:216:HOH:O	2.54	0.41
1:C:140:ALA:O	1:C:151:PRO:HA	2.21	0.41
1:D:49:MET:SD	1:D:51:GLN:HB2	2.62	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	162/170 (95%)	153 (94%)	9 (6%)	0	<b>100</b> <b>100</b>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	162/170 (95%)	156 (96%)	6 (4%)	0	100	100
1	C	137/170 (81%)	127 (93%)	9 (7%)	1 (1%)	22	10
1	D	149/170 (88%)	138 (93%)	11 (7%)	0	100	100
All	All	610/680 (90%)	574 (94%)	35 (6%)	1 (0%)	47	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	80	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/139 (99%)	136 (99%)	2 (1%)	67	58
1	B	138/139 (99%)	135 (98%)	3 (2%)	52	39
1	C	123/139 (88%)	119 (97%)	4 (3%)	38	23
1	D	128/139 (92%)	125 (98%)	3 (2%)	50	37
All	All	527/556 (95%)	515 (98%)	12 (2%)	50	37

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ASP
1	A	49	MET
1	B	17	ASP
1	B	49	MET
1	B	106	ASP
1	C	29	ASP
1	C	49	MET
1	C	85	MET
1	C	95	THR
1	D	17	ASP

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	49	MET
1	D	60	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	37	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

16 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	FFL	D	28	1	5,9,10	0.61	0	5,10,12	1.12	1 (20%)
1	FFL	B	83	1	5,9,10	0.79	0	5,10,12	3.25	3 (60%)
1	FFL	D	76	1	5,9,10	0.80	0	5,10,12	0.54	0
1	FFL	A	76	1	5,9,10	0.42	0	5,10,12	0.66	0
1	FFL	A	115	1	5,9,10	0.75	0	5,10,12	0.82	0
1	FFL	C	28	1	5,9,10	0.61	0	5,10,12	0.87	0
1	FFL	A	83	1	5,9,10	0.71	0	5,10,12	0.25	0
1	FFL	B	76	1	5,9,10	0.50	0	5,10,12	0.35	0
1	FFL	B	115	1	5,9,10	0.72	0	5,10,12	0.23	0
1	FFL	D	83	1	5,9,10	0.63	0	5,10,12	1.97	1 (20%)
1	FFL	B	108	1	5,9,10	0.62	0	5,10,12	1.24	1 (20%)
1	FFL	D	108	1	5,9,10	0.67	0	5,10,12	0.70	0
1	FFL	B	28	1	5,9,10	0.74	0	5,10,12	0.45	0
1	FFL	A	28	1	5,9,10	0.61	0	5,10,12	0.88	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	FFL	A	108	1	5,9,10	0.85	0	5,10,12	1.50	1 (20%)
1	FFL	C	83	1	5,9,10	0.81	0	5,10,12	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FFL	D	28	1	-	1/9/10/12	-
1	FFL	B	83	1	-	4/9/10/12	-
1	FFL	D	76	1	-	3/9/10/12	-
1	FFL	A	76	1	-	1/9/10/12	-
1	FFL	A	115	1	-	6/9/10/12	-
1	FFL	C	28	1	-	1/9/10/12	-
1	FFL	A	83	1	-	2/9/10/12	-
1	FFL	B	76	1	-	1/9/10/12	-
1	FFL	B	115	1	-	2/9/10/12	-
1	FFL	D	83	1	-	4/9/10/12	-
1	FFL	B	108	1	-	0/9/10/12	-
1	FFL	D	108	1	-	4/9/10/12	-
1	FFL	B	28	1	-	0/9/10/12	-
1	FFL	A	28	1	-	0/9/10/12	-
1	FFL	A	108	1	-	0/9/10/12	-
1	FFL	C	83	1	-	2/9/10/12	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	83	FFL	CG-CB-CA	6.27	122.95	114.52
1	D	83	FFL	CG-CB-CA	4.03	119.94	114.52
1	A	108	FFL	CG-CB-CA	3.08	118.66	114.52
1	B	83	FFL	CB-CG-CD1	2.31	117.06	111.18
1	D	28	FFL	CG-CB-CA	2.14	117.39	114.52
1	B	108	FFL	CG-CB-CA	2.07	117.30	114.52
1	B	83	FFL	CD1-CG-CD2	2.02	114.62	110.95

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	115	FFL	F1-CD1-CG-CB
1	A	115	FFL	F1-CD1-CG-CD2
1	A	115	FFL	F2-CD2-CG-CB
1	A	115	FFL	F2-CD2-CG-CD1
1	B	83	FFL	CA-CB-CG-CD1
1	B	83	FFL	CA-CB-CG-CD2
1	B	83	FFL	F2-CD2-CG-CB
1	B	83	FFL	F2-CD2-CG-CD1
1	D	76	FFL	CA-CB-CG-CD2
1	D	76	FFL	F2-CD2-CG-CB
1	D	76	FFL	F2-CD2-CG-CD1
1	D	83	FFL	CA-CB-CG-CD1
1	D	83	FFL	CA-CB-CG-CD2
1	D	83	FFL	F1-CD1-CG-CD2
1	D	83	FFL	F2-CD2-CG-CB
1	D	108	FFL	CA-CB-CG-CD1
1	D	108	FFL	CA-CB-CG-CD2
1	B	76	FFL	CA-CB-CG-CD2
1	D	28	FFL	CA-CB-CG-CD2
1	C	28	FFL	C-CA-CB-CG
1	C	83	FFL	C-CA-CB-CG
1	D	108	FFL	C-CA-CB-CG
1	C	83	FFL	N-CA-CB-CG
1	A	76	FFL	CA-CB-CG-CD2
1	A	83	FFL	C-CA-CB-CG
1	A	115	FFL	C-CA-CB-CG
1	B	115	FFL	C-CA-CB-CG
1	A	83	FFL	N-CA-CB-CG
1	A	115	FFL	N-CA-CB-CG
1	B	115	FFL	N-CA-CB-CG
1	D	108	FFL	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	76	FFL	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	201	-	4,4,4	0.20	0	6,6,6	0.32	0
2	SO4	C	201	-	4,4,4	0.21	0	6,6,6	0.60	0
2	SO4	A	201	-	4,4,4	0.23	0	6,6,6	0.27	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	164/170 (96%)	-0.46	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	10, 17, 32, 40	0
1	B	163/170 (95%)	-0.46	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	11, 17, 34, 41	0
1	C	145/170 (85%)	0.18	5 (3%) <span style="border: 1px solid red; padding: 2px;">45</span> <span style="border: 1px solid red; padding: 2px;">39</span>	16, 29, 46, 56	0
1	D	153/170 (90%)	0.42	7 (4%) <span style="border: 1px solid red; padding: 2px;">32</span> <span style="border: 1px solid red; padding: 2px;">27</span>	16, 33, 48, 66	0
All	All	625/680 (91%)	-0.10	12 (1%) <span style="border: 1px solid blue; padding: 2px;">66</span> <span style="border: 1px solid blue; padding: 2px;">63</span>	10, 23, 42, 66	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	27	PHE	3.6
1	D	90	ALA	3.6
1	C	122	VAL	3.0
1	D	107	PHE	3.0
1	C	84	ALA	2.9
1	D	84	ALA	2.8
1	D	68	ILE	2.7
1	C	72	ALA	2.4
1	C	68	ILE	2.4
1	C	90	ALA	2.3
1	D	122	VAL	2.2
1	D	66	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	FFL	D	76	10/11	0.67	0.19	49,52,55,56	0
1	FFL	C	28	10/11	0.74	0.19	32,40,42,42	0
1	FFL	B	115	10/11	0.74	0.22	36,38,48,49	0
1	FFL	D	28	10/11	0.80	0.23	37,41,44,50	0
1	FFL	D	108	10/11	0.84	0.16	43,46,50,55	0
1	FFL	D	83	10/11	0.88	0.11	25,27,31,33	0
1	FFL	C	83	10/11	0.89	0.10	24,26,40,40	0
1	FFL	A	28	10/11	0.90	0.12	12,14,30,33	0
1	FFL	A	115	10/11	0.93	0.12	18,24,31,32	0
1	FFL	B	76	10/11	0.94	0.10	19,22,24,24	0
1	FFL	B	83	10/11	0.94	0.10	10,12,19,19	0
1	FFL	B	28	10/11	0.94	0.11	14,16,30,34	0
1	FFL	A	83	10/11	0.95	0.09	10,11,14,18	0
1	FFL	B	108	10/11	0.96	0.07	14,17,19,19	0
1	FFL	A	108	10/11	0.96	0.09	13,16,21,24	0
1	FFL	A	76	10/11	0.97	0.07	14,16,23,23	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	201	5/5	0.96	0.08	50,51,56,61	0
2	SO4	A	201	5/5	0.99	0.05	23,24,28,29	0
2	SO4	C	201	5/5	0.99	0.06	23,27,29,30	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.