



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2023 – 09:09 pm GMT

PDB ID : 8QB3
Title : ADDobody zinc containing condition
Authors : Buzas, D.; Toelzer, C.; Berger, I.; Schaffitzel, C.
Deposited on : 2023-08-24
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	
1	B	310	
1	C	310	
1	D	310	
1	E	310	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 8617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADDobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	233	1810	1167	303	334	6	0	0	0
1	B	233	1806	1164	307	329	6	0	0	0
1	C	216	1653	1075	278	294	6	0	0	0
1	D	206	1525	984	256	280	5	0	0	0
1	E	233	1822	1176	306	333	7	0	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	101.48Å 103.07Å 167.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	87.74 – 2.90 87.74 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (87.74-2.90) 89.8 (87.74-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 2.91Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.271 , 0.291 0.273 , 0.286	Depositor DCC
R_{free} test set	2000 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtrriage
Anisotropy	0.898	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 8.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.039 for k,h,-l	Xtrriage
F_o, F_c correlation	0.75	EDS
Total number of atoms	8617	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1856	0.49	0/2529
1	B	0.26	0/1849	0.49	0/2515
1	C	0.26	0/1696	0.50	0/2313
1	D	0.25	0/1560	0.48	0/2131
1	E	0.28	0/1868	0.51	0/2540
All	All	0.26	0/8829	0.50	0/12028

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1810	0	1681	22	0
1	B	1806	0	1697	17	0
1	C	1653	0	1532	17	0
1	D	1525	0	1375	14	0
1	E	1822	0	1723	22	0
2	A	1	0	0	0	0
All	All	8617	0	8008	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:83:ARG:H	1:C:87:VAL:HG22	1.56	0.70
1:B:177:ALA:HB1	1:B:196:LYS:HE3	1.72	0.70
1:A:69:LEU:HD21	1:D:122:PHE:HE2	1.63	0.63
1:B:122:PHE:HE2	1:C:69:LEU:HD21	1.66	0.61
1:D:148:ARG:NH1	1:D:243:THR:OG1	2.33	0.61
1:E:148:ARG:HD3	1:E:243:THR:HG21	1.81	0.61
1:B:15:MET:O	1:E:237:ARG:NH2	2.33	0.61
1:A:148:ARG:NH1	1:A:243:THR:OG1	2.34	0.60
1:C:122:PHE:HE2	1:D:69:LEU:HD21	1.66	0.60
1:B:67:ILE:HD11	1:B:124:PRO:HB3	1.83	0.59
1:C:67:ILE:HD11	1:C:124:PRO:HB3	1.84	0.59
1:A:122:PHE:HE2	1:E:69:LEU:HD21	1.68	0.58
1:A:95:LYS:NZ	1:A:295:ILE:O	2.27	0.58
1:E:67:ILE:HD11	1:E:124:PRO:HB3	1.85	0.57
1:E:93:GLY:HA2	1:E:129:LEU:HB2	1.87	0.57
1:D:101:PHE:O	1:D:102:ARG:HG2	2.06	0.56
1:D:67:ILE:HD11	1:D:124:PRO:HB3	1.88	0.55
1:B:42:HIS:O	1:B:44:GLU:N	2.38	0.55
1:A:101:PHE:O	1:A:102:ARG:HG2	2.05	0.54
1:E:75:ILE:HD13	1:E:287:ARG:HD2	1.89	0.54
1:E:101:PHE:O	1:E:102:ARG:HG2	2.08	0.54
1:B:58:GLU:OE2	1:E:237:ARG:NH2	2.41	0.54
1:A:93:GLY:HA2	1:A:129:LEU:HB2	1.91	0.53
1:C:237:ARG:NH2	1:D:58:GLU:OE2	2.41	0.52
1:D:125:ASP:OD1	1:D:126:ILE:N	2.43	0.52
1:A:67:ILE:HD11	1:A:124:PRO:HB3	1.91	0.52
1:B:124:PRO:HG2	1:B:145:LEU:O	2.11	0.51
1:B:80:GLU:HG3	1:B:81:ILE:HG22	1.92	0.51
1:B:101:PHE:O	1:B:102:ARG:HG2	2.11	0.51
1:B:82:GLY:C	1:B:87:VAL:HG22	2.32	0.49
1:A:126:ILE:HD13	1:A:134:VAL:HG21	1.93	0.49
1:C:101:PHE:O	1:C:102:ARG:HG2	2.13	0.48
1:C:126:ILE:HD13	1:C:134:VAL:HG21	1.96	0.47
1:A:40:TYR:HE2	1:A:42:HIS:ND1	2.12	0.47
1:E:125:ASP:OD1	1:E:126:ILE:N	2.47	0.47
1:A:218:THR:HG22	1:A:220:TYR:H	1.80	0.47
1:B:112:ILE:HB	1:B:172:LEU:HD23	1.95	0.46
1:A:207:ARG:NH1	1:A:290:GLU:OE1	2.39	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:TRP:CD1	1:A:236:ILE:HG12	2.50	0.46
1:A:58:GLU:OE2	1:D:237:ARG:NH2	2.49	0.46
1:E:124:PRO:HG2	1:E:145:LEU:O	2.16	0.46
1:E:218:THR:HG22	1:E:220:TYR:H	1.81	0.45
1:D:126:ILE:HD13	1:D:134:VAL:HG21	1.99	0.45
1:C:227:TYR:O	1:C:237:ARG:HD2	2.15	0.45
1:B:101:PHE:HB2	1:B:225:LEU:HD11	1.98	0.45
1:C:22:ALA:HB3	1:C:54:PHE:HE2	1.82	0.45
1:A:203:ASP:OD1	1:A:207:ARG:N	2.43	0.45
1:C:148:ARG:HD3	1:C:243:THR:HG21	1.99	0.45
1:A:40:TYR:CE2	1:A:42:HIS:ND1	2.85	0.44
1:B:78:TYR:HE1	1:B:87:VAL:HG23	1.82	0.44
1:B:97:ASP:O	1:B:125:ASP:N	2.50	0.44
1:A:223:TRP:HD1	1:A:236:ILE:HG12	1.82	0.44
1:D:97:ASP:O	1:D:125:ASP:N	2.48	0.44
1:A:164:LEU:HD22	1:A:223:TRP:HA	1.98	0.44
1:B:61:PHE:HB3	1:B:65:MET:HB3	2.00	0.44
1:D:124:PRO:HG2	1:D:145:LEU:O	2.18	0.43
1:A:212:LEU:HD21	1:A:219:ALA:HB2	2.01	0.43
1:C:101:PHE:HB2	1:C:225:LEU:HD11	2.00	0.43
1:A:209:TYR:O	1:A:218:THR:HG23	2.18	0.43
1:D:93:GLY:HA2	1:D:129:LEU:HB2	2.00	0.43
1:E:212:LEU:HD21	1:E:219:ALA:HB2	2.01	0.43
1:E:223:TRP:HD1	1:E:236:ILE:HG12	1.84	0.43
1:E:148:ARG:O	1:E:241:LEU:N	2.45	0.42
1:A:124:PRO:HG2	1:A:145:LEU:O	2.19	0.42
1:E:126:ILE:HD13	1:E:134:VAL:HG21	2.01	0.42
1:E:15:MET:HG3	1:E:17:SER:H	1.84	0.42
1:B:31:PRO:HG3	1:B:46:ILE:HD11	2.01	0.42
1:E:16:PHE:O	1:E:18:ASN:N	2.53	0.42
1:E:218:THR:HG21	1:E:220:TYR:HD2	1.85	0.42
1:E:213:GLU:O	1:E:214:ASP:CB	2.68	0.42
1:A:218:THR:HG21	1:A:220:TYR:HD2	1.86	0.41
1:C:122:PHE:CE2	1:D:69:LEU:HD21	2.51	0.41
1:C:22:ALA:HB3	1:C:54:PHE:CE2	2.56	0.41
1:B:82:GLY:O	1:B:87:VAL:HG22	2.20	0.41
1:C:97:ASP:O	1:C:125:ASP:N	2.52	0.41
1:C:124:PRO:HG2	1:C:145:LEU:O	2.21	0.41
1:D:102:ARG:O	1:D:102:ARG:HG3	2.20	0.41
1:E:158:LYS:HD3	1:E:160:MET:SD	2.60	0.41
1:E:158:LYS:O	1:E:158:LYS:HG3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:TYR:CE1	1:A:195:LEU:HD23	2.56	0.40
1:C:102:ARG:HA	1:C:112:ILE:HD11	2.02	0.40
1:E:66:THR:O	1:E:70:MET:HG3	2.21	0.40
1:C:78:TYR:HE1	1:C:87:VAL:HG23	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/310 (73%)	209 (93%)	16 (7%)	0	100	100
1	B	223/310 (72%)	210 (94%)	13 (6%)	0	100	100
1	C	208/310 (67%)	195 (94%)	13 (6%)	0	100	100
1	D	196/310 (63%)	186 (95%)	10 (5%)	0	100	100
1	E	225/310 (73%)	205 (91%)	19 (8%)	1 (0%)	34	66
All	All	1077/1550 (70%)	1005 (93%)	71 (7%)	1 (0%)	51	82

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	32	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/276 (66%)	182 (100%)	1 (0%)	88	96
1	B	182/276 (66%)	180 (99%)	2 (1%)	73	92
1	C	162/276 (59%)	158 (98%)	4 (2%)	47	78
1	D	145/276 (52%)	140 (97%)	5 (3%)	37	71
1	E	187/276 (68%)	182 (97%)	5 (3%)	44	77
All	All	859/1380 (62%)	842 (98%)	17 (2%)	55	82

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	GLU
1	B	76	ASP
1	B	140	ARG
1	C	19	LYS
1	C	60	ASN
1	C	90	SER
1	C	95	LYS
1	D	24	VAL
1	D	45	ASP
1	D	97	ASP
1	D	112	ILE
1	D	176	THR
1	E	16	PHE
1	E	38	ASP
1	E	203	ASP
1	E	222	SER
1	E	245	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	60	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/310 (75%)	-0.22	1 (0%) 92 93	10, 23, 39, 45	0
1	B	233/310 (75%)	-0.17	4 (1%) 70 69	11, 25, 49, 76	0
1	C	216/310 (69%)	0.15	4 (1%) 66 65	23, 45, 66, 73	0
1	D	206/310 (66%)	0.19	7 (3%) 45 40	23, 49, 72, 104	0
1	E	233/310 (75%)	-0.23	1 (0%) 92 93	9, 19, 38, 59	0
All	All	1121/1550 (72%)	-0.07	17 (1%) 73 73	9, 29, 62, 104	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	60	ASN	4.1
1	A	301	THR	3.6
1	D	126	ILE	3.5
1	B	16	PHE	3.3
1	C	229	TYR	3.1
1	C	81	ILE	3.0
1	B	300	PRO	3.0
1	D	103	LEU	3.0
1	B	13	GLU	2.9
1	B	14	TYR	2.6
1	D	138	GLU	2.6
1	D	89	GLU	2.5
1	D	141	LEU	2.4
1	E	178	TYR	2.3
1	D	90	SER	2.3
1	D	137	THR	2.1
1	C	102	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	401	1/1	0.99	0.10	23,23,23,23	1

6.5 Other polymers [i](#)

There are no such residues in this entry.