



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 11:12 am BST

PDB ID : 5QDN  
Title : PanDDA analysis group deposition – Crystal structure of PTP1B in complex with compound\_FMOPL000163a  
Authors : Keedy, D.A.; Hill, Z.B.; Biel, J.T.; Kang, E.; Rettenmaier, T.J.; Brandao-Neto, J.; von Delft, F.; Wells, J.A.; Fraser, J.S.  
Deposited on : 2018-08-30  
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TRS	A	402	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10618 atoms, of which 5159 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	282	10321	3252	5113	911	1017	28	0	113	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	SER	CYS	engineered mutation	UNP P18031
A	92	VAL	CYS	engineered mutation	UNP P18031

- Molecule 2 is N-{4-[(2S)-butan-2-yl]phenyl}methanesulfonamide (three-letter code: JFY) (formula: C<sub>11</sub>H<sub>17</sub>NO<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
2	A	1	64	22	34	2	4	2	0	1

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code:

TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
3	A	1	20	4	12	1	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	213	Total	O	0	0
			213	213		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.54Å 89.54Å 106.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.77 – 1.82 44.77 – 1.82	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.77-1.82) 100.0 (44.77-1.82)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 1.82Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.193 , 0.213 0.194 , 0.214	Depositor DCC
$R_{free}$ test set	1758 reflections (3.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.0	Xtrriage
Anisotropy	0.162	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.031 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10618	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JFY, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.50	1/5305 (0.0%)	0.63	4/7083 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	155	VAL	CB-CG2	-5.63	1.41	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	79	ARG	NE-CZ-NH2	-5.67	117.47	120.30
1	A	121	CYS	CA-CB-SG	5.18	123.33	114.00
1	A	79	ARG	NE-CZ-NH1	5.13	122.86	120.30
1	A	213	VAL	CA-CB-CG2	5.12	118.57	110.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5208	5113	5047	82	6
2	A	30	34	0	8	0
3	A	8	12	11	0	6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	213	0	0	24	0
All	All	5459	5159	5058	82	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29[C]:ASP:OD1	4:A:501:HOH:O	1.74	1.02
1:A:29[D]:ASP:OD1	4:A:501:HOH:O	1.74	1.02
1:A:73[C]:LYS:HE3	4:A:531:HOH:O	1.66	0.96
1:A:73[D]:LYS:HE3	4:A:531:HOH:O	1.66	0.96
1:A:201:SER:HA	2:A:401[C]:JFY:C15	1.98	0.93
1:A:201:SER:HA	2:A:401[D]:JFY:C15	1.98	0.93
1:A:76:GLU:OE1	1:A:238[D]:ARG:NH2	2.07	0.87
1:A:12[C]:LYS:HB3	1:A:12[C]:LYS:HZ2	1.41	0.83
1:A:12[D]:LYS:HZ2	1:A:12[D]:LYS:HB3	1.41	0.83
1:A:128[C]:LYS:NZ	4:A:502:HOH:O	1.74	0.80
1:A:128[D]:LYS:NZ	4:A:502:HOH:O	1.74	0.80
1:A:9[C]:GLN:O	1:A:13[C]:SER:OG	2.01	0.79
1:A:9[D]:GLN:O	1:A:13[D]:SER:OG	2.01	0.79
1:A:201:SER:CA	2:A:401[C]:JFY:C15	2.63	0.76
1:A:201:SER:CA	2:A:401[D]:JFY:C15	2.63	0.76
1:A:9[D]:GLN:O	1:A:9[D]:GLN:HG2	1.88	0.72
1:A:9[C]:GLN:O	1:A:9[C]:GLN:HG2	1.88	0.72
1:A:201:SER:CB	2:A:401[D]:JFY:C15	2.67	0.72
1:A:201:SER:CB	2:A:401[C]:JFY:C15	2.67	0.72
1:A:2[C]:GLU:HG3	1:A:2[C]:GLU:O	1.91	0.70
1:A:2[D]:GLU:O	1:A:2[D]:GLU:HG3	1.91	0.70
1:A:10[C]:ILE:O	1:A:14[C]:GLY:N	2.26	0.69
1:A:10[D]:ILE:O	1:A:14[D]:GLY:N	2.26	0.69
1:A:201:SER:HB2	2:A:401[C]:JFY:C15	2.24	0.68
1:A:201:SER:HB2	2:A:401[D]:JFY:C15	2.24	0.68
1:A:2[A]:GLU:HG2	1:A:3[A]:MET:N	2.11	0.65
1:A:2[B]:GLU:HG2	1:A:3[B]:MET:N	2.11	0.65
1:A:75[C]:GLU:HG3	4:A:648:HOH:O	1.96	0.65
1:A:75[D]:GLU:HG3	4:A:648:HOH:O	1.96	0.65
1:A:153[B]:TYR:HE1	1:A:155:VAL:CG2	2.09	0.65
1:A:21[C]:GLN:HG2	4:A:513:HOH:O	1.97	0.65
1:A:21[D]:GLN:HG2	4:A:513:HOH:O	1.97	0.65

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29[A]:ASP:OD1	4:A:501:HOH:O	2.15	0.63
1:A:29[B]:ASP:OD1	4:A:501:HOH:O	2.15	0.63
1:A:252[C]:GLU:OE1	4:A:504:HOH:O	2.17	0.60
1:A:252[D]:GLU:OE1	4:A:504:HOH:O	2.17	0.60
1:A:105[C]:ARG:HG2	1:A:105[C]:ARG:HH11	1.66	0.59
1:A:105[D]:ARG:HH11	1:A:105[D]:ARG:HG2	1.66	0.59
1:A:2[A]:GLU:HG2	1:A:3[A]:MET:H	1.68	0.59
1:A:2[B]:GLU:HG2	1:A:3[B]:MET:H	1.68	0.59
1:A:83:LEU:HD11	1:A:226[B]:CYS:SG	2.43	0.59
1:A:153[B]:TYR:HE1	1:A:155:VAL:HG23	1.69	0.56
1:A:241[D]:PRO:HG2	1:A:282[D]:MET:HE1	1.88	0.56
1:A:241[C]:PRO:HG2	1:A:282[C]:MET:HE1	1.88	0.56
1:A:227:LEU:HB2	1:A:253[D]:MET:HE1	1.90	0.54
1:A:3[B]:MET:HE1	1:A:274:VAL:HG12	1.90	0.53
1:A:3[A]:MET:HE1	1:A:274:VAL:HG12	1.90	0.53
1:A:227:LEU:HB2	1:A:253[C]:MET:HE1	1.90	0.52
1:A:92:VAL:HG12	1:A:135:PHE:CE1	2.45	0.52
1:A:241[C]:PRO:HG2	1:A:282[C]:MET:CE	2.39	0.51
1:A:241[D]:PRO:HG2	1:A:282[D]:MET:CE	2.39	0.51
1:A:153[B]:TYR:CE1	1:A:155:VAL:HG23	2.44	0.50
1:A:12[C]:LYS:HB3	1:A:12[C]:LYS:NZ	2.15	0.49
1:A:12[D]:LYS:NZ	1:A:12[D]:LYS:HB3	2.15	0.49
1:A:280[D]:PHE:HD2	1:A:281[D]:ILE:HD13	1.77	0.49
1:A:280[C]:PHE:HD2	1:A:281[C]:ILE:HD13	1.77	0.49
1:A:241[C]:PRO:HG3	4:A:553:HOH:O	2.13	0.48
1:A:241[D]:PRO:HG3	4:A:553:HOH:O	2.13	0.48
1:A:39[C]:LYS:HE3	4:A:585:HOH:O	2.13	0.47
1:A:39[D]:LYS:HE3	4:A:585:HOH:O	2.13	0.47
1:A:6[A]:GLU:OE2	1:A:247[A]:LYS:HE3	2.16	0.46
1:A:6[B]:GLU:OE2	1:A:247[B]:LYS:HE3	2.16	0.46
1:A:76:GLU:CD	1:A:238[C]:ARG:HH21	2.09	0.43
1:A:76:GLU:CD	1:A:238[D]:ARG:HH21	2.09	0.43
1:A:163[C]:LEU:O	4:A:506:HOH:O	2.21	0.43
1:A:163[D]:LEU:O	4:A:506:HOH:O	2.21	0.43
1:A:252[C]:GLU:CD	4:A:504:HOH:O	2.56	0.42
1:A:252[D]:GLU:CD	4:A:504:HOH:O	2.56	0.42
1:A:73[C]:LYS:CE	4:A:531:HOH:O	2.43	0.41
1:A:73[D]:LYS:CE	4:A:531:HOH:O	2.43	0.41
1:A:45[C]:ARG:NH2	1:A:121:CYS:HA	2.36	0.41
1:A:45[D]:ARG:NH2	1:A:121:CYS:HA	2.36	0.41
1:A:147[C]:GLU:HG3	1:A:156:ARG:HG2	2.03	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147[D]:GLU:HG3	1:A:156:ARG:HG2	2.03	0.41
1:A:153[B]:TYR:CE1	1:A:155:VAL:CG2	2.97	0.41
1:A:153[A]:TYR:OH	1:A:197[A]:LYS:HG3	2.20	0.41
1:A:12[C]:LYS:CB	1:A:12[C]:LYS:HZ2	2.21	0.40
1:A:12[D]:LYS:CB	1:A:12[D]:LYS:HZ2	2.21	0.40
1:A:280[C]:PHE:CD2	1:A:281[C]:ILE:HD13	2.56	0.40
1:A:280[D]:PHE:CD2	1:A:281[D]:ILE:HD13	2.56	0.40
1:A:2[A]:GLU:CG	1:A:3[A]:MET:N	2.82	0.40
1:A:2[B]:GLU:CG	1:A:3[B]:MET:N	2.82	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128[C]:LYS:HZ3	3:A:402:TRS:HO2[4_565]	0.59	1.01
1:A:128[D]:LYS:HZ3	3:A:402:TRS:HO2[4_565]	0.59	1.01
1:A:128[C]:LYS:NZ	3:A:402:TRS:HO2[4_565]	1.51	0.09
1:A:128[D]:LYS:NZ	3:A:402:TRS:HO2[4_565]	1.51	0.09
1:A:128[C]:LYS:HZ3	3:A:402:TRS:O2[4_565]	1.52	0.08
1:A:128[D]:LYS:HZ3	3:A:402:TRS:O2[4_565]	1.52	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	613/321 (191%)	589 (96%)	15 (2%)	9 (2%)	<b>10</b> <b>2</b>

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3[A]	MET
1	A	3[B]	MET

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	3[C]	MET
1	A	3[D]	MET
1	A	63[A]	ASP
1	A	63[B]	ASP
1	A	63[C]	ASP
1	A	63[D]	ASP
1	A	261	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	578/294 (197%)	563 (97%)	15 (3%)	46 32

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61[A]	GLN
1	A	61[B]	GLN
1	A	61[C]	GLN
1	A	61[D]	GLN
1	A	72	ILE
1	A	79	ARG
1	A	121	CYS
1	A	265[A]	ASP
1	A	265[B]	ASP
1	A	265[C]	ASP
1	A	265[D]	ASP
1	A	281[A]	ILE
1	A	281[B]	ILE
1	A	281[C]	ILE
1	A	281[D]	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	JFY	A	401[D]	-	15,15,15	3.05	7 (46%)	21,21,21	1.74	3 (14%)
3	TRS	A	402	-	7,7,7	0.84	0	9,9,9	0.96	1 (11%)
2	JFY	A	401[C]	-	15,15,15	3.05	7 (46%)	21,21,21	1.74	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JFY	A	401[D]	-	-	4/11/11/11	0/1/1/1
3	TRS	A	402	-	-	3/9/9/9	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JFY	A	401[C]	-	-	4/11/11/11	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[D]	JFY	C15-S12	-7.09	1.58	1.75
2	A	401[C]	JFY	C15-S12	-7.09	1.58	1.75
2	A	401[D]	JFY	S12-N11	-6.18	1.55	1.63
2	A	401[C]	JFY	S12-N11	-6.18	1.55	1.63
2	A	401[D]	JFY	C07-C06	-3.62	1.32	1.38
2	A	401[C]	JFY	C07-C06	-3.62	1.32	1.38
2	A	401[D]	JFY	O14-S12	-3.59	1.37	1.43
2	A	401[C]	JFY	O14-S12	-3.59	1.37	1.43
2	A	401[D]	JFY	C06-C05	-3.24	1.33	1.39
2	A	401[C]	JFY	C06-C05	-3.24	1.33	1.39
2	A	401[D]	JFY	O13-S12	-2.27	1.39	1.43
2	A	401[C]	JFY	O13-S12	-2.27	1.39	1.43
2	A	401[D]	JFY	C07-C08	-2.15	1.35	1.39
2	A	401[C]	JFY	C07-C08	-2.15	1.35	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[D]	JFY	C15-S12-N11	-5.11	100.80	106.63
2	A	401[C]	JFY	C15-S12-N11	-5.11	100.80	106.63
2	A	401[D]	JFY	O14-S12-N11	3.73	114.70	107.10
2	A	401[C]	JFY	O14-S12-N11	3.73	114.70	107.10
2	A	401[D]	JFY	C09-C08-N11	2.38	125.25	120.09
2	A	401[C]	JFY	C09-C08-N11	2.38	125.25	120.09
3	A	402	TRS	O3-C3-C	-2.32	103.63	111.00

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401[D]	JFY	C08-N11-S12-C15
2	A	401[D]	JFY	C08-N11-S12-O13
3	A	402	TRS	C2-C-C1-O1
3	A	402	TRS	C3-C-C1-O1
3	A	402	TRS	N-C-C1-O1
2	A	401[C]	JFY	C08-N11-S12-C15

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	401[C]	JFY	C08-N11-S12-O13
2	A	401[D]	JFY	C01-C02-C03-C04
2	A	401[C]	JFY	C01-C02-C03-C04
2	A	401[D]	JFY	C01-C02-C03-C05
2	A	401[C]	JFY	C01-C02-C03-C05

There are no ring outliers.

3 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401[D]	JFY	4	0
3	A	402	TRS	0	6
2	A	401[C]	JFY	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	6

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	63[C]:ASP	C	64:ASN	N	1.19
1	A	63[D]:ASP	C	64:ASN	N	1.19
1	A	89:PRO	C	90[C]:ASN	N	1.19
1	A	89:PRO	C	90[D]:ASN	N	1.19
1	A	28:SER	C	29[C]:ASP	N	1.17
1	A	28:SER	C	29[D]:ASP	N	1.17

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	282/321 (87%)	0.23	18 (6%) 19 15	22, 34, 56, 68	1 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281[A]	ILE	7.0
1	A	282[A]	MET	5.8
1	A	280[A]	PHE	4.5
1	A	63[A]	ASP	4.2
1	A	5[A]	LYS	3.8
1	A	121	CYS	3.8
1	A	62[A]	GLU	3.8
1	A	152[A]	TYR	3.6
1	A	12[A]	LYS	3.1
1	A	47[A]	ARG	2.9
1	A	1[A]	MET	2.6
1	A	165[A]	THR	2.6
1	A	25[A]	HIS	2.5
1	A	2[A]	GLU	2.3
1	A	13[A]	SER	2.1
1	A	279[A]	LYS	2.0
1	A	14[A]	GLY	2.0
1	A	10[A]	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	JFY	A	401[D]	15/15	0.66	0.30	31,54,73,73	32
2	JFY	A	401[C]	15/15	0.66	0.30	31,54,73,73	32
3	TRS	A	402	8/8	0.89	0.12	25,65,106,106	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.