



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 01:39 AM EDT

PDB ID : 2QF1
Title : Rat cytosolic PEPCK in complex with oxaloacetic acid.
Authors : Sullivan, S.M.; Holyoak, T.
Deposited on : 2007-06-26
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	624	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoenolpyruvate carboxykinase, cytosolic [GTP].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	606	5026	3249	844	900	33	0	46	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	cloning artifact	UNP P07379
A	0	SER	-	cloning artifact	UNP P07379


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is OXALOACETATE ION (three-letter code: OAA) (formula: C₄H₃O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 18 8 10	0	1


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	363	Total O 363 363	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoenolpyruvate carboxykinase, cytosolic [GTP]



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.22Å 119.34Å 60.81Å 90.00° 108.71° 90.00°	Depositor
Resolution (Å)	24.48 – 1.80 23.33 – 1.80	Depositor EDS
% Data completeness (in resolution range)	89.3 (24.48-1.80) 89.3 (23.33-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.80Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.194 , 0.228 0.193 , 0.227	Depositor DCC
R_{free} test set	2554 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtrriage
Anisotropy	0.096	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5410	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OAA, NA, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/5283	0.54	0/7140

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5026	0	5149	49	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
4	A	18	0	4	0	0
5	A	363	0	0	2	0
All	All	5410	0	5153	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:ILE:HG22	1:A:408[B]:THR:HG21	1.18	1.09
1:A:297[A]:ASN:H	1:A:297[A]:ASN:HD22	1.25	0.81
1:A:88[B]:ILE:HG21	1:A:91[B]:LYS:HD2	1.64	0.80
1:A:351:ILE:HG22	1:A:408[B]:THR:CG2	2.08	0.71
1:A:243:LYS:O	1:A:247:ALA:HB3	1.93	0.68
1:A:443:LEU:HB2	1:A:577[A]:LEU:HD22	1.81	0.62
1:A:194:HIS:HE1	1:A:233:SER:O	1.85	0.60
1:A:55:ARG:HD2	5:A:892:HOH:O	2.02	0.59
1:A:355:THR:HG23	1:A:408[B]:THR:HG22	1.85	0.58
1:A:355:THR:CG2	1:A:408[B]:THR:HG22	2.34	0.58
1:A:590:GLU:HG2	1:A:594[B]:LYS:HE2	1.90	0.55
1:A:161:THR:OG1	1:A:194:HIS:HD2	1.89	0.54
1:A:100[B]:ARG:NH2	5:A:1046:HOH:O	2.34	0.54
1:A:219[A]:ILE:HD11	1:A:245:CYS:SG	2.48	0.54
1:A:215[B]:GLU:CD	1:A:215[B]:GLU:H	2.13	0.52
1:A:477:HIS:HE2	1:A:591:GLU:CD	2.13	0.52
1:A:315:MET:HG2	1:A:325:ALA:HB2	1.94	0.50
1:A:89:GLU:OE1	1:A:483:ARG:NH2	2.43	0.49
1:A:542:GLY:O	1:A:547:GLU:HB2	2.13	0.49
1:A:503[B]:ARG:HG2	1:A:506:ALA:HB2	1.96	0.48
1:A:296:MET:O	1:A:305[B]:VAL:HG22	2.13	0.48
1:A:431:ILE:HB	1:A:511[A]:ILE:HD13	1.96	0.48
1:A:391:TRP:CD2	1:A:399:CYS:HB3	2.49	0.47
1:A:295:MET:CE	1:A:413:CYS:SG	3.03	0.46
1:A:142:ILE:HG23	1:A:176[B]:MET:CG	2.46	0.46
1:A:219[A]:ILE:CD1	1:A:245:CYS:SG	3.04	0.46
1:A:615:LEU:O	1:A:619[A]:ILE:HG12	2.16	0.45
1:A:565:ASN:C	1:A:565:ASN:HD22	2.20	0.45
1:A:17:GLN:NE2	1:A:52:GLU:OE1	2.50	0.45
1:A:526:LEU:HB3	1:A:577[A]:LEU:HG	1.97	0.45
1:A:261:LEU:HD11	1:A:500[A]:MET:SD	2.57	0.45
1:A:420:TRP:CE2	1:A:421:GLU:HG2	2.52	0.44
1:A:497:TRP:HA	1:A:500[A]:MET:CE	2.47	0.44
1:A:497:TRP:HA	1:A:500[A]:MET:HE2	1.99	0.44
1:A:295:MET:HE3	1:A:413:CYS:SG	2.58	0.43
1:A:593:ASP:HA	1:A:612:LEU:HD21	1.99	0.43
1:A:349:LYS:HD3	1:A:414:PRO:HG3	2.00	0.43
1:A:213:ASN:OD1	1:A:215[B]:GLU:HG2	2.18	0.43
1:A:60:MET:HB3	1:A:66:ILE:HG12	2.01	0.42
1:A:246:PHE:C	1:A:246:PHE:CD1	2.93	0.42
1:A:550:ALA:HB1	1:A:557:TYR:HB3	2.01	0.42
1:A:335[B]:VAL:HG22	1:A:337:PRO:HD2	2.00	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297[A]:ASN:H	1:A:297[A]:ASN:ND2	2.05	0.42
1:A:450:TRP:HZ2	1:A:500[A]:MET:HE3	1.85	0.41
1:A:149[B]:LEU:HD21	1:A:180:VAL:HG22	2.03	0.41
1:A:243:LYS:HE3	1:A:333:PHE:HB3	2.03	0.40
1:A:431:ILE:HB	1:A:511[A]:ILE:CD1	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	648/624 (104%)	628 (97%)	19 (3%)	1 (0%)	47 33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	552/520 (106%)	523 (95%)	29 (5%)	22 9

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51[A]	GLU
1	A	51[B]	GLU
1	A	103	VAL
1	A	119[A]	GLU
1	A	119[B]	GLU
1	A	203	LYS
1	A	219[A]	ILE
1	A	219[B]	ILE
1	A	243	LYS
1	A	246	PHE
1	A	253	ARG
1	A	276[A]	LYS
1	A	276[B]	LYS
1	A	297[A]	ASN
1	A	297[B]	ASN
1	A	304	LYS
1	A	377	LEU
1	A	395	ASP
1	A	421	GLU
1	A	473	LYS
1	A	522	ASN
1	A	547	GLU
1	A	552	LEU
1	A	565	ASN
1	A	569	LEU
1	A	572	VAL
1	A	577[A]	LEU
1	A	577[B]	LEU
1	A	587	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	59	HIS
1	A	61	GLN
1	A	194	HIS
1	A	209	ASN
1	A	330	ASN
1	A	565	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	OAA	A	800[B]	2	8,8,8	2.23	3 (37%)	9,10,10	1.17	0
4	OAA	A	800[A]	2	8,8,8	2.24	3 (37%)	9,10,10	1.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OAA	A	800[B]	2	-	4/8/8/8	-
4	OAA	A	800[A]	2	-	1/8/8/8	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	800[B]	OAA	O3-C3	4.26	1.31	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	800[A]	OAA	O3-C3	4.11	1.31	1.23
4	A	800[A]	OAA	O4-C4	3.57	1.32	1.22
4	A	800[B]	OAA	O4-C4	3.56	1.32	1.22
4	A	800[A]	OAA	O1-C1	2.79	1.31	1.22
4	A	800[B]	OAA	O1-C1	2.79	1.31	1.22

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	800[A]	OAA	C2-C3-C4-O5
4	A	800[B]	OAA	C1-C2-C3-O3
4	A	800[B]	OAA	C1-C2-C3-C4
4	A	800[B]	OAA	C2-C3-C4-O5
4	A	800[B]	OAA	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	606/624 (97%)	0.13	27 (4%) 33 27	24, 29, 35, 44	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	394	GLN	5.8
1	A	472	GLY	5.7
1	A	360	VAL	4.8
1	A	287	ALA	4.7
1	A	341	VAL	4.2
1	A	393	PRO	3.9
1	A	522	ASN	3.7
1	A	464	ALA	3.6
1	A	392	ARG	3.6
1	A	504	PRO	3.5
1	A	288	CYS	3.3
1	A	396	GLU	3.2
1	A	293	LEU	3.0
1	A	395	ASP	2.9
1	A	342	LYS	2.8
1	A	577[A]	LEU	2.8
1	A	443	LEU	2.8
1	A	562	ASP	2.8
1	A	389[A]	LYS	2.8
1	A	521	LYS	2.7
1	A	361	ALA	2.6
1	A	525	PHE	2.4
1	A	561	GLU	2.3
1	A	356[A]	ILE	2.3
1	A	397	GLU	2.3
1	A	569	LEU	2.1
1	A	517	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	OAA	A	800[A]	9/9	0.90	0.15	28,30,32,32	9
4	OAA	A	800[B]	9/9	0.90	0.15	29,32,32,33	9
3	NA	A	702	1/1	0.97	0.15	35,35,35,35	0
2	MN	A	701	1/1	0.98	0.08	25,25,25,25	1
2	MN	A	700	1/1	0.99	0.09	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.