

Full wwPDB X-ray Structure Validation Report (i)

Mar 4, 2024 - 09:39 AM EST

PDB ID : 4QJR

Title : Crystal structure of human nuclear receptor sf-1 (nr5a1) bound to its hormone

pip3 at 2.4 a resolution

Authors: Joint Center for Structural Genomics (JCSG); Partnership for Stem Cell Bi-

ology (STEMCELL)

Deposited on : 2014-06-04

Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	245	93%	7%
2	В	14	29%	14%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2294 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Steroidogenic factor 1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	244	Total	С	N	О	S	0	19	0
1	A	244	2013	1287	343	370	13	U	12	U

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q13285
A	247	SER	CYS	engineered mutation	UNP Q13285
A	412	SER	CYS	engineered mutation	UNP Q13285

• Molecule 2 is a protein called Peroxisome proliferator-activated receptor gamma coactivator 1-alpha.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	В	12	Total 89	C 61	N 14	O 14	0	0	0


• Molecule 3 is (2S)-3-{[(R)-{[(1S,2S,3R,4S,5S,6S)-2,6-dihydroxy-3,4,5-tris(phosphonooxy)cyclohexyl]oxy}(hydroxy)phosphoryl]oxy}propane -1,2-diyl dihexadecanoate (three-letter code: PIZ) (formula: $C_{41}H_{82}O_{22}P_4$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 67	C 41	O 22	P 4	0	0

 \bullet Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

 \bullet Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	117	Total O 117 117	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Steroidogenic factor 1



• Molecule 2: Peroxisome proliferator-activated receptor gamma coactivator 1-alpha







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	74.85Å 74.85Å 139.63Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.16 - 2.40	Depositor
rtesolution (A)	29.17 - 2.40	EDS
% Data completeness	99.5 (29.16-2.40)	Depositor
(in resolution range)	99.5 (29.17-2.40)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$< I/\sigma(I) > 1$	2.43 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
P. P.	0.190 , 0.230	Depositor
R, R_{free}	0.191 , 0.233	DCC
R_{free} test set	863 reflections (5.33%)	wwPDB-VP
Wilson B-factor (Å ²)	70.7	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.28 , 48.7	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2294	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PIZ, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.22	0/2083	0.36	0/2816	
2	В	0.18	0/90	0.30	0/119	
All	All	0.22	0/2173	0.36	0/2935	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2013	0	2069	11	0
2	В	89	0	110	0	0
3	A	67	0	75	3	0
4	A	8	0	6	0	0
5	A	117	0	0	1	0
All	All	2294	0	2260	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
1:A:427[A]:ARG:NH1	5:A:693:HOH:O	2.24	0.66
1:A:306:LEU:HB3	3:A:501:PIZ:H76	1.76	0.66
1:A:384:LEU:HD22	1:A:390:VAL:HG21	1.80	0.64
1:A:350[A]:ARG:NE	1:A:425[A]:GLU:OE2	2.34	0.61
3:A:501:PIZ:H15	3:A:501:PIZ:H78	1.85	0.58
1:A:351:ALA:HB2	3:A:501:PIZ:H3	1.88	0.56
1:A:409:TYR:CD1	1:A:415:LYS:HD2	2.43	0.53
1:A:304[B]:GLU:HG3	1:A:426:VAL:HG12	1.97	0.46
1:A:297:LEU:HD21	1:A:384:LEU:HD21	2.01	0.42
1:A:304[B]:GLU:HG2	1:A:375:ILE:HD13	2.02	0.41
1:A:258:GLN:NE2	1:A:445:GLU:OE1	2.44	0.41
1:A:220:GLY:HA2	1:A:408:HIS:NE2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$254/245\ (104\%)$	250 (98%)	4 (2%)	0	100	100
2	В	10/14~(71%)	9 (90%)	1 (10%)	0	100	100
All	All	264/259 (102%)	259 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	$228/217 \ (105\%)$	228 (100%)	0	100	100	
2	В	10/12 (83%)	10 (100%)	0	100	100	
All	All	238/229 (104%)	238 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain I		Dec	Res Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	A	502	-	3,3,3	0.76	0	3,3,3	1.31	0
4	ACT	A	503	-	3,3,3	0.77	0	3,3,3	1.27	0
3	PIZ	A	501	-	67,67,67	0.89	2 (2%)	83,88,88	0.87	6 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PIZ	A	501	-	-	22/65/89/89	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(ext{\AA})$
3	A	501	PIZ	OB1-CB1	3.32	1.43	1.34
3	A	501	PIZ	OA1-CA1	2.68	1.41	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
3	A	501	PIZ	C5-C6-C1	3.09	115.37	108.96
3	A	501	PIZ	OB1-CB1-OB2	-2.25	118.26	123.70
3	A	501	PIZ	OA1-CA1-CA2	2.24	118.93	111.91
3	A	501	PIZ	C3-C2-C1	2.21	113.55	108.96
3	A	501	PIZ	C6-C1-C2	2.19	114.02	110.85
3	A	501	PIZ	OB1-CB1-CB2	2.13	116.09	111.50

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	PIZ	C8-O13-P1-O12
3	A	501	PIZ	OB2-CB1-OB1-C7
3	A	501	PIZ	CB2-CB1-OB1-C7
3	A	501	PIZ	C8-O13-P1-O1
3	A	501	PIZ	CB9-CBA-CBB-CBC
3	A	501	PIZ	CB5-CB6-CB7-CB8
3	A	501	PIZ	CBC-CBD-CBE-CBF
3	A	501	PIZ	CB6-CB7-CB8-CB9
3	A	501	PIZ	CBA-CBB-CBC-CBD
3	A	501	PIZ	CB4-CB5-CB6-CB7
3	A	501	PIZ	CA1-CA2-CA3-CA4
3	A	501	PIZ	OB1-C7-C9-OA1
3	A	501	PIZ	C8-C7-C9-OA1
3	A	501	PIZ	CB1-CB2-CB3-CB4
3	A	501	PIZ	CAB-CAC-CAD-CAE
3	A	501	PIZ	CBD-CBE-CBF-CBG
3	A	501	PIZ	CAC-CAD-CAE-CAF

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	501	PIZ	C8-C7-OB1-CB1
3	A	501	PIZ	C8-O13-P1-O11
3	A	501	PIZ	C7-C8-O13-P1
3	A	501	PIZ	CA6-CA7-CA8-CA9
3	A	501	PIZ	CA8-CA9-CAA-CAB

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	PIZ	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	244/245 (99%)	-0.13	9 (3%) 41 41		46, 68, 110, 134	0
2	В	12/14 (85%)	1.56	4 (33%) 0 0		87, 107, 145, 146	0
All	All	$256/259 \ (98\%)$	-0.06	13 (5%) 28 26	6	46, 68, 119, 146	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	151	PRO	5.3
2	В	152	ALA	4.4
1	A	306	LEU	3.8
2	В	141	PRO	3.8
1	A	459	LYS	3.4
1	A	305	LEU	3.1
2	В	149	LEU	2.8
1	A	307	VAL	2.5
1	A	303	SER	2.5
1	A	276	ILE	2.3
1	A	383	PHE	2.3
1	A	461	THR	2.2
1	A	411	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	ACT	A	502	4/4	0.67	0.32	95,97,99,99	4
4	ACT	A	503	4/4	0.91	0.41	82,85,86,92	0
3	PIZ	A	501	67/67	0.95	0.16	50,69,108,118	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers (i)

There are no such residues in this entry.

