



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 11, 2024 – 01:25 pm GMT

PDB ID : 8QKG
Title : PvSub1 Catalytic Domain in Complex with Peptidomimetic Inhibitor (MAM-125)
Authors : Batista, F.A.; Martinez, M.; Bouillon, A.; Mechaly, A.; Alzari, P.M.; Haouz, A.; Barale, J.C.
Deposited on : 2023-09-15
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	631	
1	B	631	
2	C	7	
2	D	7	

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 5693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called subtilisin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2572	1613	452	495	12	0	1	0
1	B	336	2599	1629	458	500	12	0	0	0

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	MET	-	initiating methionine	UNP E6Y8B9
A	7	LYS	-	expression tag	UNP E6Y8B9
A	8	LEU	-	expression tag	UNP E6Y8B9
A	9	CYS	-	expression tag	UNP E6Y8B9
A	10	ILE	-	expression tag	UNP E6Y8B9
A	11	LEU	-	expression tag	UNP E6Y8B9
A	12	LEU	-	expression tag	UNP E6Y8B9
A	13	ALA	-	expression tag	UNP E6Y8B9
A	14	VAL	-	expression tag	UNP E6Y8B9
A	15	VAL	-	expression tag	UNP E6Y8B9
A	16	ALA	-	expression tag	UNP E6Y8B9
A	17	PHE	-	expression tag	UNP E6Y8B9
A	18	VAL	-	expression tag	UNP E6Y8B9
A	19	GLY	-	expression tag	UNP E6Y8B9
A	20	LEU	-	expression tag	UNP E6Y8B9
A	21	SER	-	expression tag	UNP E6Y8B9
A	22	LEU	-	expression tag	UNP E6Y8B9
A	23	GLY	-	expression tag	UNP E6Y8B9
A	24	ARG	-	expression tag	UNP E6Y8B9
A	25	SER	-	expression tag	UNP E6Y8B9
A	361	SER	ASN	engineered mutation	UNP E6Y8B9
A	432	SER	ASN	engineered mutation	UNP E6Y8B9
A	445	SER	ASN	engineered mutation	UNP E6Y8B9
A	631	HIS	-	expression tag	UNP E6Y8B9
A	632	HIS	-	expression tag	UNP E6Y8B9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	633	HIS	-	expression tag	UNP E6Y8B9
A	634	HIS	-	expression tag	UNP E6Y8B9
A	635	HIS	-	expression tag	UNP E6Y8B9
A	636	HIS	-	expression tag	UNP E6Y8B9
B	6	MET	-	initiating methionine	UNP E6Y8B9
B	7	LYS	-	expression tag	UNP E6Y8B9
B	8	LEU	-	expression tag	UNP E6Y8B9
B	9	CYS	-	expression tag	UNP E6Y8B9
B	10	ILE	-	expression tag	UNP E6Y8B9
B	11	LEU	-	expression tag	UNP E6Y8B9
B	12	LEU	-	expression tag	UNP E6Y8B9
B	13	ALA	-	expression tag	UNP E6Y8B9
B	14	VAL	-	expression tag	UNP E6Y8B9
B	15	VAL	-	expression tag	UNP E6Y8B9
B	16	ALA	-	expression tag	UNP E6Y8B9
B	17	PHE	-	expression tag	UNP E6Y8B9
B	18	VAL	-	expression tag	UNP E6Y8B9
B	19	GLY	-	expression tag	UNP E6Y8B9
B	20	LEU	-	expression tag	UNP E6Y8B9
B	21	SER	-	expression tag	UNP E6Y8B9
B	22	LEU	-	expression tag	UNP E6Y8B9
B	23	GLY	-	expression tag	UNP E6Y8B9
B	24	ARG	-	expression tag	UNP E6Y8B9
B	25	SER	-	expression tag	UNP E6Y8B9
B	361	SER	ASN	engineered mutation	UNP E6Y8B9
B	432	SER	ASN	engineered mutation	UNP E6Y8B9
B	445	SER	ASN	engineered mutation	UNP E6Y8B9
B	631	HIS	-	expression tag	UNP E6Y8B9
B	632	HIS	-	expression tag	UNP E6Y8B9
B	633	HIS	-	expression tag	UNP E6Y8B9
B	634	HIS	-	expression tag	UNP E6Y8B9
B	635	HIS	-	expression tag	UNP E6Y8B9
B	636	HIS	-	expression tag	UNP E6Y8B9

- Molecule 2 is a protein called Peptidomimetic Inhibitor (MAM-125).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			43	26	6	11			
2	D	7	Total	C	N	O	0	0	0
			43	26	6	11			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	A	3	3	3	0	0
4	B	3	3	3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	O S	0	0
			5	4 1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	212	Total	O	0	0
			212	212		
6	B	194	Total	O	0	0
			194	194		
6	C	2	Total	O	0	0
			2	2		
6	D	3	Total	O	0	0
			3	3		




- Molecule 2: Peptidomimetic Inhibitor (MAM-125)

Chain C: 71% 14% 14%



- Molecule 2: Peptidomimetic Inhibitor (MAM-125)

Chain D: 57% 29% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	53.38Å 55.16Å 69.03Å 69.20° 78.30° 74.75°	Depositor
Resolution (Å)	30.75 – 1.54 30.75 – 1.54	Depositor EDS
% Data completeness (in resolution range)	94.0 (30.75-1.54) 94.0 (30.75-1.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.54Å)	Xtrriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.179 , 0.204 0.174 , 0.196	Depositor DCC
R_{free} test set	4856 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtrriage
Anisotropy	0.280	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5693	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5XU, SO4, ACE, NAG, TBG, VEF, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2623	0.61	0/3560
1	B	0.49	0/2648	0.61	0/3594
2	C	1.98	2/18 (11.1%)	1.28	0/22
2	D	2.04	2/18 (11.1%)	1.25	0/22
All	All	0.53	4/5307 (0.1%)	0.62	0/7198

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	3
2	D	0	3
All	All	0	6

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	6	ASP	CG-OD2	6.14	1.39	1.25
2	D	6	ASP	CG-OD1	5.88	1.38	1.25
2	C	6	ASP	CG-OD1	5.81	1.38	1.25
2	C	6	ASP	CG-OD2	5.79	1.38	1.25

There are no bond angle outliers.

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	5	VEF	Mainchain,Peptide
2	C	6	ASP	Sidechain
2	D	5	VEF	Mainchain,Peptide
2	D	6	ASP	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2572	0	2521	5	0
1	B	2599	0	2542	10	0
2	C	43	0	35	0	0
2	D	43	0	35	1	0
3	A	14	0	13	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
5	B	5	0	0	0	0
6	A	212	0	0	1	0
6	B	194	0	0	2	0
6	C	2	0	0	0	0
6	D	3	0	0	0	0
All	All	5693	0	5146	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:311:LYS:H	1:B:429:HIS:HD2	1.19	0.89
1:A:311:LYS:H	1:A:429:HIS:HD2	1.20	0.89
1:A:311:LYS:H	1:A:429:HIS:CD2	2.05	0.70
1:B:311:LYS:H	1:B:429:HIS:CD2	2.05	0.70
1:B:532:ASN:H	1:B:546:ASN:HD22	1.40	0.68
1:B:532:ASN:H	1:B:546:ASN:ND2	1.90	0.68
1:B:506:ASP:O	1:B:508:ASP:O	2.23	0.57
1:B:278:LYS:HG2	1:B:299:GLN:HG3	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:LYS:HG3	1:B:278:LYS:O	2.15	0.46
6:B:1001:HOH:O	2:D:3:THR:HG23	2.15	0.45
1:A:605:ASN:ND2	6:A:804:HOH:O	2.50	0.45
1:B:460:VAL:HG21	1:B:487:PRO:HD2	1.99	0.44
1:A:290:LEU:HD22	1:A:295:LEU:HD22	2.01	0.42
1:B:393:VAL:HB	1:B:557:ALA:CB	2.52	0.40
1:B:410:LEU:HD23	6:B:1001:HOH:O	2.21	0.40
1:A:460:VAL:HG21	1:A:487:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/631 (52%)	318 (97%)	8 (2%)	3 (1%)	17	3
1	B	332/631 (53%)	319 (96%)	10 (3%)	3 (1%)	17	3
2	C	3/7 (43%)	3 (100%)	0	0	100	100
2	D	3/7 (43%)	3 (100%)	0	0	100	100
All	All	667/1276 (52%)	643 (96%)	18 (3%)	6 (1%)	17	3

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	371	TYR
1	B	371	TYR
1	A	520	SER
1	B	520	SER
1	A	463	SER
1	B	463	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/551 (52%)	285 (99%)	2 (1%)	84	68
1	B	289/551 (52%)	288 (100%)	1 (0%)	92	84
2	C	2/2 (100%)	2 (100%)	0	100	100
2	D	2/2 (100%)	2 (100%)	0	100	100
All	All	580/1106 (52%)	577 (100%)	3 (0%)	88	77

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	550	MET
1	A	579	ARG
1	B	550	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	285	ASN
1	A	429	HIS
1	A	484	HIS
1	A	540	ASN
1	A	605	ASN
1	B	285	ASN
1	B	338	HIS
1	B	429	HIS
1	B	516	ASN
1	B	540	ASN
1	B	546	ASN
1	B	605	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TBG	D	2	2	6,7,8	0.50	0	7,10,12	0.81	0
2	VEF	C	5	2,1	4,6,8	0.56	0	3,7,12	0.24	0
2	5XU	D	7	2	3,4,4	0.37	0	2,4,4	0.57	0
2	TBG	C	2	2	6,7,8	0.46	0	7,10,12	0.77	0
2	VEF	D	5	2,1	4,6,8	0.46	0	3,7,12	0.42	0
2	5XU	C	7	2	3,4,4	0.39	0	2,4,4	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TBG	D	2	2	-	0/6/8/10	-
2	VEF	C	5	2,1	-	0/5/6/12	-
2	5XU	D	7	2	-	0/0/2/2	-
2	TBG	C	2	2	-	0/6/8/10	-
2	VEF	D	5	2,1	-	0/5/6/12	-
2	5XU	C	7	2	-	0/0/2/2	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	B	904	-	4,4,4	0.20	0	6,6,6	0.74	0
3	NAG	A	701	1	14,14,15	0.32	0	17,19,21	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	701	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	NAG	C4-C5-C6-O6
3	A	701	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/631 (52%)	0.00	8 (2%) 59 64	13, 19, 30, 46	0
1	B	336/631 (53%)	0.08	13 (3%) 39 45	14, 20, 34, 50	0
2	C	3/7 (42%)	-0.35	0 100 100	18, 18, 19, 21	0
2	D	3/7 (42%)	0.01	0 100 100	21, 21, 22, 25	0
All	All	674/1276 (52%)	0.04	21 (3%) 49 56	13, 19, 32, 50	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	277	TYR	11.0
1	A	507	LEU	5.5
1	B	509	GLU	4.0
1	B	507	LEU	3.9
1	B	474	ASP	3.9
1	A	476	ALA	3.7
1	B	613	ALA	3.5
1	A	590	GLU	3.4
1	B	612	ALA	3.4
1	B	466	ALA	3.3
1	B	475	ILE	3.0
1	A	380	ILE	2.7
1	B	590	GLU	2.5
1	B	276	GLY	2.4
1	B	508	ASP	2.4
1	A	465	CYS	2.4
1	B	380	ILE	2.3
1	B	307	VAL	2.3
1	B	539	MET	2.2
1	A	466	ALA	2.1
1	A	558	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	5XU	C	7	5/5	0.84	0.21	23,29,32,34	0
2	5XU	D	7	5/5	0.85	0.18	23,28,31,33	0
2	TBG	D	2	8/9	0.92	0.10	19,20,21,21	0
2	TBG	C	2	8/9	0.94	0.08	17,17,19,19	0
2	VEF	C	5	7/9	0.97	0.06	15,15,15,15	0
2	VEF	D	5	7/9	0.97	0.06	16,17,18,19	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	701	14/15	0.77	0.23	31,37,39,40	0
4	CA	B	903	1/1	0.99	0.04	19,19,19,19	0
5	SO4	B	904	5/5	0.99	0.07	25,25,27,27	0
4	CA	A	704	1/1	1.00	0.04	19,19,19,19	0
4	CA	B	901	1/1	1.00	0.03	15,15,15,15	0
4	CA	B	902	1/1	1.00	0.04	18,18,18,18	0
4	CA	A	702	1/1	1.00	0.06	15,15,15,15	0
4	CA	A	703	1/1	1.00	0.04	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.