



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2023 – 03:00 pm GMT

PDB ID : 1QL6  
Title : THE CATALYTIC MECHANISM OF PHOSPHORYLASE KINASE  
PROBED BY MUTATIONAL STUDIES  
Authors : Skamnaki, V.T.; Owen, D.J.; Noble, M.E.M.; Lowe, E.D.; Oikonomakos, N.G.;  
Johnson, L.N.  
Deposited on : 1999-08-24  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	298	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2412 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHOSPHORYLASE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	281	2270	1453	377	424	16	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	SER	GLU	engineered mutation	UNP P00518

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	10	5	13	3	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mn	0	0
			2	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	104	Total	O	0	0
			104	104		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: PHOSPHORYLASE KINASE



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.56Å 68.17Å 112.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40	Depositor
% Data completeness (in resolution range)	86.6 (20.00-2.40)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.05	Depositor
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.240 , 0.330	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2412	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN, ATP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/2322	1.79	43/3137 (1.4%)

There are no bond length outliers.

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	34	ARG	NE-CZ-NH2	-18.77	110.91	120.30
1	A	127	ARG	NE-CZ-NH1	-13.93	113.34	120.30
1	A	175	ASP	CB-CG-OD1	12.50	129.55	118.30
1	A	275	ARG	CD-NE-CZ	10.56	138.38	123.60
1	A	35	ARG	NE-CZ-NH1	-10.55	115.03	120.30
1	A	275	ARG	NE-CZ-NH1	9.62	125.11	120.30
1	A	266	ARG	NE-CZ-NH2	-8.49	116.06	120.30
1	A	178	GLU	OE1-CD-OE2	-8.45	113.17	123.30
1	A	127	ARG	NH1-CZ-NH2	8.35	128.59	119.40
1	A	27	ARG	NE-CZ-NH2	8.33	124.47	120.30
1	A	34	ARG	NE-CZ-NH1	8.33	124.46	120.30
1	A	167	ASP	CB-CG-OD1	7.84	125.36	118.30
1	A	260	VAL	CB-CA-C	-7.84	96.51	111.40
1	A	148	ARG	NE-CZ-NH2	-7.68	116.46	120.30
1	A	16	TYR	CB-CG-CD2	7.19	125.32	121.00
1	A	93	TYR	CB-CG-CD1	-7.04	116.78	121.00
1	A	35	ARG	NH1-CZ-NH2	7.04	127.14	119.40
1	A	44	GLU	OE1-CD-OE2	-6.83	115.10	123.30
1	A	23	GLU	OE1-CD-OE2	-6.82	115.12	123.30
1	A	275	ARG	NE-CZ-NH2	-6.78	116.91	120.30
1	A	158	ASP	CB-CG-OD2	6.37	124.03	118.30
1	A	249	GLY	N-CA-C	6.37	129.03	113.10
1	A	114	TYR	CB-CG-CD1	6.14	124.68	121.00
1	A	59	SER	N-CA-CB	6.10	119.64	110.50

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	THR	N-CA-CB	6.07	121.83	110.30
1	A	178	GLU	CA-CB-CG	5.95	126.49	113.40
1	A	131	ARG	NE-CZ-NH1	-5.89	117.35	120.30
1	A	131	ARG	CD-NE-CZ	5.87	131.82	123.60
1	A	125	GLU	OE1-CD-OE2	-5.73	116.42	123.30
1	A	20	GLU	OE1-CD-OE2	-5.66	116.51	123.30
1	A	91	ASP	CB-CG-OD2	5.62	123.36	118.30
1	A	256	TYR	CB-CG-CD2	5.56	124.34	121.00
1	A	141	HIS	CB-CA-C	5.55	121.50	110.40
1	A	173	GLN	CA-CB-CG	5.35	125.16	113.40
1	A	279	GLU	OE1-CD-OE2	5.23	129.57	123.30
1	A	262	ASP	CB-CG-OD2	5.21	122.99	118.30
1	A	153	GLU	CG-CD-OE1	5.20	128.71	118.30
1	A	27	ARG	NE-CZ-NH1	-5.16	117.72	120.30
1	A	35	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	268	LEU	C-N-CA	5.10	134.46	121.70
1	A	224	ALA	CB-CA-C	-5.08	102.47	110.10
1	A	193	GLU	OE1-CD-OE2	-5.08	117.20	123.30
1	A	132	ALA	N-CA-CB	5.03	117.14	110.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2270	0	2253	44	0
2	A	31	0	12	0	0
3	A	2	0	0	0	0
4	A	5	0	0	0	0
5	A	104	0	0	1	0
All	All	2412	0	2265	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:SER:HB3	1:A:251:PRO:CD	1.99	0.92
1:A:250:SER:HB3	1:A:251:PRO:HD3	1.49	0.91
1:A:175:ASP:HB3	1:A:176:PRO:HD2	1.72	0.70
1:A:12:THR:OG1	1:A:95:THR:HG21	1.93	0.69
1:A:219:MET:HG2	1:A:264:VAL:HG22	1.73	0.69
1:A:230:TRP:HA	1:A:237:MET:HE2	1.74	0.69
1:A:11:SER:OG	1:A:95:THR:HG22	1.93	0.68
1:A:250:SER:CB	1:A:251:PRO:CD	2.72	0.67
1:A:260:VAL:HG13	1:A:286:PHE:CE1	2.30	0.67
1:A:284:HIS:CG	1:A:285:PRO:HD2	2.35	0.61
1:A:154:ASN:O	1:A:166:THR:HG22	2.03	0.59
1:A:251:PRO:HD2	1:A:252:GLU:OE1	2.02	0.59
1:A:138:CYS:O	1:A:142:LYS:HG3	2.04	0.58
1:A:33:VAL:HG22	1:A:48:LYS:HB2	1.86	0.57
1:A:218:ILE:O	1:A:222:LEU:HG	2.07	0.55
1:A:72:LYS:HG2	5:A:409:HOH:O	2.06	0.55
1:A:181:ARG:NH2	1:A:203:HIS:O	2.39	0.54
1:A:175:ASP:HB3	1:A:176:PRO:CD	2.38	0.54
1:A:41:THR:O	1:A:42:CYS:HB2	2.08	0.54
1:A:48:LYS:HE2	1:A:50:ILE:HD11	1.91	0.53
1:A:52:VAL:HG22	1:A:97:THR:O	2.11	0.51
1:A:13:HIS:O	1:A:17:GLU:N	2.37	0.51
1:A:156:LEU:O	1:A:163:ILE:HA	2.10	0.50
1:A:51:ASP:OD2	1:A:55:GLY:N	2.43	0.50
1:A:59:SER:O	1:A:60:ALA:C	2.51	0.48
1:A:15:PHE:CZ	1:A:47:VAL:HG21	2.48	0.48
1:A:231:HIS:ND1	1:A:232:ARG:N	2.62	0.48
1:A:174:LEU:HD21	1:A:180:LEU:HD11	1.96	0.47
1:A:266:ARG:O	1:A:276:TYR:HB2	2.14	0.47
1:A:187:PRO:HA	1:A:190:LEU:HD12	1.97	0.46
1:A:138:CYS:SG	1:A:279:GLU:HG2	2.55	0.45
1:A:271:GLN:OE1	1:A:273:GLN:OE1	2.36	0.44
1:A:83:HIS:CG	1:A:84:PRO:HD2	2.52	0.44
1:A:87:ILE:HD12	1:A:87:ILE:HA	1.74	0.43
1:A:53:THR:HG22	1:A:56:GLY:H	1.83	0.43
1:A:27:ARG:HG2	1:A:32:VAL:HG22	2.00	0.43
1:A:233:LYS:HB3	1:A:233:LYS:HE2	1.83	0.43
1:A:183:VAL:HG21	1:A:198:SER:HB3	2.01	0.42
1:A:220:TYR:CD1	1:A:228:PRO:HD3	2.55	0.42
1:A:285:PRO:HA	1:A:288:GLN:HG2	2.02	0.42
1:A:230:TRP:HA	1:A:237:MET:CE	2.47	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ARG:CZ	1:A:71:LEU:HD11	2.50	0.41
1:A:271:GLN:HB3	1:A:273:GLN:OE1	2.20	0.41
1:A:157:LEU:HD13	1:A:161:MET:CE	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	279/298 (94%)	262 (94%)	15 (5%)	2 (1%)	22 32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	SER
1	A	250	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	254/268 (95%)	246 (97%)	8 (3%)	40 60

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	GLU
1	A	66	LEU
1	A	88	GLN
1	A	96	ASN
1	A	138	CYS
1	A	260	VAL
1	A	276	TYR
1	A	291	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	271	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ATP	A	301	3	26,33,33	1.46	4 (15%)	31,52,52	1.54	5 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	304	-	4,4,4	0.67	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	301	3	-	2/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	ATP	C2-N1	3.09	1.39	1.33
2	A	301	ATP	C8-N7	-3.04	1.29	1.34
2	A	301	ATP	C5-N7	-2.39	1.31	1.39
2	A	301	ATP	C2'-C1'	2.31	1.57	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	ATP	C1'-N9-C4	4.68	134.86	126.64
2	A	301	ATP	C5-C6-N6	3.15	125.14	120.35
2	A	301	ATP	C4-C5-N7	2.86	112.38	109.40
2	A	301	ATP	O3'-C3'-C2'	2.42	119.66	111.82
2	A	301	ATP	N3-C2-N1	-2.32	125.06	128.68

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	ATP	PB-O3B-PG-O2G
2	A	301	ATP	PB-O3B-PG-O3G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.