Full wwPDB X-ray Structure Validation Report (i)

Oct 23, 2021 – 12:28 PM EDT

PDB ID	:	1QPK
Title	:	MUTANT (D193G) MALTOTETRAOSE-FORMING EXO-AMYLASE IN COMPLEX WITH MALTOTETRAOSE
Authors	:	Yoshioka, Y.; Hasegawa, K.; Matsuura, Y.; Katsube, Y.; Kubota, M.
Deposited on	:	1999-05-26
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	% 74% 20% 5% •
2	B	4	75% 25%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (MALTOTETRAOSE-FORMING AMYLASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	418	3293	2068	596	619	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	193	GLY	ASP	engineered mutation	UNP P13507

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	4	45	24	21	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	2	2	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	173	173	173	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTEIN (MALTOTETRAOSE-FORMING AMYLASE)

Chain A:



• Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	63.21Å 167.50Å 46.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 36.66 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 90.6 (36.66-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.79 (at 2.00Å)	Xtriage
Refinement program	PROFFT	Depositor
R, R_{free}	0.178 , (Not available) 0.163 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å^2)	22.3	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 57.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	3513	wwPDB-VP
Average B, all atoms (Å^2)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.89	1/3398 (0.0%)	1.75	56/4625 (1.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82	ASP	CA-CB	6.94	1.69	1.53

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	346	ARG	CD-NE-CZ	40.21	179.89	123.60
1	A	82	ASP	CB-CG-OD2	-16.07	103.84	118.30
1	A	175	ARG	NE-CZ-NH1	15.77	128.18	120.30
1	A	417	ARG	NE-CZ-NH1	14.87	127.73	120.30
1	A	352	ARG	NE-CZ-NH1	14.33	127.47	120.30
1	A	346	ARG	NE-CZ-NH1	13.71	127.15	120.30
1	A	191	ARG	NE-CZ-NH1	13.18	126.89	120.30
1	A	417	ARG	NE-CZ-NH2	-12.69	113.95	120.30
1	A	352	ARG	NE-CZ-NH2	-10.79	114.91	120.30
1	A	281	ARG	NE-CZ-NH2	10.56	125.58	120.30
1	A	11	ARG	NE-CZ-NH1	10.07	125.34	120.30
1	A	155	ARG	NE-CZ-NH1	9.98	125.29	120.30
1	A	88	ARG	NE-CZ-NH1	9.66	125.13	120.30
1	A	358	ARG	NE-CZ-NH1	-9.18	115.71	120.30
1	A	152	ASP	CB-CG-OD1	9.02	126.41	118.30
1	A	82	ASP	CB-CG-OD1	7.95	125.46	118.30
1	A	82	ASP	CA-CB-CG	-7.86	96.10	113.40
1	A	360	ASP	CB-CG-OD1	7.75	125.27	118.30
1	A	248	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	A	82	ASP	CB-CA-C	-7.30	95.81	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	233	ARG	NE-CZ-NH2	-7.29	116.65	120.30
1	A	279	ASP	CB-CG-OD2	-6.88	112.11	118.30
1	A	16	ASP	CB-CG-OD2	6.88	124.49	118.30
1	A	155	ARG	NE-CZ-NH2	-6.77	116.91	120.30
1	A	358	ARG	NH1-CZ-NH2	6.70	126.77	119.40
1	A	343	ASP	CB-CG-OD1	6.69	124.32	118.30
1	A	212	ASP	CB-CG-OD1	6.61	124.25	118.30
1	A	353	ARG	CD-NE-CZ	6.42	132.59	123.60
1	A	152	ASP	CB-CG-OD2	-6.39	112.55	118.30
1	A	137	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	A	160	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	227	TYR	CB-CG-CD1	6.20	124.72	121.00
1	A	137	ARG	NE-CZ-NH1	6.17	123.39	120.30
1	A	122	TYR	CB-CG-CD2	-6.13	117.32	121.00
1	A	279	ASP	CB-CG-OD1	6.05	123.75	118.30
1	A	182	ARG	NE-CZ-NH2	5.91	123.25	120.30
1	A	88	ARG	NE-CZ-NH2	-5.91	117.34	120.30
1	A	338	ASP	CB-CG-OD1	5.91	123.62	118.30
1	A	346	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	A	334	ASP	CA-CB-CG	-5.79	100.66	113.40
1	A	140	CYS	CA-CB-SG	5.62	124.12	114.00
1	A	358	ARG	NE-CZ-NH2	-5.61	117.50	120.30
1	A	290	VAL	CA-CB-CG2	5.56	119.23	110.90
1	A	334	ASP	CB-CG-OD2	-5.56	113.30	118.30
1	A	175	ARG	NH1-CZ-NH2	-5.37	113.49	119.40
1	A	347	GLN	CA-CB-CG	5.27	124.99	113.40
1	A	227	TYR	CB-CG-CD2	-5.21	117.87	121.00
1	A	82	ASP	N-CA-C	5.16	124.94	111.00
1	A	81	HIS	C-N-CA	5.14	134.55	121.70
1	A	397	VAL	N-CA-CB	-5.12	100.23	111.50
1	A	16	ASP	CB-CG-OD1	-5.12	113.69	118.30
1	A	148	ASN	CB-CA-C	5.10	120.60	110.40
1	A	334	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	124	ASP	CB-CG-OD2	-5.06	113.75	118.30
1	A	209	ASP	CB-CG-OD1	5.04	122.84	118.30
1	A	212	ASP	CB-CG-OD2	-5.03	113.77	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3293	0	3012	84	0
2	B	45	0	38	1	0
3	A	2	0	0	0	0
4	A	173	0	0	7	0
All	All	3513	0	3050	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:PRO:HD2	4:A:731:HOH:O	1.51	1.10
1:A:11:ARG:HH12	1:A:204:ASN:HD22	1.02	1.02
1:A:233:ARG:HG2	1:A:241:ILE:HG12	1.47	0.96
1:A:167:HIS:HD2	1:A:169:GLN:H	1.23	0.87
1:A:334:ASP:OD2	1:A:338:ASP:OD1	1.93	0.86
1:A:317:GLN:H	1:A:317:GLN:HE21	0.93	0.86
1:A:277:ASN:HD22	1:A:279:ASP:H	1.24	0.85
1:A:11:ARG:HH12	1:A:204:ASN:ND2	1.77	0.82
1:A:317:GLN:H	1:A:317:GLN:NE2	1.74	0.82
1:A:317:GLN:HE21	1:A:317:GLN:N	1.77	0.79
1:A:81:HIS:HD2	4:A:508:HOH:O	1.66	0.78
1:A:114:VAL:HG22	4:A:723:HOH:O	1.86	0.74
1:A:145:ASN:ND2	1:A:155:ARG:H	1.84	0.73
1:A:237:SER:H	1:A:240:GLN:HE21	1.35	0.73
1:A:145:ASN:HD21	1:A:154:ASP:HB3	1.57	0.70
1:A:145:ASN:HD21	1:A:155:ARG:H	1.39	0.70
1:A:292:ASN:HD22	1:A:294:ASP:H	1.37	0.69
1:A:60:TRP:HB3	1:A:82:ASP:O	1.93	0.68
1:A:26:ASN:HD22	1:A:29:ARG:HE	1.40	0.67
1:A:11:ARG:NH1	1:A:204:ASN:HD22	1.84	0.66
1:A:167:HIS:CD2	1:A:169:GLN:H	2.11	0.65
1:A:325:SER:HB2	1:A:326:PRO:CD	2.26	0.64
1:A:237:SER:H	1:A:240:GLN:NE2	1.95	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ARG:NH1	1:A:182:ARG:HG2	2.13	0.64
1:A:86:ASN:ND2	1:A:92:ASP:H	1.96	0.63
1:A:24:HIS:HB3	1:A:334:ASP:OD1	1.99	0.62
1:A:175:ARG:NH1	1:A:176:ASP:OD2	2.33	0.61
1:A:216:CYS:O	1:A:252:PRO:HD2	2.02	0.60
1:A:319:TYR:O	1:A:323:LEU:HB2	2.03	0.59
1:A:40:ARG:NH1	4:A:772:HOH:O	2.34	0.59
1:A:26:ASN:ND2	1:A:29:ARG:HE	2.02	0.58
1:A:11:ARG:HH22	1:A:204:ASN:ND2	2.01	0.58
1:A:37:ASN:HD21	1:A:94:GLN:NE2	2.01	0.57
1:A:1:ASP:OD1	1:A:108:LYS:NZ	2.35	0.57
1:A:292:ASN:ND2	1:A:294:ASP:H	2.03	0.57
1:A:366:HIS:HE1	1:A:375:THR:OG1	1.88	0.56
1:A:88:ARG:HD3	4:A:790:HOH:O	2.04	0.56
1:A:2:GLN:NE2	1:A:358:ARG:HH12	2.04	0.55
1:A:50:GLY:HA2	1:A:353:ARG:HH12	1.72	0.54
1:A:404:GLU:HG2	4:A:771:HOH:O	2.07	0.54
1:A:2:GLN:HE22	1:A:358:ARG:NH1	2.05	0.53
1:A:292:ASN:ND2	1:A:295:THR:H	2.07	0.53
1:A:155:ARG:HA	1:A:162:ASP:OD1	2.07	0.53
1:A:143:PRO:HB2	1:A:146:TYR:HE1	1.74	0.53
1:A:311:GLN:HG2	4:A:749:HOH:O	2.09	0.52
1:A:114:VAL:O	1:A:114:VAL:HG12	2.10	0.52
1:A:2:GLN:NE2	1:A:358:ARG:NH1	2.56	0.52
1:A:182:ARG:HG2	1:A:182:ARG:HH11	1.73	0.52
1:A:332:TYR:HB3	1:A:335:HIS:HD2	1.75	0.51
1:A:325:SER:HB2	1:A:326:PRO:HD2	1.90	0.51
1:A:277:ASN:ND2	1:A:279:ASP:H	2.01	0.51
1:A:266:SER:O	1:A:269:ASP:HB2	2.11	0.50
1:A:277:ASN:O	1:A:283:ARG:HD3	2.11	0.50
1:A:388:ASN:HD22	1:A:412:GLN:HB3	1.76	0.50
1:A:24:HIS:HD2	1:A:26:ASN:H	1.59	0.50
1:A:182:ARG:HH11	1:A:182:ARG:CG	2.24	0.49
1:A:50:GLY:HA2	1:A:353:ARG:NH1	2.27	0.48
1:A:145:ASN:HD21	1:A:154:ASP:CB	2.25	0.47
1:A:195:VAL:HG22	1:A:219:GLU:O	2.15	0.47
1:A:388:ASN:ND2	1:A:412:GLN:HB3	2.30	0.46
1:A:201:GLU:OE1	1:A:248:ARG:HD2	2.16	0.46
1:A:145:ASN:ND2	1:A:155:ARG:HG2	2.31	0.45
1:A:204:ASN:HB2	1:A:249:ALA:O	2.16	0.45
1:A:137:ARG:HG3	1:A:151:ASP:HB3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:PRO:HB3	1:A:152:ASP:HA	1.99	0.45
1:A:146:TYR:HB2	1:A:147:PRO:HD2	1.99	0.45
1:A:250:LYS:HD3	1:A:250:LYS:HA	1.75	0.44
1:A:200:PRO:HB2	1:A:248:ARG:HB2	2.00	0.44
1:A:50:GLY:CA	1:A:353:ARG:NH1	2.82	0.43
1:A:261:ARG:NE	1:A:261:ARG:HA	2.33	0.43
1:A:325:SER:CB	1:A:326:PRO:CD	2.94	0.42
1:A:79:PHE:CG	2:B:3:GLC:H62	2.55	0.42
1:A:393:ASN:HB2	1:A:394:PRO:HD2	2.01	0.42
1:A:11:ARG:HH22	1:A:204:ASN:HD21	1.67	0.41
1:A:155:ARG:HB2	1:A:159:GLY:HA2	2.01	0.41
1:A:145:ASN:HD21	1:A:155:ARG:N	2.14	0.41
1:A:122:TYR:HB3	1:A:125:LYS:HD3	2.03	0.41
1:A:11:ARG:NH1	1:A:204:ASN:ND2	2.55	0.41
1:A:24:HIS:CB	1:A:334:ASP:OD1	2.68	0.41
1:A:143:PRO:HB2	1:A:146:TYR:CE1	2.54	0.40
1:A:244:ASP:O	1:A:248:ARG:HG3	2.21	0.40
1:A:130:PRO:CG	1:A:133:GLN:HE21	2.35	0.40
1:A:146:TYR:CB	1:A:147:PRO:CD	2.99	0.40
1:A:190:PHE:O	1:A:216:CYS:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	416/418 (100%)	402 (97%)	13 (3%)	1 (0%)	47 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	157	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	332/332 (100%)	313 (94%)	19 (6%)	20 16

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	TYR
1	A	70	SER
1	A	88	ARG
1	A	95	LEU
1	A	119	ASN
1	A	146	TYR
1	A	162	ASP
1	A	182	ARG
1	A	271	LYS
1	A	277	ASN
1	A	292	ASN
1	A	308	TRP
1	A	317	GLN
1	A	323	LEU
1	A	358	ARG
1	A	372	LEU
1	A	397	VAL
1	A	407	ASN
1	A	417	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	24	HIS
1	A	26	ASN
1	A	81	HIS
1	A	86	ASN
1	A	94	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	119	ASN
1	A	133	GLN
1	A	145	ASN
1	A	167	HIS
1	A	180	ASN
1	A	204	ASN
1	A	240	GLN
1	A	264	ASN
1	A	277	ASN
1	A	292	ASN
1	A	317	GLN
1	A	335	HIS
1	A	366	HIS
1	A	388	ASN
1	A	407	ASN
1	A	410	ASN
1	A	412	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GLC	B	1	2	12,12,12	1.06	0	17,17,17	2.53	5 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GLC	B	2	2	11,11,12	0.80	0	15,15,17	1.74	3 (20%)
2	GLC	B	3	2	11,11,12	0.55	0	15,15,17	1.57	2 (13%)
2	GLC	B	4	2	11,11,12	0.68	0	15,15,17	1.33	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	2/2/19/22	0/1/1/1
2	GLC	B	4	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1	GLC	C1-O5-C5	6.02	125.03	113.66
2	B	1	GLC	O5-C1-C2	5.24	119.64	110.28
2	B	2	GLC	C1-O5-C5	4.31	118.03	112.19
2	B	1	GLC	O2-C2-C1	4.31	119.14	109.16
2	B	3	GLC	O3-C3-C2	-3.00	104.25	109.99
2	B	1	GLC	O1-C1-C2	2.59	116.33	109.03
2	B	2	GLC	O2-C2-C1	2.57	114.42	109.15
2	B	4	GLC	C1-O5-C5	2.52	115.60	112.19
2	B	1	GLC	C1-C2-C3	2.48	115.46	110.31
2	B	3	GLC	C1-O5-C5	2.37	115.40	112.19
2	B	4	GLC	C1-C2-C3	2.30	112.49	109.67
2	B	2	GLC	O5-C5-C4	-2.00	105.96	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	3	GLC	O5-C5-C6-O6
2	B	3	GLC	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	418/418 (100%)	-0.32	3 (0%)	87	87	10, 17, 30, 46	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLN	2.7
1	A	68	ASP	2.6
1	A	146	TYR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GLC	B	1	12/12	0.96	0.15	12,15,16,18	0
2	GLC	B	3	11/12	0.97	0.08	17,18,19,19	0
2	GLC	B	2	11/12	0.98	0.10	14,15,15,16	0
2	GLC	B	4	11/12	0.98	0.07	20,21,22,23	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CA	A	452	1/1	0.99	0.05	18,18,18,18	0
3	CA	A	451	1/1	1.00	0.08	15,15,15,15	0

6.5 Other polymers (i)

There are no such residues in this entry.