



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2022 – 03:04 pm GMT

PDB ID : 7QRN  
Title : Crystal structure of Ovalbumin-related protein X (OVAX) complexed with fondaparinux  
Authors : Coste, F.; Bruneau, G.; Rehault-Godbert, S.  
Deposited on : 2022-01-11  
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	 3% 83% 7% 9%
2	F	5	 100%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3083 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ovalbumin-related protein X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	364	2820	1815	457	528	20	0	4	0

- Molecule 2 is an oligosaccharide called 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-3,6-di-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-methyl 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranoside.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	F	5	91	31	3	49	8	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	123	Total O 123 123	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ovalbumin-related protein X



- Molecule 2: 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-4)-2-deoxy-3,6-di-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-methyl 2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranoside



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.50Å 68.07Å 108.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.09 – 1.83 57.66 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.1 (38.09-1.83) 95.0 (57.66-1.83)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.56 (at 1.83Å)	Xtriage
Refinement program	PHENIX 1.19_4092	Depositor
R, $R_{free}$	0.177 , 0.212 0.177 , 0.211	Depositor DCC
$R_{free}$ test set	1733 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtriage
Anisotropy	0.408	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3083	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SGN, SO4, SUS, ZDO, IDS, EDO, BDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/2889	0.55	0/3924

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2820	0	2732	15	0
2	F	91	0	27	0	0
3	A	25	0	0	0	0
4	A	24	0	36	2	0
5	A	123	0	0	2	0
All	All	3083	0	2795	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:ASN:HB3	1:A:287:THR:HG23	1.89	0.54
4:A:511:EDO:H11	5:A:606:HOH:O	2.08	0.53
1:A:305:LYS:O	4:A:509:EDO:H22	2.11	0.51
1:A:32:HIS:HE1	5:A:622:HOH:O	1.96	0.49
1:A:10:MET:O	1:A:13:ILE:HG22	2.13	0.49
1:A:54:GLY:HA3	1:A:330:ILE:HG13	1.96	0.46
1:A:114[A]:LEU:HD12	1:A:254:LEU:HD11	1.97	0.46
1:A:8:PHE:HE2	1:A:97:LEU:HD21	1.82	0.45
1:A:309:THR:HG23	1:A:321:PHE:CD1	2.51	0.45
1:A:187:LEU:HD11	1:A:336:LEU:HG	1.98	0.44
1:A:331:SER:HB3	1:A:336:LEU:HD22	1.99	0.42
1:A:200:LYS:HA	1:A:250:ALA:HA	2.01	0.42
1:A:117:ALA:HB1	1:A:163:TRP:CH2	2.55	0.41
1:A:246:GLU:HG2	1:A:257:LEU:CD2	2.50	0.41
1:A:172:ILE:HD13	1:A:303:GLU:HG3	2.03	0.41
1:A:70:ASP:N	1:A:70:ASP:OD1	2.54	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	362/402 (90%)	357 (99%)	5 (1%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/355 (84%)	294 (99%)	3 (1%)	76 68

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	209	ARG
1	A	292	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ZDO	F	1	2	20,21,21	3.37	5 (25%)	25,32,32	2.08	4 (16%)
2	IDS	F	2	2	16,16,17	1.82	6 (37%)	17,24,26	1.50	3 (17%)
2	SUS	F	3	2	22,23,24	3.86	8 (36%)	24,36,38	1.21	2 (8%)
2	BDP	F	4	2	12,12,13	1.76	3 (25%)	14,17,19	1.20	1 (7%)
2	SGN	F	5	2	18,19,20	3.71	9 (50%)	22,29,31	1.88	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ZDO	F	1	2	-	0/13/33/33	0/1/1/1
2	IDS	F	2	2	-	0/9/26/29	0/1/1/1
2	SUS	F	3	2	-	1/16/33/36	0/1/1/1
2	BDP	F	4	2	-	0/4/21/24	0/1/1/1
2	SGN	F	5	2	-	5/11/28/31	0/1/1/1

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	3	SUS	S1-N2	11.05	1.74	1.59
2	F	5	SGN	O1S-S1	8.53	1.51	1.42
2	F	3	SUS	O3S-S1	8.46	1.51	1.42
2	F	1	ZDO	OSA-S2	8.28	1.51	1.42
2	F	3	SUS	O1S-S1	8.26	1.51	1.42
2	F	5	SGN	O2S-S1	8.18	1.51	1.42
2	F	1	ZDO	OSB-S2	8.16	1.51	1.42
2	F	5	SGN	S1-N2	7.81	1.70	1.59
2	F	1	ZDO	S2-N2	7.04	1.69	1.59
2	F	1	ZDO	O6-S6	3.63	1.66	1.56
2	F	3	SUS	O6-S2	3.60	1.66	1.56
2	F	3	SUS	O3-S3	3.49	1.67	1.57
2	F	5	SGN	O6-S2	3.32	1.65	1.56
2	F	3	SUS	C3-C2	-3.19	1.48	1.52
2	F	4	BDP	O5-C5	2.97	1.48	1.43
2	F	2	IDS	O2-C2	-2.96	1.42	1.47
2	F	4	BDP	O4-C4	2.71	1.49	1.43
2	F	5	SGN	O3-C3	2.58	1.49	1.43
2	F	5	SGN	O5-C5	2.46	1.48	1.43
2	F	3	SUS	O5-C5	2.44	1.48	1.43
2	F	1	ZDO	C3-C2	-2.42	1.48	1.53
2	F	2	IDS	O3-C3	2.42	1.48	1.43
2	F	5	SGN	C3-C2	-2.41	1.47	1.52
2	F	2	IDS	C3-C2	-2.30	1.48	1.53
2	F	3	SUS	O4-C4	2.23	1.48	1.43
2	F	2	IDS	O2-S	2.16	1.63	1.57
2	F	5	SGN	C4-C3	-2.12	1.46	1.52
2	F	5	SGN	O5-C1	2.12	1.47	1.43
2	F	2	IDS	C4-C3	-2.09	1.47	1.52
2	F	2	IDS	C1-C2	2.09	1.55	1.51

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	4	BDP	C5-C6	-2.06	1.48	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	ZDO	OSA-S2-OSB	-8.13	100.94	120.16
2	F	5	SGN	O1S-S1-O2S	-6.97	103.69	120.16
2	F	2	IDS	C1-C2-C3	3.51	114.64	109.40
2	F	3	SUS	O1S-S1-O3S	-3.03	112.99	120.16
2	F	1	ZDO	C1-C2-N2	-2.67	106.46	110.96
2	F	1	ZDO	OS2-S6-OS1	-2.61	101.73	112.22
2	F	5	SGN	C1-O5-C5	2.51	115.60	112.19
2	F	3	SUS	C1-C2-C3	2.47	112.69	109.17
2	F	4	BDP	C3-C4-C5	2.45	113.44	109.25
2	F	1	ZDO	C4-C3-C2	2.42	113.89	110.34
2	F	5	SGN	C6-C5-C4	-2.13	107.65	112.09
2	F	5	SGN	O6S-S2-O5S	-2.08	101.27	108.49
2	F	2	IDS	C4-C3-C2	2.06	113.92	110.24
2	F	2	IDS	O5-C1-C2	2.00	113.47	109.41

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	5	SGN	C4-C5-C6-O6
2	F	5	SGN	O5-C5-C6-O6
2	F	5	SGN	C6-O6-S2-O5S
2	F	5	SGN	C6-O6-S2-O4S
2	F	3	SUS	C3-O3-S3-O9S
2	F	5	SGN	C2-N2-S1-O1S

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	508	-	3,3,3	0.46	0	2,2,2	0.45	0
3	SO4	A	504	-	4,4,4	0.13	0	6,6,6	0.07	0
4	EDO	A	506	-	3,3,3	0.45	0	2,2,2	0.32	0
3	SO4	A	501	-	4,4,4	0.15	0	6,6,6	0.10	0
4	EDO	A	510	-	3,3,3	0.44	0	2,2,2	0.40	0
4	EDO	A	509	-	3,3,3	0.53	0	2,2,2	0.22	0
4	EDO	A	507	-	3,3,3	0.45	0	2,2,2	0.37	0
3	SO4	A	505	-	4,4,4	0.13	0	6,6,6	0.06	0
3	SO4	A	503	-	4,4,4	0.15	0	6,6,6	0.08	0
4	EDO	A	511	-	3,3,3	0.48	0	2,2,2	0.25	0
3	SO4	A	502	-	4,4,4	0.16	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	508	-	-	1/1/1/1	-
4	EDO	A	506	-	-	0/1/1/1	-
4	EDO	A	510	-	-	0/1/1/1	-
4	EDO	A	509	-	-	1/1/1/1	-
4	EDO	A	507	-	-	1/1/1/1	-
4	EDO	A	511	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	509	EDO	O1-C1-C2-O2
4	A	507	EDO	O1-C1-C2-O2
4	A	508	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	509	EDO	1	0
4	A	511	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	364/402 (90%)	0.12	14 (3%) 40 37	23, 40, 77, 136	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	THR	7.5
1	A	250	ALA	5.5
1	A	203	PHE	3.6
1	A	13	ILE	3.4
1	A	207	ASP	3.4
1	A	8	PHE	3.3
1	A	204	ASN	3.2
1	A	107	ALA	3.1
1	A	199	TRP	2.9
1	A	350	SER	2.7
1	A	112	TYR	2.4
1	A	200	LYS	2.4
1	A	202	ALA	2.2
1	A	285	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BDP	F	4	12/13	0.81	0.12	45,66,75,82	0
2	IDS	F	2	16/17	0.93	0.13	42,48,62,66	0
2	SGN	F	5	19/20	0.93	0.12	51,82,116,119	0
2	SUS	F	3	23/24	0.94	0.13	44,55,66,81	0
2	ZDO	F	1	21/21	0.95	0.14	37,47,70,82	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	504	5/5	0.70	0.31	117,117,118,118	0
3	SO4	A	503	5/5	0.74	0.39	114,114,116,132	0
4	EDO	A	511	4/4	0.78	0.16	61,70,71,72	0
4	EDO	A	509	4/4	0.82	0.23	41,51,55,67	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	A	507	4/4	0.82	0.13	49,55,56,66	0
3	SO4	A	501	5/5	0.83	0.16	68,75,84,89	0
3	SO4	A	505	5/5	0.84	0.36	103,103,104,105	0
4	EDO	A	510	4/4	0.89	0.18	49,53,58,70	0
4	EDO	A	506	4/4	0.91	0.16	50,53,53,57	0
4	EDO	A	508	4/4	0.92	0.08	42,45,52,62	0
3	SO4	A	502	5/5	0.94	0.17	74,79,85,86	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.