Full wwPDB X-ray Structure Validation Report (i)

Aug 17, 2022 – 03:47 PM EDT

PDB ID	:	4QTI
Title	:	Crystal structure of human uPAR in complex with anti-uPAR Fab 8B12
Authors	:	Zhao, B.; Yuan, C.; Luo, Z.; Huang, M.
Deposited on	:	2014-07-08
Resolution	:	3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.29
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	224	2% 83% 13% •
2	L	214	5% 81% 16% • •
3	U	283	6% 73% 16% • 8%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-uPAR antibody, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	218	1651	1043	269	330	9	0	0	0

- Molecule 2 is a protein called anti-uPAR antibody, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	212	1643	1016	278	343	6	0	0	0

- Molecule 3 is a protein called Urokinase plasminogen activator surface receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	U	260	1996	1195	370	397	34	0	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: anti-uPAR antibody, heavy chain

Chain H:



• Molecule 2: anti-uPAR antibody, light chain

Chain L:



• Molecule 3: Urokinase plasminogen activator surface receptor

Chain U:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α, β, γ	45.55Å 95.40Å 189.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.77 – 3.00 42.44 – 3.00	Depositor EDS
% Data completeness (in resolution range)	94.3 (94.77-3.00) 94.3 (42.44-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.50 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.250 , 0.296 0.249 , 0.297	Depositor DCC
R_{free} test set	823 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å^2)	105.3	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 72.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	5290	wwPDB-VP
Average B, all atoms (Å^2)	118.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	H	0.30	0/1694	0.56	0/2316
2	L	0.30	0/1676	0.53	0/2272
3	U	0.30	0/2027	0.55	0/2729
All	All	0.30	0/5397	0.55	0/7317

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1651	0	1599	11	0
2	L	1643	0	1575	19	0
3	U	1996	0	1863	25	0
All	All	5290	0	5037	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:163:TRP:HE1	2:L:175:MET:HE3	1.13	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:163:TRP:HE1	2:L:175:MET:CE	1.68	1.05
2:L:163:TRP:NE1	2:L:175:MET:CE	2.25	0.99
2:L:163:TRP:NE1	2:L:175:MET:HE3	1.82	0.90
2:L:163:TRP:NE1	2:L:175:MET:HE2	1.91	0.83
3:U:190:ASN:ND2	3:U:217:GLY:O	2.26	0.67
1:H:134:ALA:O	1:H:222:ARG:NH1	2.30	0.65
2:L:194:CYS:SG	2:L:194:CYS:O	2.55	0.64
3:U:55:LEU:HD21	3:U:144:LEU:HD23	1.78	0.64
3:U:187:LEU:O	3:U:216:ARG:NH1	2.34	0.61
1:H:73:ASP:OD1	1:H:75:SER:OG	2.13	0.59
2:L:169:LYS:HA	2:L:169:LYS:HE3	1.86	0.57
3:U:55:LEU:HD13	3:U:123:LEU:HD12	1.90	0.54
1:H:145:VAL:O	1:H:145:VAL:HG13	2.08	0.53
3:U:177:ASN:HD22	3:U:177:ASN:H	1.56	0.53
2:L:66:ARG:HG3	2:L:67:SER:N	2.22	0.53
3:U:225:ALA:CB	3:U:252:LEU:HD13	2.40	0.52
2:L:167:ASP:O	2:L:171:SER:HA	2.10	0.51
2:L:163:TRP:CD1	2:L:175:MET:CE	2.94	0.51
3:U:189:GLN:HE21	3:U:189:GLN:N	2.09	0.50
2:L:1:ASP:O	2:L:3:GLN:NE2	2.45	0.49
2:L:35:TRP:CE2	2:L:73:LEU:HB2	2.49	0.48
3:U:1:LEU:HD12	3:U:22:ASP:HA	1.95	0.48
2:L:163:TRP:CD1	2:L:175:MET:HE3	2.48	0.47
2:L:189:HIS:O	2:L:211:ARG:HD3	2.16	0.46
3:U:153:CYS:O	3:U:155:GLY:N	2.49	0.45
3:U:160:HIS:CE1	3:U:241:CYS:HB2	2.51	0.45
3:U:30:ARG:C	3:U:31:LEU:HD22	2.36	0.45
2:L:18:ARG:HA	2:L:76:SER:O	2.17	0.45
1:H:128:PRO:HB3	1:H:151:VAL:HG13	1.97	0.44
1:H:100:PHE:HB2	1:H:107:LEU:HB3	2.00	0.44
2:L:147:LYS:N	2:L:195:GLU:O	2.49	0.44
3:U:129:TRP:HA	3:U:130:ILE:HA	1.76	0.44
3:U:150:LEU:O	3:U:153:CYS:HB2	2.18	0.44
1:H:41:PRO:O	1:H:43:HIS:N	2.51	0.43
3:U:96:ILE:H	3:U:177:ASN:HD21	1.65	0.43
2:L:8:PRO:O	2:L:102:THR:HG23	2.18	0.43
3:U:273:HIS:CG	3:U:274:PRO:HD2	2.53	0.43
2:L:91:TYR:HA	2:L:96:TYR:CD1	2.53	0.43
1:H:41:PRO:C	1:H:43:HIS:H	2.22	0.42
1:H:186:LEU:C	1:H:186:LEU:HD12	2.40	0.42
3:U:157:ASN:HB3	3:U:245:SER:OG	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:50:GLU:OE2	1:H:59:ASN:ND2	2.51	0.42
3:U:153:CYS:N	3:U:154:PRO:CD	2.82	0.42
3:U:91:ARG:HG2	3:U:114:GLN:HG2	2.02	0.41
3:U:91:ARG:CG	3:U:114:GLN:HG2	2.51	0.41
3:U:126:VAL:HG13	3:U:143:HIS:CE1	2.56	0.41
3:U:228:THR:HA	3:U:235:SER:HA	2.02	0.41
1:H:41:PRO:C	1:H:43:HIS:N	2.73	0.41
3:U:96:ILE:HG23	3:U:110:HIS:HB3	2.03	0.40
2:L:163:TRP:CD1	2:L:175:MET:HG3	2.56	0.40
3:U:177:ASN:HD22	3:U:177:ASN:N	2.17	0.40
1:H:196:THR:O	1:H:196:THR:CG2	2.68	0.40
3:U:24:CYS:HB2	3:U:76:CYS:SG	2.61	0.40
3:U:225:ALA:HB2	3:U:252:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	214/224 (96%)	200 (94%)	11 (5%)	3 (1%)	11 43
2	L	210/214 (98%)	198 (94%)	10 (5%)	2 (1%)	15 53
3	U	254/283 (90%)	226 (89%)	25 (10%)	3 (1%)	13 48
All	All	678/721 (94%)	624 (92%)	46 (7%)	8 (1%)	13 48

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	104	ARG
1	H	170	SER
2	L	143	ASP
2	L	183	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	U	202	THR
3	U	154	PRO
1	H	181	SER
3	U	129	TRP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	189/192 (98%)	175 (93%)	14 (7%)	13	44
2	L	190/192 (99%)	172 (90%)	18 (10%)	8	32
3	U	232/250 (93%)	212 (91%)	20 (9%)	10	37
All	All	611/634 (96%)	559 (92%)	52 (8%)	10	38

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	1	GLN
1	H	11	LEU
1	H	12	MET
1	H	13	LYS
1	H	58	THR
1	H	87	THR
1	H	106	ASN
1	H	107	LEU
1	H	111	VAL
1	H	146	THR
1	H	160	THR
1	H	189	SER
1	H	203	THR
1	H	218	LYS
2	L	7	SER
2	L	18	ARG
2	L	21	LEU
2	L	24	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	55	ASP
2	L	66	ARG
2	L	84	VAL
2	L	105	GLU
2	L	107	LYS
2	L	110	ASP
2	L	156	GLN
2	L	163	TRP
2	L	169	LYS
2	L	180	THR
2	L	182	THR
2	L	200	THR
2	L	206	VAL
2	L	209	PHE
3	U	1	LEU
3	U	2	ARG
3	U	5	GLN
3	U	13	ARG
3	U	69	VAL
3	U	71	CYS
3	U	91	ARG
3	U	93	LEU
3	U	105	CYS
3	U	113	LEU
3	U	116	ARG
3	U	125	VAL
3	U	130	ILE
3	U	142	ARG
3	U	177	ASN
3	U	189	GLN
3	U	206	SER
3	U	209	GLU
3	U	245	SER
3	U	267	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	1	GLN
1	H	5	GLN
1	H	77	ASN
1	H	106	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	3	GLN
3	U	160	HIS
3	U	177	ASN
3	U	189	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	H	218/224 (97%)	-0.20	4 (1%)	68	40	72, 98, 127, 153	16 (7%)
2	L	212/214 (99%)	0.08	10 (4%)	31	11	73, 109, 159, 174	14 (6%)
3	U	260/283 (91%)	0.12	17 (6%)	18	5	87, 131, 183, 222	0
All	All	690/721 (95%)	0.00	31 (4%)	33	12	72, 111, 167, 222	30 (4%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	U	249	HIS	6.9
3	U	34	GLU	6.7
3	U	251	HIS	6.5
3	U	260	HIS	5.4
3	U	19	LEU	4.9
2	L	194	CYS	4.8
3	U	182	LEU	4.4
2	L	208	SER	4.0
3	U	40	LEU	3.8
3	U	252	LEU	3.8
2	L	154	GLU	3.6
3	U	253	GLY	3.6
3	U	35	GLY	3.6
2	L	158	GLY	3.4
2	L	209	PHE	3.3
1	H	147	LEU	3.2
3	U	32	TRP	2.9
1	H	149	CYS	2.9
2	L	212	ASN	2.8
3	U	20	GLY	2.6
2	L	200	THR	2.6
3	U	256	PHE	2.5
1	H	42	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	192	TYR	2.4
3	U	47	HIS	2.3
3	U	41	VAL	2.2
1	H	187	SER	2.2
3	U	250	ALA	2.0
2	L	201	SER	2.0
3	U	2	ARG	2.0
2	L	175	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.