



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2021 – 02:45 AM EDT

PDB ID : 2QWR
Title : Crystal structure of disulfide-bond-crosslinked complex of bovine hsc70 (1-394aa)R171C and bovine Auxilin (810-910aa)D876C in the AMPPNP intact form
Authors : Jiang, J.; Maes, E.G.; Wang, L.; Taylor, A.B.; Hinck, A.P.; Lafer, E.M.; Sousa, R.
Deposited on : 2007-08-10
Resolution : 2.21 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	394	 78% 18% . .
2	B	92	 78% 20% .

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heat shock cognate 71 kDa protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	383	2959	1856	517	577	9	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	171	CYS	ARG	engineered mutation	UNP P19120

- Molecule 2 is a protein called Putative tyrosine-protein phosphatase auxilin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	92	754	488	126	135	5	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	876	CYS	ASP	engineered mutation	UNP Q27974

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	31	10	6	12	3	0	0

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	3	2	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 5 3 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	98	Total O 98 98	0	0
6	B	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Heat shock cognate 71 kDa protein



- Molecule 2: Putative tyrosine-protein phosphatase auxilin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.80Å 56.49Å 225.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.67 – 2.21 31.67 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.7 (31.67-2.21) 99.7 (31.67-2.21)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.230 , 0.294 0.282 , 0.314	Depositor DCC
R_{free} test set	1310 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtrriage
Anisotropy	0.418	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3863	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, ACY, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.69	1/3005 (0.0%)	0.74	2/4058 (0.0%)
2	B	0.49	0/773	0.57	0/1046
All	All	0.66	1/3778 (0.0%)	0.71	2/5104 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	340	SER	CB-OG	5.01	1.48	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	ASP	CB-CG-OD1	6.14	123.83	118.30
1	A	228	LEU	CA-CB-CG	5.84	128.73	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2959	0	2955	44	0
2	B	754	0	760	13	0
3	A	31	0	13	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	3	0	0	0	0
5	A	11	0	13	1	0
6	A	98	0	0	3	0
6	B	7	0	0	0	0
All	All	3863	0	3741	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ASN:O	1:A:260:VAL:HG23	1.82	0.79
1:A:233:PHE:HA	1:A:306:ASN:HD21	1.47	0.78
1:A:151:ASN:H	1:A:154:GLN:HE21	1.34	0.75
1:A:255:GLU:HA	1:A:255:GLU:OE1	1.89	0.73
2:B:823:ILE:HG12	2:B:893:GLU:HG2	1.73	0.71
1:A:233:PHE:HA	1:A:306:ASN:ND2	2.10	0.66
1:A:128:LYS:HE2	1:A:139:VAL:O	1.96	0.65
1:A:58:GLN:HE22	1:A:62:ASN:HD22	1.43	0.65
1:A:358:GLU:HG3	6:A:3213:HOH:O	1.99	0.62
1:A:151:ASN:H	1:A:154:GLN:NE2	1.98	0.60
2:B:828:ARG:HG2	2:B:897:ALA:HA	1.83	0.59
1:A:258:ARG:O	1:A:262:ARG:HG3	2.03	0.58
1:A:256:ASN:HB3	1:A:259:ALA:HB3	1.85	0.56
1:A:56:LYS:O	1:A:59:VAL:HG12	2.06	0.55
2:B:874:HIS:HB3	2:B:877:LYS:HD3	1.89	0.55
1:A:176:PRO:HB2	1:A:197:ILE:HG13	1.89	0.55
1:A:283:GLU:HG2	1:A:294:TYR:CD1	2.41	0.55
2:B:877:LYS:H	2:B:877:LYS:CD	2.22	0.53
1:A:197:ILE:HD13	1:A:335:VAL:HB	1.92	0.52
2:B:877:LYS:CD	2:B:877:LYS:N	2.73	0.51
1:A:315:ASP:HB2	1:A:316:PRO:HD3	1.92	0.50
1:A:175:GLU:HG2	1:A:369:VAL:HG11	1.93	0.50
1:A:112:LYS:HE2	6:A:3230:HOH:O	2.11	0.50
2:B:831:ARG:HG2	2:B:856:LEU:HB3	1.95	0.49
1:A:118:GLU:O	1:A:122:MET:HG3	2.13	0.49
1:A:236:ARG:HH22	1:A:308:ASP:CG	2.17	0.48
1:A:252:ASP:OD1	1:A:254:SER:HB2	2.13	0.48
1:A:255:GLU:OE1	1:A:255:GLU:CA	2.62	0.48
1:A:338:GLY:HA2	3:A:487:ANP:O1A	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:877:LYS:N	2:B:877:LYS:HD2	2.30	0.47
1:A:178:ALA:O	1:A:372:GLY:HA3	2.15	0.47
2:B:847:LYS:HA	2:B:847:LYS:NZ	2.31	0.46
1:A:363:ILE:O	1:A:364:ASN:C	2.55	0.46
1:A:195:VAL:HG21	1:A:212:ILE:HD11	1.99	0.45
1:A:102:LYS:HE3	1:A:115:TYR:CZ	2.52	0.45
6:A:3202:HOH:O	2:B:877:LYS:HE3	2.17	0.45
1:A:128:LYS:HG3	1:A:167:LEU:HD21	2.00	0.44
2:B:877:LYS:HD3	2:B:877:LYS:H	1.83	0.43
2:B:826:LYS:HE3	2:B:836:THR:HG21	2.01	0.43
1:A:366:ASP:OD1	1:A:366:ASP:N	2.47	0.43
1:A:71:LYS:HG2	5:A:3147:GOL:H32	2.00	0.43
1:A:253:ILE:HG22	1:A:288:TYR:CG	2.53	0.43
1:A:150:PHE:CD2	1:A:154:GLN:HB3	2.54	0.42
1:A:271:LYS:HG3	1:A:302:PHE:CZ	2.54	0.42
1:A:336:LEU:HD22	1:A:341:THR:HB	2.01	0.42
1:A:103:VAL:O	1:A:113:SER:HA	2.19	0.41
1:A:275:SER:HA	1:A:344:PRO:HD3	2.02	0.41
2:B:818:LYS:NZ	2:B:839:THR:O	2.48	0.41
1:A:195:VAL:HG21	1:A:212:ILE:CD1	2.51	0.41
1:A:295:THR:OG1	1:A:296:SER:N	2.53	0.41
1:A:5:PRO:HD3	1:A:137:LYS:HE3	2.02	0.41
1:A:264:ARG:HH11	1:A:264:ARG:HD2	1.72	0.40
1:A:336:LEU:HD11	1:A:347:GLN:HG2	2.03	0.40
2:B:819:ILE:CD1	2:B:886:TYR:HB3	2.52	0.40
1:A:288:TYR:O	1:A:289:GLU:C	2.59	0.40
1:A:342:ARG:HH11	1:A:342:ARG:HD3	1.75	0.40
1:A:378:ALA:HB1	1:A:383:ASP:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/394 (97%)	364 (96%)	17 (4%)	0	100	100
2	B	90/92 (98%)	85 (94%)	5 (6%)	0	100	100
All	All	471/486 (97%)	449 (95%)	22 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/328 (97%)	304 (96%)	13 (4%)	30	37
2	B	81/81 (100%)	74 (91%)	7 (9%)	10	9
All	All	398/409 (97%)	378 (95%)	20 (5%)	24	28

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	THR
1	A	58	GLN
1	A	85	SER
1	A	89	HIS
1	A	104	GLN
1	A	228	LEU
1	A	254	SER
1	A	255	GLU
1	A	258	ARG
1	A	279	GLN
1	A	281	SER
1	A	319	LYS
1	A	385	SER
2	B	815	GLU
2	B	824	GLU
2	B	829	ASN
2	B	845	GLU
2	B	847	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	851	VAL
2	B	877	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	58	GLN
1	A	104	GLN
1	A	154	GLN
1	A	306	ASN
1	A	351	GLN
1	A	376	GLN
2	B	829	ASN
2	B	874	HIS
2	B	895	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	3148	-	3,4,5	0.12	0	1,4,5	0.28	0
5	GOL	A	3147	-	5,5,5	0.38	0	5,5,5	0.84	0
3	ANP	A	487	-	29,33,33	1.83	5 (17%)	31,52,52	2.57	8 (25%)
4	ACY	A	1022	-	2,2,3	1.25	0	1,1,3	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	3148	-	-	2/2/2/4	-
5	GOL	A	3147	-	-	2/4/4/4	-
3	ANP	A	487	-	-	2/14/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	487	ANP	PG-O1G	6.16	1.55	1.46
3	A	487	ANP	PG-N3B	3.72	1.73	1.63
3	A	487	ANP	PB-O1B	3.26	1.51	1.46
3	A	487	ANP	O4'-C1'	3.20	1.45	1.41
3	A	487	ANP	PG-O3G	2.10	1.62	1.56

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	487	ANP	O1B-PB-N3B	-7.17	101.22	111.77
3	A	487	ANP	O3G-PG-O2G	6.00	123.61	107.64
3	A	487	ANP	N3-C2-N1	-5.56	119.99	128.68
3	A	487	ANP	O2G-PG-O1G	-5.31	100.09	113.45
3	A	487	ANP	O2B-PB-O1B	3.48	117.21	109.92
3	A	487	ANP	O3A-PB-N3B	2.48	113.46	106.59
3	A	487	ANP	C4-C5-N7	-2.43	106.87	109.40
3	A	487	ANP	O2A-PA-O1A	2.22	123.24	112.24

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	487	ANP	PA-O3A-PB-O1B
3	A	487	ANP	PA-O3A-PB-O2B
5	A	3147	GOL	C1-C2-C3-O3
5	A	3148	GOL	C1-C2-C3-O3
5	A	3148	GOL	O2-C2-C3-O3
5	A	3147	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	3147	GOL	1	0
3	A	487	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.