



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 01:56 pm BST

PDB ID : 2QWX  
Title : Crystal Structure of Quinone Reductase II  
Authors : Calamini, B.; Santarsiero, B.D.; Boutin, J.A.; Mesecar, A.D.  
Deposited on : 2007-08-10  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	B	231	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ribosyldihyronicotinamide dehydrogenase [quinone].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	230	Total 1891	C 1216	N 315	O 351	S 9	0	8	0
1	B	230	Total 1896	C 1219	N 315	O 352	S 10	0	9	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Zn 1	0	0
2	A	1	Total 1	Zn 1	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	53	27	9	15	2	0	0
3	B	1	53	27	9	15	2	0	0

- Molecule 4 is N-[2-(5-methoxy-1H-indol-3-yl)ethyl]acetamide (three-letter code: ML1) (formula: C<sub>13</sub>H<sub>16</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	17	13	2	2	0	0
4	B	1	17	13	2	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	259	259	259	0	0
5	B	220	220	220	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosyldihyronicotinamide dehydrogenase [quinone]



- Molecule 1: Ribosyldihyronicotinamide dehydrogenase [quinone]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.63Å 83.55Å 106.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.50 46.88 – 1.47	Depositor EDS
% Data completeness (in resolution range)	98.4 (20.00-1.50) 97.9 (46.88-1.47)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 1.47Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.192 , 0.213 0.187 , 0.209	Depositor DCC
$R_{free}$ test set	4226 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.464	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4408	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ML1, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/1942	0.81	4/2631 (0.2%)
1	B	0.60	0/1947	0.76	0/2638
All	All	0.60	0/3889	0.79	4/5269 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	49	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	A	49	ARG	NE-CZ-NH2	-7.35	116.62	120.30
1	A	182[A]	LYS	N-CA-C	-5.47	96.23	111.00
1	A	182[B]	LYS	N-CA-C	-5.47	96.23	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1891	0	1839	39	0
1	B	1896	0	1841	42	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	53	0	31	5	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	53	0	31	5	0
4	A	17	0	16	5	0
4	B	17	0	16	4	0
5	A	259	0	0	12	0
5	B	220	0	0	14	0
All	All	4408	0	3774	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:LYS:HE2	1:A:53:LYS:HA	1.43	0.98
1:A:151:THR:H	1:A:154:MET:HE3	1.28	0.96
1:A:17:PHE:HB2	3:A:232:FAD:H51A	1.55	0.86
1:B:127:ASP:HB3	5:B:373:HOH:O	1.76	0.85
1:B:4[A]:LYS:HE2	5:B:438:HOH:O	1.81	0.79
1:B:4[B]:LYS:HE3	5:B:425:HOH:O	1.83	0.79
1:B:201:LYS:HD3	3:B:232:FAD:H61A	1.50	0.77
1:B:45[A]:ASN:OD1	5:B:326:HOH:O	2.08	0.72
1:B:17:PHE:HB2	3:B:232:FAD:H52A	1.72	0.71
1:B:199:GLU:O	1:B:203[A]:MET:HG3	1.91	0.70
1:A:216:LYS:HG2	5:A:307:HOH:O	1.91	0.70
1:A:53:LYS:HE2	1:A:53:LYS:CA	2.21	0.70
1:A:85[A]:THR:HG22	5:A:300:HOH:O	1.92	0.68
1:A:63:GLU:HG2	5:A:492:HOH:O	1.94	0.68
1:A:216:LYS:HE2	5:A:316:HOH:O	1.93	0.68
1:B:218:GLU:HB2	5:B:444:HOH:O	1.94	0.67
1:B:151:THR:H	1:B:154:MET:HE3	1.60	0.66
1:B:196:SER:OG	1:B:199:GLU:HG3	1.96	0.65
1:A:4[B]:LYS:HD3	5:A:482:HOH:O	2.01	0.60
1:B:151:THR:HG23	1:B:154:MET:HE3	1.83	0.60
1:A:24:VAL:HG11	1:A:204:VAL:CG1	2.31	0.59
4:A:233:ML1:H31	1:B:106[B]:PHE:CE1	2.37	0.58
1:A:4[A]:LYS:HD2	5:A:424:HOH:O	2.02	0.58
1:A:17:PHE:HB2	3:A:232:FAD:C5B	2.31	0.58
1:B:24[B]:VAL:HG22	5:B:329:HOH:O	2.03	0.57
1:A:151:THR:N	1:A:154:MET:HE3	2.11	0.57
1:A:141:LEU:HD23	1:A:182[B]:LYS:HB2	1.87	0.56
4:B:233:ML1:H51	5:B:449:HOH:O	2.06	0.56

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:GLU:OE2	3:B:232:FAD:H8A	2.05	0.56
1:B:68:GLY:HA3	4:B:233:ML1:H31	1.86	0.55
1:B:17:PHE:HB2	3:B:232:FAD:C5B	2.37	0.55
1:A:85[B]:THR:HG21	5:A:279:HOH:O	2.07	0.54
1:B:182:LYS:HG2	5:B:420:HOH:O	2.07	0.54
4:A:233:ML1:H62	1:B:149:GLY:HA3	1.89	0.54
1:A:85[B]:THR:HG23	5:A:300:HOH:O	2.08	0.54
1:B:4[A]:LYS:HG2	1:B:35:THR:HB	1.89	0.53
1:A:154:MET:SD	4:B:233:ML1:H52	2.49	0.52
1:A:200:ARG:HD2	3:A:232:FAD:O4B	2.09	0.51
1:A:82:SER:HA	1:A:85[B]:THR:HG22	1.91	0.51
1:B:4[B]:LYS:HG2	1:B:35:THR:HB	1.94	0.50
1:B:216:LYS:N	1:B:216:LYS:HD2	2.27	0.50
1:B:4[B]:LYS:CE	5:B:425:HOH:O	2.51	0.50
1:A:151:THR:HG23	1:A:154:MET:HE3	1.95	0.49
1:B:106[B]:PHE:HE2	1:B:155:TYR:CE1	2.30	0.49
1:A:230:GLN:OE1	1:A:230:GLN:C	2.51	0.48
1:B:3:LYS:NZ	5:B:427:HOH:O	2.46	0.47
1:A:165:ARG:HA	1:A:168:LEU:HD12	1.97	0.47
1:B:194:ILE:HG13	5:B:387:HOH:O	2.14	0.46
1:B:230:GLN:O	1:B:230:GLN:HG3	2.16	0.46
1:B:126:PHE:CZ	4:B:233:ML1:C11	2.99	0.46
1:A:53:LYS:CE	1:A:53:LYS:CA	2.91	0.46
1:A:52:ASP:OD1	1:A:53:LYS:HE3	2.16	0.46
1:B:169:TRP:HB3	1:B:170:PRO:HD3	1.97	0.45
1:A:82:SER:HA	1:A:85[B]:THR:CG2	2.47	0.45
1:A:216:LYS:HG2	1:A:216:LYS:O	2.16	0.44
1:B:138:GLN:HA	1:B:180:GLY:O	2.17	0.44
1:A:216:LYS:HE3	5:A:307:HOH:O	2.17	0.44
1:A:73[B]:GLU:OE1	5:A:475:HOH:O	2.21	0.44
4:A:233:ML1:H31	1:B:106[B]:PHE:CZ	2.53	0.43
3:A:232:FAD:HM72	5:B:288:HOH:O	2.17	0.43
1:A:170:PRO:HB3	1:B:106[A]:PHE:HB3	2.01	0.43
1:B:198:GLU:H	1:B:198:GLU:CD	2.22	0.43
1:A:101:PHE:CZ	1:A:146:VAL:HG22	2.54	0.43
1:A:216:LYS:CE	5:A:316:HOH:O	2.61	0.43
1:A:178:PHE:CE1	4:A:233:ML1:H32	2.53	0.43
1:A:169:TRP:HB3	1:A:170:PRO:HD3	2.01	0.42
4:A:233:ML1:H53	1:B:154:MET:CE	2.48	0.42
1:A:4[B]:LYS:HG2	1:A:35:THR:HB	2.02	0.42
1:A:20:SER:O	1:A:24:VAL:HG13	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:LYS:HD3	3:B:232:FAD:N6A	2.26	0.42
1:B:189:SER:HA	1:B:203[A]:MET:HE2	2.02	0.42
1:B:143:LEU:C	1:B:143:LEU:HD23	2.40	0.41
1:B:20:SER:O	1:B:24[A]:VAL:HG22	2.21	0.41
1:A:104:TYR:HA	3:A:232:FAD:C5X	2.50	0.41
1:B:195:ALA:HB1	1:B:199:GLU:HB2	2.01	0.41
1:B:4[B]:LYS:HE3	5:B:438:HOH:O	2.20	0.41
1:B:167:PHE:O	1:B:170:PRO:HD2	2.21	0.40
1:A:151:THR:HG23	1:A:154:MET:CE	2.50	0.40
1:A:216:LYS:CE	5:A:307:HOH:O	2.69	0.40
1:A:64:VAL:HG22	1:B:15:LYS:HD2	2.03	0.40
1:B:4[A]:LYS:HD2	5:B:372:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/231 (102%)	227 (96%)	9 (4%)	0	100	100
1	B	237/231 (103%)	225 (95%)	11 (5%)	1 (0%)	34	13
All	All	473/462 (102%)	452 (96%)	20 (4%)	1 (0%)	47	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	229	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/195 (104%)	201 (100%)	1 (0%)	88	78
1	B	203/195 (104%)	201 (99%)	2 (1%)	76	57
All	All	405/390 (104%)	402 (99%)	3 (1%)	81	69

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	230	GLN
1	B	129	PRO
1	B	153	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	212	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ML1	B	233	-	17,18,18	1.14	3 (17%)	18,24,24	1.02	1 (5%)
4	ML1	A	233	-	17,18,18	0.91	1 (5%)	18,24,24	1.06	2 (11%)
3	FAD	A	232	-	51,58,58	2.10	9 (17%)	60,89,89	2.03	5 (8%)
3	FAD	B	232	-	51,58,58	2.15	10 (19%)	60,89,89	1.92	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ML1	B	233	-	-	3/8/8/8	0/2/2/2
4	ML1	A	233	-	-	2/8/8/8	0/2/2/2
3	FAD	A	232	-	-	1/30/50/50	0/6/6/6
3	FAD	B	232	-	-	6/30/50/50	0/6/6/6

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	232	FAD	C4X-N5	7.84	1.44	1.33
3	B	232	FAD	C4X-N5	7.67	1.44	1.33
3	B	232	FAD	C10-N1	6.25	1.41	1.33
3	B	232	FAD	C4X-C10	5.20	1.44	1.38
3	B	232	FAD	C4-N3	5.17	1.42	1.33
3	A	232	FAD	C9A-N10	5.16	1.45	1.38
3	A	232	FAD	C10-N1	5.07	1.39	1.33
3	A	232	FAD	C4X-C10	4.84	1.43	1.38
3	B	232	FAD	C9A-N10	4.50	1.44	1.38
3	B	232	FAD	C4A-N3A	4.23	1.41	1.35
3	A	232	FAD	C4-N3	4.23	1.40	1.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	232	FAD	C4A-N3A	3.77	1.40	1.35
3	A	232	FAD	C5X-N5	3.15	1.40	1.35
4	B	233	ML1	C11-C10	2.59	1.43	1.38
3	B	232	FAD	C9-C8	2.58	1.44	1.37
3	B	232	FAD	O4B-C1B	2.41	1.44	1.41
4	B	233	ML1	C9-C10	2.37	1.41	1.37
3	A	232	FAD	C9-C8	2.22	1.43	1.37
3	A	232	FAD	C5A-C4A	-2.14	1.35	1.40
4	B	233	ML1	C12-C11	2.12	1.41	1.36
3	B	232	FAD	C2A-N1A	2.09	1.37	1.33
4	A	233	ML1	C9-C10	2.09	1.40	1.37
3	B	232	FAD	C5X-N5	2.04	1.38	1.35

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	232	FAD	C4-N3-C2	11.99	125.27	115.14
3	B	232	FAD	C4-N3-C2	11.38	124.75	115.14
3	B	232	FAD	C4X-C4-N3	-5.91	115.34	123.43
3	A	232	FAD	C4X-C4-N3	-5.87	115.41	123.43
3	B	232	FAD	C1'-N10-C9A	3.66	121.17	118.29
3	A	232	FAD	C1'-N10-C9A	3.23	120.83	118.29
4	B	233	ML1	C11-C12-C13	-2.85	117.25	120.84
3	A	232	FAD	C4-C4X-C10	-2.80	118.10	119.95
4	A	233	ML1	C11-C12-C13	-2.59	117.57	120.84
4	A	233	ML1	C1-C6-C7	2.22	115.58	111.28
3	A	232	FAD	C9A-N10-C10	-2.06	119.21	121.91

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	233	ML1	C5-C4-N1-C1
4	B	233	ML1	C1-C6-C7-C8
3	B	232	FAD	C5B-O5B-PA-O1A
3	B	232	FAD	C5B-O5B-PA-O2A
3	B	232	FAD	O4B-C4B-C5B-O5B
3	B	232	FAD	C3B-C4B-C5B-O5B
4	A	233	ML1	C5-C4-N1-C1
4	A	233	ML1	O2-C4-N1-C1
4	B	233	ML1	O2-C4-N1-C1
3	A	232	FAD	O4B-C4B-C5B-O5B

Continued on next page...

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	B	232	FAD	C5B-O5B-PA-O3P
3	B	232	FAD	C4'-C5'-O5'-P


There are no ring outliers.

4 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	233	ML1	4	0
4	A	233	ML1	5	0
3	A	232	FAD	5	0
3	B	232	FAD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/231 (99%)	0.09	11 (4%) 30 33	11, 19, 40, 53	0
1	B	230/231 (99%)	0.47	20 (8%) 10 10	12, 23, 46, 61	0
All	All	460/462 (99%)	0.28	31 (6%) 17 19	11, 20, 44, 61	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	8.0
1	B	230	GLN	6.5
1	A	230	GLN	6.1
1	B	229	GLY	5.7
1	B	194	ILE	5.3
1	A	1	ALA	5.3
1	B	225	HIS	5.0
1	A	198	GLU	4.8
1	B	198	GLU	4.6
1	A	194	ILE	4.4
1	B	57	GLY	3.9
1	A	2	GLY	3.8
1	B	197	GLU	3.7
1	B	159	GLY	3.6
1	B	58	THR	3.2
1	A	197	GLU	3.2
1	A	200	ARG	3.1
1	B	158	THR	2.9
1	B	62	PRO	2.9
1	B	59	LEU	2.8
1	B	228	PHE	2.7
1	A	153	GLU	2.7
1	B	166	TYR	2.7
1	B	150	GLY	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	157	LYS	2.5
1	A	201	LYS	2.5
1	A	216	LYS	2.3
1	B	2	GLY	2.1
1	A	63	GLU	2.1
1	B	195	ALA	2.1
1	B	153	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ML1	A	233	17/17	0.63	0.27	56,57,61,62	0
4	ML1	B	233	17/17	0.74	0.27	55,56,62,63	0
3	FAD	B	232	53/53	0.87	0.18	17,25,60,60	0
3	FAD	A	232	53/53	0.90	0.15	19,25,52,52	0
2	ZN	B	231	1/1	0.94	0.04	35,35,35,35	0
2	ZN	A	231	1/1	0.99	0.03	23,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FAD B 232:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD A 232:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.