



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 2, 2023 – 01:29 PM EDT

PDB ID : 3QX8  
Title : Crystal structure of MID domain from hAGO2 in complex with m7GpppG  
Authors : Frank, F.; Fabian, M.R.; Stepinski, J.; Jemielity, J.; Darzynkiewicz, E.; Sonnenberg, N.; Nagar, B.  
Deposited on : 2011-03-01  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	
1	B	138	
1	C	138	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein argonaute-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	133	Total 1036	C 659	N 185	O 182	S 10	0	0	0
1	B	133	Total 1036	C 659	N 185	O 182	S 10	0	0	0
1	C	133	Total 1036	C 659	N 185	O 182	S 10	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	438	SER	-	expression tag	UNP Q9UKV8
B	438	SER	-	expression tag	UNP Q9UKV8
C	438	SER	-	expression tag	UNP Q9UKV8

- Molecule 2 is 7-METHYL-GUANOSINE-5'-TRIPHOSPHATE-5'-GUANOSINE (three-letter code: GTG) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>10</sub>O<sub>18</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	23	0
			52	21	10	18	3		
2	B	1	Total	C	N	O	P	23	0
			52	21	10	18	3		
2	C	1	Total	C	N	O	P	23	0
			52	21	10	18	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	37	Total	O	0	0
			37	37		
3	B	32	Total	O	0	0
			32	32		
3	C	25	Total	O	0	0
			25	25		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein argonaute-2



- Molecule 1: Protein argonaute-2



- Molecule 1: Protein argonaute-2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.69Å 46.97Å 66.12Å 86.09° 73.33° 83.38°	Depositor
Resolution (Å)	30.00 – 2.30 38.22 – 1.69	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.30) 83.4 (38.22-1.69)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.00 (at 1.69Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.226 , 0.279 0.211 , 0.263	Depositor DCC
$R_{free}$ test set	2776 reflections (5.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.1	Xtrriage
Anisotropy	0.314	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,-h+l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3358	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1055	0.62	0/1425
1	B	0.39	0/1055	0.62	0/1425
1	C	0.38	0/1055	0.63	0/1425
All	All	0.39	0/3165	0.62	0/4275

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1036	0	1077	21	0
1	B	1036	0	1077	41	0
1	C	1036	0	1077	23	0
2	A	52	0	26	5	0
2	B	52	0	26	16	0
2	C	52	0	26	2	0
3	A	37	0	0	0	0
3	B	32	0	0	0	0
3	C	25	0	0	0	0
All	All	3358	0	3309	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:524:GLY:O	2:B:28:GTG:HC71	1.40	1.18
1:B:522:LEU:HB2	2:B:28:GTG:HC72	1.30	1.12
1:B:522:LEU:CB	2:B:28:GTG:HC72	1.91	1.00
1:A:440:LYS:HD3	1:A:441:GLN:H	1.24	0.96
1:A:440:LYS:HD3	1:A:441:GLN:N	1.83	0.93
1:B:524:GLY:O	2:B:28:GTG:C7X	2.21	0.87
1:B:498:ALA:HB1	1:B:531:GLU:HG2	1.70	0.74
1:B:459:GLN:HB3	1:B:494:TYR:CZ	2.27	0.69
1:C:521:ILE:HD12	1:C:552:VAL:HG21	1.74	0.68
1:B:524:GLY:CA	2:B:28:GTG:HC73	2.23	0.68
1:B:502:GLU:HB3	1:B:503:PRO:HD3	1.76	0.66
1:A:526:THR:H	2:A:28:GTG:HC71	1.62	0.65
1:B:531:GLU:HA	1:B:531:GLU:OE1	1.96	0.64
1:C:521:ILE:HD13	1:C:547:VAL:HG13	1.80	0.64
1:C:448:ILE:HD12	1:C:485:ILE:HG12	1.80	0.63
1:B:526:THR:N	2:B:28:GTG:O6A	2.31	0.63
1:C:544:THR:O	1:C:570:LYS:HE2	1.99	0.62
1:A:475:ARG:HD3	1:A:487:GLY:HA3	1.81	0.61
1:A:525:LYS:HA	2:A:28:GTG:HC73	1.82	0.61
1:B:450:VAL:HG12	1:B:515:LEU:HA	1.81	0.61
1:A:525:LYS:HA	2:A:28:GTG:C7X	2.30	0.60
1:C:521:ILE:HD13	1:C:547:VAL:CG1	2.32	0.60
1:B:524:GLY:O	2:B:28:GTG:O6A	2.20	0.59
1:B:443:HIS:CD2	1:B:572:GLY:HA3	2.38	0.59
1:B:522:LEU:HB3	2:B:28:GTG:HC72	1.80	0.57
1:B:475:ARG:HD3	1:B:487:GLY:HA3	1.87	0.56
1:A:549:MET:O	1:A:553:GLN:HG3	2.05	0.55
1:B:526:THR:HG22	2:B:28:GTG:O6A	2.06	0.55
1:A:448:ILE:HD12	1:A:485:ILE:HG12	1.87	0.55
1:A:528:VAL:O	1:A:532:VAL:HG23	2.07	0.54
1:B:450:VAL:CG1	1:B:515:LEU:HA	2.38	0.53
1:B:529:TYR:CD1	2:B:28:GTG:N2A	2.77	0.53
1:C:525:LYS:HD3	2:C:28:GTG:HC73	1.91	0.52
1:B:524:GLY:N	2:B:28:GTG:HC73	2.25	0.52
1:A:491:PHE:CZ	1:A:493:LYS:HD2	2.45	0.52
1:B:479:ARG:NE	1:B:480:ASP:OD1	2.37	0.52
1:C:457:ALA:HA	1:C:523:PRO:HB3	1.91	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:450:VAL:HB	1:B:514:GLY:O	2.11	0.50
1:B:524:GLY:C	2:B:28:GTG:C7X	2.81	0.49
1:B:545:GLN:HG2	1:B:567:ILE:CG1	2.42	0.49
1:C:441:GLN:HG3	1:C:442:PHE:N	2.28	0.49
1:B:526:THR:OG1	1:B:527:PRO:HD2	2.12	0.49
1:B:549:MET:O	1:B:553:GLN:HG3	2.13	0.48
1:B:533:LYS:O	1:B:537:ASP:HB2	2.12	0.48
1:C:525:LYS:HA	2:C:28:GTG:HC73	1.95	0.48
1:B:486:GLN:HG2	1:B:488:GLN:H	1.79	0.48
1:B:520:VAL:O	1:B:546:CYS:HA	2.13	0.48
1:A:554:ARG:HD3	1:B:480:ASP:HA	1.96	0.48
1:B:524:GLY:HA2	2:B:28:GTG:HC73	1.97	0.47
1:B:538:THR:O	1:B:538:THR:HG22	2.15	0.47
1:B:479:ARG:HD2	1:B:479:ARG:C	2.35	0.47
1:B:491:PHE:CE1	1:B:493:LYS:HD2	2.51	0.46
1:C:491:PHE:CD2	1:C:508:LEU:HD21	2.51	0.46
1:C:563:LEU:O	1:C:567:ILE:HG13	2.15	0.45
1:A:520:VAL:HG11	1:A:532:VAL:HG21	1.99	0.45
1:B:448:ILE:HD13	1:B:485:ILE:HG12	1.99	0.45
1:B:524:GLY:H	2:B:28:GTG:C7X	2.30	0.45
1:B:464:GLU:HG2	1:B:468:LYS:HE2	1.97	0.45
1:A:531:GLU:HA	1:A:531:GLU:OE1	2.17	0.44
1:C:483:MET:O	1:C:483:MET:HG3	2.16	0.44
1:A:502:GLU:HB3	1:A:503:PRO:HD3	1.99	0.44
1:B:491:PHE:CD2	1:B:508:LEU:HD21	2.52	0.44
1:B:548:GLN:HG3	2:B:28:GTG:O2D	2.17	0.43
1:A:486:GLN:O	1:A:486:GLN:HG3	2.19	0.43
1:A:533:LYS:HD3	1:A:533:LYS:HA	1.71	0.43
1:B:501:VAL:HG12	1:B:535:VAL:HG21	1.99	0.43
1:C:443:HIS:ND1	1:C:572:GLY:HA3	2.34	0.43
1:A:521:ILE:HG12	1:A:552:VAL:HG21	2.01	0.43
1:C:475:ARG:HH11	1:C:475:ARG:HG2	1.84	0.43
1:C:456:PHE:HA	1:C:495:ALA:O	2.19	0.42
1:C:446:ILE:HD12	1:C:568:ASN:ND2	2.35	0.42
2:B:28:GTG:O1B	2:B:28:GTG:O2A	2.37	0.41
1:A:524:GLY:O	2:A:28:GTG:HC73	2.19	0.41
1:B:520:VAL:HG11	1:B:532:VAL:HG21	2.02	0.41
1:C:520:VAL:HG11	1:C:532:VAL:HG21	2.01	0.41
1:A:525:LYS:HE2	1:A:525:LYS:HB3	1.73	0.41
1:A:525:LYS:HA	2:A:28:GTG:HC71	2.01	0.41
1:B:516:GLN:O	1:B:543:ALA:HB3	2.21	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:LYS:CD	1:A:441:GLN:N	2.71	0.41
1:C:475:ARG:HG2	1:C:475:ARG:NH1	2.36	0.41
1:C:531:GLU:OE2	1:C:531:GLU:HA	2.21	0.41
1:C:440:LYS:HE3	1:C:440:LYS:HB2	1.80	0.40
1:C:475:ARG:HD3	1:C:487:GLY:HA3	2.04	0.40
1:C:502:GLU:HB3	1:C:503:PRO:HD3	2.02	0.40
1:C:491:PHE:CZ	1:C:493:LYS:HD2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/138 (95%)	127 (97%)	4 (3%)	0	100	100
1	B	131/138 (95%)	125 (95%)	5 (4%)	1 (1%)	19	23
1	C	131/138 (95%)	126 (96%)	5 (4%)	0	100	100
All	All	393/414 (95%)	378 (96%)	14 (4%)	1 (0%)	41	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	487	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/119 (97%)	113 (98%)	2 (2%)	60	76
1	B	115/119 (97%)	111 (96%)	4 (4%)	36	50
1	C	115/119 (97%)	113 (98%)	2 (2%)	60	76
All	All	345/357 (97%)	337 (98%)	8 (2%)	50	67

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	500	SER
1	A	554	ARG
1	B	449	LYS
1	B	460	ARG
1	B	461	GLN
1	B	506	ARG
1	C	486	GLN
1	C	554	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	441	GLN
1	B	443	HIS
1	B	496	GLN
1	B	510	ASN
1	B	548	GLN
1	B	551	ASN
1	B	558	GLN
1	B	562	ASN
1	C	443	HIS
1	C	459	GLN
1	C	488	GLN
1	C	551	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GTG	B	28	-	46,57,57	1.42	6 (13%)	47,90,90	1.88	13 (27%)
2	GTG	C	28	-	46,57,57	1.77	11 (23%)	47,90,90	2.08	15 (31%)
2	GTG	A	28	-	46,57,57	1.99	6 (13%)	47,90,90	2.49	18 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTG	B	28	-	-	6/24/64/64	0/6/6/6
2	GTG	C	28	-	-	4/24/64/64	0/6/6/6
2	GTG	A	28	-	-	9/24/64/64	0/6/6/6

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	28	GTG	O4E-C1E	8.54	1.53	1.41
2	A	28	GTG	PG-O1G	5.80	1.71	1.50
2	C	28	GTG	O4D-C1D	5.39	1.48	1.41
2	B	28	GTG	O4E-C1E	4.47	1.47	1.41
2	C	28	GTG	O4E-C1E	4.41	1.47	1.41
2	A	28	GTG	O4E-C4E	3.74	1.53	1.45
2	A	28	GTG	PB-O1B	3.68	1.64	1.50
2	C	28	GTG	PG-O1G	3.28	1.62	1.50

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	28	GTG	PB-O1B	3.26	1.62	1.50
2	C	28	GTG	O4D-C4D	-3.16	1.37	1.45
2	C	28	GTG	C8A-N7A	-3.12	1.27	1.33
2	B	28	GTG	PG-O1G	3.00	1.61	1.50
2	C	28	GTG	PB-O1B	2.76	1.60	1.50
2	B	28	GTG	C8A-N7A	-2.66	1.28	1.33
2	A	28	GTG	O4D-C1D	2.64	1.44	1.41
2	C	28	GTG	C8B-N7B	-2.54	1.30	1.35
2	C	28	GTG	C2D-C1D	2.39	1.57	1.53
2	B	28	GTG	C8B-N7B	-2.30	1.31	1.35
2	A	28	GTG	C8A-N7A	-2.27	1.29	1.33
2	B	28	GTG	O4D-C1D	2.17	1.44	1.41
2	C	28	GTG	C5A-C4A	2.08	1.43	1.39
2	C	28	GTG	C3D-C4D	2.01	1.58	1.53
2	C	28	GTG	PA-O2A	2.00	1.64	1.55

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	28	GTG	PG-O3B-PB	-6.45	110.68	132.83
2	A	28	GTG	C2E-C3E-C4E	6.03	114.36	102.64
2	A	28	GTG	O4E-C1E-C2E	-5.88	98.33	106.93
2	C	28	GTG	O4D-C4D-C5D	5.77	128.35	109.37
2	A	28	GTG	PG-O3B-PB	-5.38	114.37	132.83
2	A	28	GTG	PB-O3A-PA	-4.82	116.28	132.83
2	B	28	GTG	O4D-C4D-C5D	4.64	124.65	109.37
2	A	28	GTG	O4D-C4D-C5D	4.63	124.61	109.37
2	B	28	GTG	PB-O3A-PA	-4.58	117.12	132.83
2	B	28	GTG	PG-O3B-PB	-4.52	117.32	132.83
2	A	28	GTG	O4E-C4E-C3E	-4.14	96.92	105.11
2	A	28	GTG	O3E-C3E-C4E	-4.06	99.32	111.05
2	A	28	GTG	C3E-C2E-C1E	4.03	107.05	100.98
2	C	28	GTG	C3D-C2D-C1D	-3.58	95.58	100.98
2	C	28	GTG	C8B-N7B-C5B	3.49	109.65	102.99
2	A	28	GTG	C8B-N7B-C5B	3.47	109.59	102.99
2	B	28	GTG	C8B-N7B-C5B	3.46	109.58	102.99
2	B	28	GTG	O4E-C1E-C2E	-3.36	102.01	106.93
2	A	28	GTG	O5E-PG-O1G	-3.28	96.25	109.07
2	B	28	GTG	PA-O5D-C5D	-3.01	104.00	121.68
2	C	28	GTG	PB-O3A-PA	-3.00	122.53	132.83
2	C	28	GTG	C5B-C6B-N1B	2.82	118.92	113.95
2	B	28	GTG	O4D-C1D-C2D	-2.77	102.88	106.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	28	GTG	C5B-C6B-N1B	2.74	118.79	113.95
2	C	28	GTG	O4E-C1E-C2E	-2.74	102.93	106.93
2	B	28	GTG	C5B-C6B-N1B	2.67	118.66	113.95
2	A	28	GTG	C2B-N1B-C6B	-2.63	120.25	125.10
2	B	28	GTG	C3D-C2D-C1D	-2.53	97.17	100.98
2	C	28	GTG	PG-O5E-C5E	-2.53	106.86	121.68
2	C	28	GTG	C2B-N1B-C6B	-2.51	120.48	125.10
2	A	28	GTG	O4D-C1D-C2D	-2.50	103.27	106.93
2	C	28	GTG	O4D-C1D-C2D	-2.47	103.31	106.93
2	B	28	GTG	C2B-N1B-C6B	-2.42	120.64	125.10
2	B	28	GTG	C2A-N1A-C6A	-2.42	120.64	125.10
2	C	28	GTG	C2D-C3D-C4D	-2.42	97.94	102.64
2	A	28	GTG	C2A-N1A-C6A	-2.34	120.79	125.10
2	A	28	GTG	O4E-C4E-C5E	-2.30	101.80	109.37
2	C	28	GTG	PA-O5D-C5D	-2.29	108.24	121.68
2	B	28	GTG	PG-O5E-C5E	-2.26	108.44	121.68
2	C	28	GTG	O5D-PA-O1A	-2.18	100.54	109.07
2	B	28	GTG	O5D-PA-O1A	-2.17	100.58	109.07
2	A	28	GTG	PA-O5D-C5D	-2.16	109.00	121.68
2	A	28	GTG	C3D-C2D-C1D	-2.14	97.75	100.98
2	C	28	GTG	C2A-N1A-C6A	-2.14	121.15	125.10
2	A	28	GTG	O2G-PG-O5E	2.01	117.09	107.75
2	C	28	GTG	O4E-C4E-C5E	2.01	115.97	109.37

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	28	GTG	C5D-O5D-PA-O1A
2	A	28	GTG	C5E-O5E-PG-O1G
2	B	28	GTG	C3D-C4D-C5D-O5D
2	A	28	GTG	O4E-C4E-C5E-O5E
2	A	28	GTG	C3E-C4E-C5E-O5E
2	B	28	GTG	O4D-C4D-C5D-O5D
2	A	28	GTG	PB-O3A-PA-O5D
2	A	28	GTG	PG-O3B-PB-O2B
2	B	28	GTG	PB-O3B-PG-O1G
2	C	28	GTG	C5D-O5D-PA-O1A
2	C	28	GTG	C5D-O5D-PA-O2A
2	B	28	GTG	PB-O3A-PA-O2A
2	B	28	GTG	PA-O3A-PB-O1B
2	A	28	GTG	C5E-O5E-PG-O3B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	28	GTG	C5D-O5D-PA-O3A
2	A	28	GTG	O4D-C4D-C5D-O5D
2	A	28	GTG	PG-O3B-PB-O1B
2	B	28	GTG	PB-O3A-PA-O1A
2	C	28	GTG	PB-O3B-PG-O2G


There are no ring outliers.

3 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	28	GTG	16	0
2	C	28	GTG	2	0
2	A	28	GTG	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/138 (96%)	-0.48	1 (0%) 86 89	19, 28, 42, 69	0
1	B	133/138 (96%)	-0.04	3 (2%) 60 67	24, 36, 50, 63	0
1	C	133/138 (96%)	-0.27	3 (2%) 60 67	22, 32, 46, 68	0
All	All	399/414 (96%)	-0.26	7 (1%) 68 74	19, 32, 50, 69	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	525	LYS	3.3
1	B	538	THR	3.1
1	C	486	GLN	2.6
1	C	487	GLY	2.5
1	C	440	LYS	2.2
1	B	459	GLN	2.2
1	B	499	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GTG	B	28	52/52	0.74	0.32	59,77,89,89	23
2	GTG	A	28	52/52	0.78	0.24	51,62,73,75	23
2	GTG	C	28	52/52	0.82	0.19	54,70,76,78	23

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around GTG A 28:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GTG C 28:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.