



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2021 – 10:44 AM EST

PDB ID : 5R0F  
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment F2X-Entry D06, DMSO-free  
Authors : Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.; Weiss, M.S.  
Deposited on : 2020-02-12  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.17.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.17.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	
2	B	308	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	1994	1278	334	371	11	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2568	1645	420	483	20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	170	SER	LEU	conflict	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ASP	deletion	UNP P32357

- Molecule 3 is (3-methoxyphenyl)(pyrrolidin-1-yl)methanone (three-letter code: R9G) (formula: C<sub>12</sub>H<sub>15</sub>NO<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	15	12	1	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	45	Total	O	0	0
			45	45		
4	B	21	Total	O	0	0
			21	21		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.59Å 82.24Å 91.64Å 90.00° 108.25° 90.00°	Depositor
Resolution (Å)	22.34 – 1.97 44.61 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.5 (22.34-1.97) 99.5 (44.61-1.97)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.95 (at 1.97Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.261 , 0.304 0.273 , 0.314	Depositor DCC
$R_{free}$ test set	2100 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.1	Xtrriage
Anisotropy	0.161	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 61.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4643	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R9G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.86	4/2041 (0.2%)	1.02	14/2765 (0.5%)
2	B	0.74	1/2638 (0.0%)	0.83	2/3563 (0.1%)
All	All	0.79	5/4679 (0.1%)	0.92	16/6328 (0.3%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1887	GLY	C-O	-7.76	1.11	1.23
1	A	1899	TRP	CE3-CZ3	5.87	1.48	1.38
1	A	1992	TYR	CD1-CE1	-5.19	1.31	1.39
1	A	1992	TYR	CD2-CE2	-5.11	1.31	1.39
2	B	74	LYS	CD-CE	5.09	1.64	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1940	MET	CG-SD-CE	-14.77	76.56	100.20
1	A	1996	LEU	CB-CG-CD1	-7.43	98.36	111.00
2	B	112	ASP	CB-CG-OD1	6.23	123.91	118.30
1	A	1993	ASP	CB-CA-C	5.81	122.02	110.40
1	A	2007	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	A	1986	MET	C-N-CA	5.43	135.28	121.70
1	A	1973	LYS	CB-CA-C	5.26	120.91	110.40
1	A	1997	ASP	CB-CG-OD1	-5.23	113.59	118.30
1	A	1987	VAL	N-CA-CB	-5.20	100.06	111.50
1	A	2026	LEU	CB-CG-CD2	-5.19	102.17	111.00
1	A	2011	LEU	CB-CG-CD2	-5.19	102.18	111.00
2	B	74	LYS	CD-CE-NZ	5.13	123.49	111.70
1	A	1961	LEU	CB-CG-CD2	-5.12	102.30	111.00
1	A	1986	MET	CB-CG-SD	-5.12	97.04	112.40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2035	LYS	O-C-N	-5.11	114.53	122.70
1	A	1940	MET	CB-CG-SD	5.07	127.61	112.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1994	0	2020	28	0
2	B	2568	0	2441	33	0
3	A	15	0	0	0	0
4	A	45	0	0	1	0
4	B	21	0	0	1	0
All	All	4643	0	4461	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1842:GLU:OE1	4:A:2601:HOH:O	1.92	0.87
2:B:1:MET:N	4:B:401:HOH:O	2.09	0.86
1:A:1918[B]:SER:OG	1:A:1922:ARG:NH2	2.11	0.82
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.80	0.81
1:A:1951:PHE:HB3	1:A:1954:ILE:HD12	1.69	0.74
1:A:2058:LEU:C	1:A:2058:LEU:HD23	2.19	0.63
2:B:47:GLN:NE2	2:B:52:SER:O	2.32	0.62
1:A:1962:ARG:O	1:A:1962:ARG:HG2	1.99	0.61
2:B:287:ARG:O	2:B:291:ILE:HD13	1.99	0.61
2:B:277:GLU:CD	2:B:277:GLU:H	2.05	0.59
1:A:1930:PRO:O	1:A:1954:ILE:HG12	2.03	0.59
2:B:18:ASP:OD2	2:B:105:TYR:OH	2.17	0.58
1:A:2044:THR:OG1	1:A:2047:GLN:HG3	2.07	0.55

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2028:SER:O	1:A:2030:PRO:HD3	2.07	0.54
2:B:96:PHE:HB2	2:B:102:MET:HE3	1.92	0.52
1:A:1910:LYS:HA	1:A:1940:MET:SD	2.50	0.51
2:B:41[B]:VAL:HG12	2:B:123:MET:HG2	1.93	0.51
2:B:16:GLY:HA3	2:B:45:HIS:CE1	2.46	0.50
2:B:203[A]:TYR:CZ	2:B:207:THR:HG21	2.45	0.50
2:B:61:ASP:HB2	2:B:123:MET:HG3	1.94	0.50
1:A:1902:GLN:HB2	1:A:1905:LEU:CD2	2.43	0.49
2:B:53:SER:O	2:B:54[A]:MET:HB3	2.12	0.49
2:B:59:TRP:HH2	2:B:229:LEU:CD1	2.26	0.49
1:A:1848:ILE:H	1:A:1931[A]:LYS:NZ	2.11	0.48
2:B:251:CYS:SG	2:B:292:CYS:HB3	2.53	0.48
1:A:1891:LEU:HD22	1:A:1974:LEU:HD22	1.95	0.48
2:B:144:MET:HE2	2:B:176:LEU:HG	1.94	0.48
1:A:2052:GLU:O	1:A:2056[B]:ARG:HG3	2.14	0.48
1:A:1951:PHE:CB	1:A:1954:ILE:HD12	2.42	0.47
2:B:140:VAL:HG12	2:B:141:ASP:N	2.29	0.47
1:A:1913:THR:HB	1:A:1940:MET:HE1	1.97	0.47
2:B:141:ASP:OD1	2:B:141:ASP:C	2.53	0.46
2:B:59:TRP:HH2	2:B:229:LEU:HD12	1.80	0.46
2:B:251:CYS:O	2:B:296:SER:HB2	2.15	0.46
2:B:250:ILE:HG21	2:B:264:LEU:HD22	1.98	0.45
2:B:195:MET:HG2	2:B:199:LEU:HD12	1.99	0.45
1:A:2021:SER:HB3	1:A:2058:LEU:HD12	1.99	0.44
2:B:6:PHE:CE2	2:B:69:ILE:HG21	2.53	0.44
1:A:2064:GLY:O	1:A:2068:ASN:N	2.45	0.44
1:A:1941:LEU:O	1:A:1945:GLU:HB2	2.18	0.44
2:B:18:ASP:OD2	2:B:42:HIS:ND1	2.42	0.44
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.52	0.44
1:A:1910:LYS:HG2	1:A:1940:MET:SD	2.57	0.44
1:A:1859:ARG:HB2	1:A:1875:ILE:HG13	2.00	0.44
1:A:2021:SER:HB3	1:A:2058:LEU:CD1	2.48	0.44
1:A:1851:PHE:O	1:A:1881:THR:HA	2.19	0.43
2:B:231:ALA:O	2:B:235:GLY:N	2.48	0.43
2:B:267:ILE:O	2:B:271:GLN:HG3	2.18	0.43
2:B:214:PHE:O	2:B:215:LYS:HB2	2.19	0.42
1:A:1975:SER:O	1:A:1979[A]:MET:SD	2.78	0.42
2:B:10:PRO:O	2:B:26:GLU:HG3	2.19	0.41
2:B:214:PHE:O	2:B:215:LYS:CB	2.68	0.41
2:B:285:ASN:OD1	2:B:288:VAL:HG23	2.20	0.41
2:B:15:ILE:O	2:B:21:SER:HA	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:70:GLN:OE1	2:B:81:MET:HE1	2.20	0.41
1:A:1835:MET:HE3	1:A:1961:LEU:HD12	2.01	0.41
1:A:1902:GLN:HB2	1:A:1905:LEU:HD21	2.02	0.41
1:A:1835:MET:HE3	1:A:1835:MET:HB3	1.96	0.41
2:B:292:CYS:HA	2:B:296:SER:HB3	2.02	0.40
1:A:2056[B]:ARG:O	1:A:2060:LEU:HG	2.20	0.40
2:B:225:GLN:O	2:B:229:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/258 (96%)	239 (97%)	8 (3%)	0	100	100
2	B	305/308 (99%)	285 (93%)	19 (6%)	1 (0%)	41	29
All	All	552/566 (98%)	524 (95%)	27 (5%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	215	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/233 (97%)	217 (96%)	8 (4%)	35	23
2	B	286/284 (101%)	279 (98%)	7 (2%)	49	41
All	All	511/517 (99%)	496 (97%)	15 (3%)	40	31

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1847	ASP
1	A	1903	LYS
1	A	1962	ARG
1	A	1977	VAL
1	A	1986	MET
1	A	2021	SER
1	A	2065	ARG
2	B	40	HIS
2	B	53	SER
2	B	77	LEU
2	B	124	ASP
2	B	280	SER
2	B	315	GLU
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	R9G	A	2501	-	16,16,16	2.04	6 (37%)	21,21,21	0.96	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	R9G	A	2501	-	-	0/10/17/17	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2501	R9G	C11-N	-3.76	1.39	1.47
3	A	2501	R9G	C10-C9	-3.52	1.23	1.48
3	A	2501	R9G	C8-N	-3.47	1.39	1.47
3	A	2501	R9G	O1-C7	-3.02	1.16	1.22
3	A	2501	R9G	O-C	-2.37	1.35	1.42
3	A	2501	R9G	C2-C1	-2.10	1.34	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2501	R9G	O-C1-C2	-2.42	108.26	119.82
3	A	2501	R9G	O-C1-C6	2.16	130.26	119.94


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	237/258 (91%)	1.50	61 (25%) 0 0	39, 74, 116, 151	0
2	B	300/308 (97%)	1.79	93 (31%) 0 0	48, 84, 143, 178	0
All	All	537/566 (94%)	1.66	154 (28%) 0 0	39, 81, 131, 178	0

All (154) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	173	ALA	18.1
2	B	52	SER	7.4
2	B	1	MET	6.8
1	A	1878	CYS	6.8
2	B	174	HIS	6.7
2	B	150	ASN	6.6
2	B	172	PRO	6.5
2	B	53	SER	6.2
2	B	109	ASP	6.1
2	B	149	GLU	5.8
2	B	175	SER	5.7
2	B	170	SER	5.6
2	B	281	ASP	5.5
1	A	1890	PHE	5.1
2	B	54[A]	MET	5.0
2	B	108	ILE	5.0
2	B	313	TYR	4.9
2	B	92	ILE	4.8
2	B	279	TYR	4.8
2	B	22	PHE	4.8
2	B	147	VAL	4.7
1	A	1969	MET	4.6
2	B	146	THR	4.5
2	B	291	ILE	4.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	B	316	LEU	4.5
1	A	2046	GLU	4.5
2	B	89	PHE	4.1
2	B	193	HIS	4.1
2	B	11	ILE	4.0
1	A	2015	LEU	4.0
1	A	2064	GLY	3.9
1	A	2057[A]	ASP	3.9
1	A	1940	MET	3.9
2	B	132	LYS	3.9
2	B	171	ASP	3.8
1	A	1834	ALA	3.8
2	B	213	ILE	3.7
1	A	2047	GLN	3.7
1	A	2044	THR	3.7
1	A	2063	TYR	3.6
1	A	1989	PHE	3.6
1	A	2032	ILE	3.6
1	A	1840	TYR	3.6
1	A	2065	ARG	3.6
2	B	148	GLN	3.5
2	B	280	SER	3.5
2	B	133	ASP	3.5
1	A	2069	VAL	3.5
2	B	87	ALA	3.5
1	A	1891	LEU	3.5
2	B	77	LEU	3.5
1	A	2017[A]	THR	3.4
2	B	96	PHE	3.4
1	A	1833	GLY	3.4
2	B	86	GLY	3.4
1	A	2058	LEU	3.3
1	A	1988	LEU	3.3
1	A	2037	TYR	3.3
2	B	250	ILE	3.3
1	A	1971	ILE	3.2
2	B	103	VAL	3.2
1	A	2068	ASN	3.2
2	B	74	LYS	3.2
2	B	275	LEU	3.2
1	A	2024[A]	MET	3.1
1	A	1881	THR	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	B	41[A]	VAL	3.1
2	B	203[A]	TYR	3.1
2	B	191	PRO	3.0
1	A	1865	THR	3.0
2	B	196	GLU	3.0
1	A	1934	ILE	3.0
2	B	91	ASN	3.0
2	B	207	THR	2.9
1	A	2051	ILE	2.9
1	A	1879	ILE	2.9
2	B	237	TYR	2.8
2	B	84	ARG	2.8
1	A	2049	ILE	2.8
2	B	181	ILE	2.8
2	B	235	GLY	2.8
2	B	107	LYS	2.7
1	A	1905	LEU	2.7
2	B	94	HIS	2.7
2	B	101	MET	2.7
1	A	1991	ILE	2.6
2	B	188	ALA	2.6
2	B	198	PHE	2.6
1	A	2061	THR	2.6
2	B	278	GLN	2.6
2	B	221	PHE	2.5
2	B	73	PRO	2.5
1	A	1837	SER	2.5
1	A	1889	LEU	2.5
1	A	1848	ILE	2.5
1	A	1999	ILE	2.5
2	B	253	SER	2.5
1	A	1895	HIS	2.5
2	B	243	TRP	2.5
2	B	100	GLN	2.5
1	A	1921	VAL	2.5
1	A	2018	ASN	2.4
2	B	88	LYS	2.4
1	A	1843	LEU	2.4
2	B	129	ILE	2.4
1	A	1987	VAL	2.4
2	B	138	SER	2.3
1	A	1882	LEU	2.3

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	2027	LEU	2.3
1	A	1909	ALA	2.3
2	B	205	LEU	2.3
2	B	249	LEU	2.3
2	B	297	PHE	2.3
2	B	190	ARG	2.3
2	B	276	PRO	2.3
1	A	2048	TRP	2.3
2	B	126	ILE	2.3
2	B	232	MET	2.3
1	A	1951	PHE	2.3
2	B	309	MET	2.2
1	A	2031	THR	2.2
2	B	230[A]	ASN	2.2
2	B	295	SER	2.2
1	A	1949	LEU	2.2
2	B	152	LEU	2.2
2	B	176	LEU	2.2
2	B	294	TYR	2.2
1	A	1880	PHE	2.2
2	B	122[A]	GLN	2.2
2	B	123	MET	2.2
1	A	2033	THR	2.2
2	B	212	GLY	2.2
2	B	228	PHE	2.2
1	A	1930	PRO	2.1
2	B	17	ILE	2.1
1	A	1927	GLU	2.1
1	A	1979[A]	MET	2.1
1	A	1888	HIS	2.1
2	B	208	VAL	2.1
1	A	1850	LEU	2.1
2	B	9	ALA	2.1
2	B	227	ALA	2.1
2	B	119	GLU	2.1
2	B	121	VAL	2.1
2	B	128	LYS	2.1
1	A	2006	SER	2.1
2	B	293	LEU	2.1
2	B	43	VAL	2.1
1	A	2059	ILE	2.1
2	B	272	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	2035	LYS	2.1
2	B	110	GLU	2.0
2	B	111	ASP	2.0
2	B	26	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	R9G	A	2501	15/15	0.93	0.17	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.