



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 08:55 pm BST

PDB ID : 2R14  
Title : Structure of morphinone reductase in complex with tetrahydroNAD  
Authors : Costello, C.L.; Scrutton, N.S.; Leys, D.  
Deposited on : 2007-08-22  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Morphinone reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	363	2807	1767	500	533	7	0	0	0

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).







## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.27Å 118.89Å 180.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.00 – 1.40 26.70 – 1.30	Depositor EDS
% Data completeness (in resolution range)	89.2 (12.00-1.40) 93.1 (26.70-1.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.37 (at 1.30Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.156 , 0.181 0.157 , 0.182	Depositor DCC
$R_{free}$ test set	6208 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.3	Xtrriage
Anisotropy	0.847	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 55.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3569	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, TXD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.82	1/2877 (0.0%)	0.95	7/3914 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	35	ARG	CG-CD	-6.29	1.36	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	169	ARG	NE-CZ-NH1	9.78	125.19	120.30
1	A	375	ARG	NE-CZ-NH2	-7.20	116.70	120.30
1	A	375	ARG	NE-CZ-NH1	6.87	123.73	120.30
1	A	81	ASP	CB-CG-OD1	6.38	124.04	118.30
1	A	35	ARG	CG-CD-NE	6.01	124.43	111.80
1	A	169	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	A	342	ARG	NE-CZ-NH2	5.43	123.02	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2807	0	2691	38	0
2	A	31	0	19	1	0
3	A	47	0	8	0	0
4	A	684	0	0	10	0
All	All	3569	0	2718	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:GLN:HE21	1:A:207:GLN:H	1.16	0.91
1:A:334:ASN:HD21	1:A:348:ASN:H	1.17	0.91
1:A:113:HIS:HD2	1:A:115:LEU:H	1.27	0.83
1:A:190:ALA:H	1:A:194:ASN:HD22	1.32	0.76
1:A:24:ASN:HD22	1:A:26:VAL:H	1.36	0.72
1:A:266:ARG:HD2	4:A:961:HOH:O	1.89	0.72
1:A:334:ASN:HD21	1:A:348:ASN:N	1.88	0.70
1:A:184:GLU:HB2	4:A:1272:HOH:O	1.92	0.70
1:A:51:GLN:HE21	1:A:52:ARG:HH12	1.40	0.69
1:A:202:ASN:C	1:A:202:ASN:HD22	2.02	0.62
1:A:8:ASN:N	4:A:1086:HOH:O	2.31	0.61
1:A:334:ASN:ND2	1:A:348:ASN:H	1.93	0.60
1:A:113:HIS:CD2	1:A:115:LEU:H	2.17	0.57
1:A:188:ALA:HB1	1:A:240:THR:HG23	1.85	0.57
1:A:285:THR:N	4:A:977:HOH:O	2.38	0.57
1:A:214:ASN:ND2	1:A:217:ARG:HH21	2.03	0.56
1:A:278:ASP:HB2	4:A:739:HOH:O	2.05	0.55
1:A:59:ILE:HG23	4:A:1272:HOH:O	2.07	0.54
1:A:132:GLU:OE2	4:A:891:HOH:O	2.19	0.51
1:A:214:ASN:HD22	1:A:217:ARG:HH21	1.59	0.51
1:A:285:THR:HG22	4:A:1110:HOH:O	2.12	0.50
1:A:278:ASP:OD1	4:A:907:HOH:O	2.20	0.48
1:A:24:ASN:ND2	1:A:26:VAL:H	2.09	0.47
1:A:190:ALA:H	1:A:194:ASN:ND2	2.06	0.47
1:A:285:THR:OG1	1:A:286:TYR:N	2.48	0.46
1:A:51:GLN:NE2	1:A:52:ARG:HH12	2.12	0.46
1:A:240:THR:HG22	1:A:275:ASN:HB3	1.98	0.45
1:A:192:LEU:HB3	1:A:193:PRO:HD3	2.00	0.43
1:A:202:ASN:ND2	1:A:204:ARG:H	2.16	0.43
1:A:294:MET:HE3	1:A:294:MET:HA	2.01	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:TYR:CE1	2:A:401:FMN:HM72	2.54	0.42
1:A:202:ASN:HD22	1:A:204:ARG:H	1.66	0.42
1:A:69:ALA:HB1	1:A:107:HIS:HA	2.02	0.41
1:A:240:THR:CG2	1:A:275:ASN:HB3	2.50	0.41
1:A:296:GLN:HG2	4:A:1073:HOH:O	2.20	0.41
1:A:207:GLN:HE21	1:A:207:GLN:N	1.99	0.41
1:A:24:ASN:HD21	1:A:56:GLY:H	1.69	0.41
1:A:202:ASN:C	1:A:202:ASN:ND2	2.71	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	359/377 (95%)	348 (97%)	11 (3%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	284/296 (96%)	278 (98%)	6 (2%)	53 21

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	186	HIS
1	A	202	ASN
1	A	203	ARG
1	A	207	GLN
1	A	331	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	51	GLN
1	A	113	HIS
1	A	194	ASN
1	A	202	ASN
1	A	207	GLN
1	A	214	ASN
1	A	307	ASN
1	A	334	ASN
1	A	348	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TXD	A	600[A]	-	42,48,48	3.11	8 (19%)	49,73,73	2.11	8 (16%)
2	FMN	A	401	-	31,33,33	1.51	5 (16%)	40,50,50	2.36	9 (22%)
3	TXD	A	600[B]	-	42,48,48	3.12	8 (19%)	49,73,73	2.09	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TXD	A	600[A]	-	-	3/26/72/72	0/5/5/5
2	FMN	A	401	-	-	1/18/18/18	0/3/3/3
3	TXD	A	600[B]	-	-	3/26/72/72	0/5/5/5

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	600[A]	TXD	C2N-C3N	-13.11	1.39	1.52
3	A	600[B]	TXD	C2N-C3N	-13.11	1.39	1.52
3	A	600[A]	TXD	O7N-C7N	7.71	1.38	1.23
3	A	600[B]	TXD	O7N-C7N	7.71	1.38	1.23
3	A	600[B]	TXD	C4N-C5N	7.70	1.51	1.32
3	A	600[A]	TXD	C4N-C5N	7.34	1.51	1.32
3	A	600[A]	TXD	C2N-N1N	-6.75	1.36	1.47
3	A	600[B]	TXD	C2N-N1N	-6.75	1.36	1.47
3	A	600[A]	TXD	C3N-C4N	-5.17	1.40	1.52
3	A	600[B]	TXD	C3N-C4N	-4.88	1.40	1.52
2	A	401	FMN	C10-N1	3.91	1.38	1.33
3	A	600[B]	TXD	C6N-N1N	-3.86	1.38	1.47
2	A	401	FMN	C4-N3	3.81	1.39	1.33
3	A	600[A]	TXD	C6N-N1N	-3.81	1.38	1.47
2	A	401	FMN	C5'-C4'	3.12	1.56	1.51
3	A	600[A]	TXD	C3N-C7N	-2.61	1.50	1.53
3	A	600[B]	TXD	C3N-C7N	-2.61	1.50	1.53
2	A	401	FMN	C4A-N5	2.45	1.36	1.33
3	A	600[A]	TXD	C2A-N3A	2.15	1.35	1.32

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	600[B]	TXD	C2A-N3A	2.15	1.35	1.32
2	A	401	FMN	O4-C4	2.03	1.29	1.24

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FMN	C4-N3-C2	9.53	123.19	115.14
3	A	600[A]	TXD	C3N-C4N-C5N	-8.66	108.53	123.09
3	A	600[B]	TXD	C2N-C3N-C4N	7.59	119.46	108.63
3	A	600[B]	TXD	C3N-C4N-C5N	-7.42	110.62	123.09
3	A	600[A]	TXD	C2N-C3N-C4N	6.94	118.54	108.63
2	A	401	FMN	C4A-C4-N3	-5.77	115.54	123.43
3	A	600[B]	TXD	C6N-C5N-C4N	-5.66	111.46	123.05
3	A	600[A]	TXD	C6N-C5N-C4N	-5.57	111.65	123.05
2	A	401	FMN	C1'-N10-C9A	4.63	121.94	118.29
2	A	401	FMN	C4A-N5-C5A	3.99	120.76	116.77
3	A	600[A]	TXD	N3A-C2A-N1A	-3.68	122.93	128.68
3	A	600[B]	TXD	N3A-C2A-N1A	-3.68	122.93	128.68
3	A	600[A]	TXD	C4N-C3N-C7N	2.90	118.11	110.08
2	A	401	FMN	C1'-N10-C10	-2.75	115.94	118.41
3	A	600[B]	TXD	C4N-C3N-C7N	2.74	117.66	110.08
2	A	401	FMN	C8M-C8-C7	2.54	125.94	120.74
3	A	600[B]	TXD	C6N-N1N-C1'A	2.54	120.25	112.83
2	A	401	FMN	C6-C5A-N5	2.52	121.83	119.05
2	A	401	FMN	C4'-C3'-C2'	2.23	118.00	113.36
3	A	600[A]	TXD	O7N-C7N-N7N	-2.21	119.16	123.00
3	A	600[B]	TXD	O7N-C7N-N7N	-2.21	119.16	123.00
3	A	600[A]	TXD	C3N-C7N-N7N	2.17	119.19	116.85
3	A	600[B]	TXD	C3N-C7N-N7N	2.17	119.19	116.85
3	A	600[A]	TXD	O4'A-C1'A-N1N	-2.12	106.41	109.47
3	A	600[B]	TXD	O4'A-C1'A-N1N	-2.12	106.41	109.47
2	A	401	FMN	C8M-C8-C9	-2.11	115.30	120.34

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	600[A]	TXD	C4N-C3N-C7N-N7N
3	A	600[A]	TXD	O4'A-C1'A-N1N-C2N
3	A	600[B]	TXD	C4N-C3N-C7N-N7N
3	A	600[B]	TXD	O4'A-C1'A-N1N-C2N
2	A	401	FMN	C4'-C5'-O5'-P

Continued on next page...

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	A	600[A]	TXD	C2'A-C1'A-N1N-C2N
3	A	600[B]	TXD	C2'A-C1'A-N1N-C2N


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/377 (96%)	-0.32	2 (0%) 89 88	13, 18, 26, 35	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	278	ASP	2.9
1	A	8	ASN	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	TXD	A	600[A]	44/44	0.97	0.07	17,21,24,24	3
2	FMN	A	401	31/31	0.97	0.08	13,15,16,17	0
3	TXD	A	600[B]	44/44	0.97	0.07	17,21,22,24	3




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers


as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around FMN A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.