



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 04:48 PM BST

PDB ID : 4R8W
Title : Crystal structure of H7 hemagglutinin from A/Anhui/1/2013 in complex with a neutralizing antibody CT149
Authors : Wu, Y.; Shi, Y.; Qi, J.; Gao, G.F.
Deposited on : 2014-09-03
Resolution : 2.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	5% (Poor fit) 83% (0 outliers) 14% (1 outlier) •• (2 outliers)
2	B	177	7% (Poor fit) 83% (0 outliers) 10% (1 outlier) • 5% (2 outliers)
3	H	231	10% (Poor fit) 42% (0 outliers) 12% (1 outlier) • (2 outliers) 45% (Not modelled)
4	L	216	3% (Poor fit) 42% (0 outliers) 7% (1 outlier) •• (2 outliers) 49% (Not modelled)

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5675 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2412	1498	436	463	15	0	0	0

- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	169	1375	849	240	279	7	0	0	0


- Molecule 3 is a protein called Heavy chain of neutralizing antibody CT149.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	H	127	964	606	169	187	2	0	0	0

- Molecule 4 is a protein called Light chain of neutralizing antibody CT149.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	L	110	857	534	157	164	2	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	10	Total	O	0	0
			10	10		
6	B	13	Total	O	0	0
			13	13		
6	L	2	Total	O	0	0
			2	2		

SER GLY THR ALA VAL VAL CYS LEU LEU ASN ASN PHE TYR PRO ARG GLU ALA LYS VAL GLN TRP LYS VAL ASP ASN ALA LEU GLN SER GLY ASN SER GLN GLU SER VAL THR GLU GLN ASP SER LYS ASP SER THR TYR LEU SER SER THR THR LEU SER LYS ALA ASP TYR

GLU LYS HIS VAL ALA CYS VAL THR HIS GLN GLY LEU SER PRO VAL THR LYS SER PHE ASN ARG GLY GLU CYS SER

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	126.86Å 126.86Å 409.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.41 – 2.79 48.41 – 2.79	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.41-2.79) 99.4 (48.41-2.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.262 , 0.311 0.276 , 0.311	Depositor DCC
R_{free} test set	1606 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	66.4	Xtrriage
Anisotropy	0.234	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	5675	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.24	0/2458	0.42	0/3322
2	B	0.25	0/1399	0.42	0/1885
3	H	0.28	0/986	0.46	0/1340
4	L	0.24	0/877	0.53	2/1191 (0.2%)
All	All	0.25	0/5720	0.44	2/7738 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	52	ALA	CB-CA-C	-5.66	101.61	110.10
4	L	108	LYS	CB-CA-C	-5.17	100.05	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2412	0	2372	21	0
2	B	1375	0	1274	12	0
3	H	964	0	934	24	0
4	L	857	0	837	16	0
5	A	28	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	14	0	13	0	0
6	A	10	0	0	0	0
6	B	13	0	0	1	0
6	L	2	0	0	0	0
All	All	5675	0	5456	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:27:TYR:CD1	3:H:28:SER:O	2.13	1.02
3:H:73:ASP:HB3	3:H:78:THR:H	1.32	0.95
4:L:109:ARG:HE	4:L:109:ARG:HA	1.32	0.94
4:L:109:ARG:NH1	4:L:110:THR:HG22	1.89	0.86
3:H:91:THR:HG22	3:H:124:VAL:H	1.42	0.84
3:H:29:PHE:O	3:H:30:SER:OG	1.96	0.83
4:L:109:ARG:HE	4:L:109:ARG:CA	2.01	0.72
3:H:73:ASP:HB3	3:H:78:THR:N	2.06	0.67
3:H:27:TYR:HD1	3:H:28:SER:O	1.76	0.66
2:B:326:ALA:HB1	2:B:329:GLY:HA3	1.81	0.63
2:B:484:ARG:NH1	6:B:605:HOH:O	2.31	0.62
1:A:37:ASN:HB2	1:A:288:ILE:HD13	1.80	0.62
2:B:376:LEU:HD22	2:B:420:LEU:HD21	1.82	0.62
1:A:121:ARG:NH1	1:A:146:ASN:O	2.33	0.61
2:B:376:LEU:HD11	2:B:424:GLU:HG3	1.81	0.61
1:A:197:SER:OG	1:A:198:SER:N	2.35	0.60
3:H:27:TYR:CE1	3:H:28:SER:O	2.55	0.59
1:A:121:ARG:NH1	1:A:145:SER:O	2.36	0.59
4:L:86:VAL:HG22	4:L:104:ARG:HD2	1.87	0.56
2:B:375:ARG:NH2	2:B:424:GLU:OE2	2.38	0.56
1:A:207:SER:O	1:A:211:ARG:NH2	2.36	0.55
3:H:109:SER:O	4:L:47:ARG:NH2	2.41	0.54
1:A:163:ARG:HD3	1:A:250:PHE:CZ	2.44	0.53
1:A:106:GLY:HA2	1:A:255:SER:HB3	1.90	0.53
4:L:109:ARG:NE	4:L:109:ARG:CA	2.72	0.53
3:H:66:GLY:O	3:H:84:ARG:NH1	2.42	0.52
2:B:380:THR:HB	2:B:382:GLN:H	1.75	0.52
1:A:53:GLN:HG2	1:A:82:GLU:HG3	1.92	0.52
1:A:110:LYS:NZ	1:A:139:GLU:OE2	2.39	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:73:ASP:N	3:H:78:THR:O	2.45	0.50
4:L:109:ARG:NE	4:L:110:THR:N	2.60	0.50
4:L:51:GLY:O	4:L:52:ALA:HB3	2.13	0.49
4:L:109:ARG:NE	4:L:110:THR:H	2.10	0.48
3:H:32:TYR:CG	3:H:98:ARG:HD3	2.48	0.48
1:A:191:LYS:HA	1:A:239:ASN:HD21	1.78	0.48
1:A:60:ILE:HG21	1:A:170:VAL:HG21	1.96	0.48
2:B:327:ILE:HB	2:B:433:ASP:OD1	2.14	0.48
4:L:109:ARG:HH11	4:L:110:THR:HG22	1.77	0.48
4:L:24:ARG:NH1	4:L:71:ASP:OD2	2.47	0.47
3:H:73:ASP:CB	3:H:77:ALA:N	2.77	0.47
3:H:32:TYR:CD1	3:H:98:ARG:HD3	2.49	0.47
4:L:38:GLN:HB2	4:L:48:LEU:HD11	1.96	0.47
1:A:139:GLU:OE1	1:A:247:ARG:HD3	2.15	0.47
3:H:51:ILE:HD13	3:H:72:THR:HG23	1.97	0.46
3:H:73:ASP:HB2	3:H:77:ALA:N	2.30	0.46
2:B:471:ALA:O	2:B:475:ASN:ND2	2.48	0.46
4:L:55:ARG:NH1	4:L:63:PHE:O	2.49	0.45
3:H:50:TRP:CH2	3:H:52:SER:HB2	2.51	0.45
2:B:323:LEU:HB3	2:B:324:PHE:H	1.45	0.45
3:H:73:ASP:HB2	3:H:77:ALA:H	1.82	0.45
1:A:43:SER:HB2	1:A:48:THR:OG1	2.17	0.44
3:H:29:PHE:O	3:H:30:SER:CB	2.65	0.44
2:B:446:GLN:NE2	2:B:476:ASN:HA	2.32	0.44
3:H:64:PHE:O	3:H:68:VAL:HG12	2.18	0.43
1:A:110:LYS:HB2	1:A:247:ARG:NH2	2.33	0.43
3:H:27:TYR:HB3	3:H:98:ARG:HH21	1.84	0.43
3:H:73:ASP:HB3	3:H:77:ALA:CA	2.49	0.43
1:A:4:CYS:HA	2:B:458:CYS:HA	2.01	0.42
1:A:137:TYR:HB2	1:A:140:MET:HB2	2.01	0.42
1:A:5:LEU:HD11	2:B:440:TYR:HA	2.01	0.42
4:L:81:PRO:HA	4:L:107:ILE:HD12	2.02	0.42
3:H:73:ASP:CB	3:H:77:ALA:H	2.32	0.42
3:H:2:VAL:HA	3:H:26:GLY:HA3	2.01	0.42
1:A:186:TYR:CZ	1:A:241:ALA:HA	2.55	0.41
1:A:35:ARG:HD3	1:A:303:ARG:HG2	2.02	0.41
4:L:109:ARG:NH1	4:L:110:THR:CG2	2.73	0.41
4:L:109:ARG:CZ	4:L:110:THR:H	2.34	0.41
3:H:24:THR:HB	3:H:77:ALA:O	2.21	0.40
1:A:170:VAL:HG22	1:A:225:TRP:HB3	2.04	0.40
1:A:153:PRO:O	1:A:155:MET:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/321 (98%)	291 (93%)	22 (7%)	1 (0%)	41	72
2	B	167/177 (94%)	155 (93%)	11 (7%)	1 (1%)	25	56
3	H	125/231 (54%)	113 (90%)	10 (8%)	2 (2%)	9	31
4	L	108/216 (50%)	100 (93%)	6 (6%)	2 (2%)	8	26
All	All	714/945 (76%)	659 (92%)	49 (7%)	6 (1%)	19	49

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	H	73	ASP
3	H	75	THR
2	B	464	LYS
4	L	109	ARG
1	A	132	SER
4	L	107	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/269 (98%)	249 (94%)	16 (6%)	19	48
2	B	145/152 (95%)	139 (96%)	6 (4%)	30	64

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	H	101/192 (53%)	93 (92%)	8 (8%)	12	34
4	L	92/187 (49%)	85 (92%)	7 (8%)	13	36
All	All	603/800 (75%)	566 (94%)	37 (6%)	18	48

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	8	HIS
1	A	19	LEU
1	A	20	THR
1	A	65	GLN
1	A	116	THR
1	A	121	ARG
1	A	126	THR
1	A	147	THR
1	A	148	ASP
1	A	149	ASN
1	A	197	SER
1	A	203	SER
1	A	227	MET
1	A	282	ASN
1	A	303	ARG
2	B	323	LEU
2	B	327	ILE
2	B	340	ASP
2	B	380	THR
2	B	433	ASP
2	B	491	ARG
3	H	10	GLU
3	H	24	THR
3	H	31	THR
3	H	78	THR
3	H	81	LEU
3	H	82	ASP
3	H	87	ARG
3	H	121	LEU
4	L	5	THR
4	L	32	THR
4	L	47	ARG
4	L	84	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	L	91	GLN
4	L	107	ILE
4	L	109	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	602	1	14,14,15	0.29	0	17,19,21	0.47	0
5	NAG	B	501	2	14,14,15	0.30	0	17,19,21	0.37	0
5	NAG	A	601	1	14,14,15	0.24	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	602	1	-	0/6/23/26	0/1/1/1
5	NAG	B	501	2	-	1/6/23/26	0/1/1/1
5	NAG	A	601	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	601	NAG	O5-C5-C6-O6
5	A	601	NAG	C4-C5-C6-O6
5	B	501	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/321 (98%)	0.46	16 (5%) 28 19	38, 67, 119, 191	0
2	B	169/177 (95%)	0.65	13 (7%) 13 7	35, 68, 107, 150	0
3	H	127/231 (54%)	0.97	22 (17%) 1 1	47, 84, 129, 173	0
4	L	110/216 (50%)	0.44	6 (5%) 25 16	51, 68, 100, 145	0
All	All	722/945 (76%)	0.59	57 (7%) 12 7	35, 71, 114, 191	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	123	ILE	5.9
3	H	122	VAL	5.7
3	H	11	VAL	5.3
2	B	464	LYS	5.0
4	L	10	THR	5.0
2	B	353	GLU	5.0
2	B	323	LEU	4.8
3	H	29	PHE	4.0
1	A	190	ASN	4.0
3	H	74	ALA	3.8
3	H	121	LEU	3.8
3	H	23	LYS	3.7
4	L	1	GLU	3.7
3	H	94	TYR	3.5
3	H	81	LEU	3.5
3	H	9	ALA	3.5
3	H	120	THR	3.3
1	A	120	ILE	3.1
2	B	485	GLU	3.1
3	H	73	ASP	3.0
2	B	348	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	5	VAL	2.9
2	B	482	LYS	2.8
1	A	151	ALA	2.8
3	H	20	VAL	2.8
3	H	4	LEU	2.8
1	A	187	GLY	2.8
3	H	34	VAL	2.7
1	A	152	PHE	2.6
3	H	8	GLY	2.6
3	H	38	ARG	2.6
4	L	11	LEU	2.6
4	L	78	ARG	2.6
1	A	143	LEU	2.6
2	B	324	PHE	2.6
3	H	43	GLN	2.6
4	L	28	ARG	2.5
2	B	332	GLU	2.5
1	A	160	LYS	2.4
3	H	124	VAL	2.4
3	H	65	GLN	2.4
2	B	465	CYS	2.4
2	B	347	HIS	2.4
1	A	2	LYS	2.3
1	A	238	PHE	2.3
1	A	236	PHE	2.3
1	A	130	ARG	2.3
2	B	325	GLY	2.3
1	A	180	ALA	2.3
1	A	223	PHE	2.3
2	B	327	ILE	2.2
4	L	5	THR	2.2
1	A	186	TYR	2.2
2	B	483	TYR	2.2
3	H	82	ASP	2.2
1	A	164	LYS	2.1
1	A	9	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	601	14/15	0.52	0.35	128,141,151,153	0
5	NAG	A	602	14/15	0.70	0.18	118,128,135,137	0
5	NAG	B	501	14/15	0.82	0.17	72,89,97,104	0

6.5 Other polymers [i](#)

There are no such residues in this entry.