



Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2020 – 08:32 pm BST

PDB ID : 1RCW
Title : Crystal structure of CT610 from Chlamydia trachomatis
Authors : Schwarzenbacher, R.; Liddington, R.C.
Deposited on : 2003-11-04
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	 2% (red), 76% (green), 14% (yellow), 8% (orange), 0% (grey)
1	B	231	 0% (red), 78% (green), 13% (yellow), 7% (orange), 0% (grey)
1	C	231	 4% (red), 75% (green), 16% (yellow), 8% (orange), 0% (grey)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CT610.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	213	Total 1739	C 1115	N 284	O 334	S 2	Se 4	0	0	0
1	B	214	Total 1747	C 1119	N 285	O 337	S 2	Se 4	0	0	0
1	C	213	Total 1700	C 1096	N 276	O 322	S 2	Se 4	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MSE	MET	MODIFIED RESIDUE	UNP O84616
A	79	MSE	MET	MODIFIED RESIDUE	UNP O84616
A	123	MSE	MET	MODIFIED RESIDUE	UNP O84616
A	191	MSE	MET	MODIFIED RESIDUE	UNP O84616
B	21	MSE	MET	MODIFIED RESIDUE	UNP O84616
B	79	MSE	MET	MODIFIED RESIDUE	UNP O84616
B	123	MSE	MET	MODIFIED RESIDUE	UNP O84616
B	191	MSE	MET	MODIFIED RESIDUE	UNP O84616
C	21	MSE	MET	MODIFIED RESIDUE	UNP O84616
C	79	MSE	MET	MODIFIED RESIDUE	UNP O84616
C	123	MSE	MET	MODIFIED RESIDUE	UNP O84616
C	191	MSE	MET	MODIFIED RESIDUE	UNP O84616

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total 2	Fe 2	0	0
2	A	2	Total 2	Fe 2	0	0
2	C	2	Total 2	Fe 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total 71	O 71	0	0
3	B	72	Total 72	O 72	0	0
3	C	26	Total 26	O 26	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	77.55Å 192.97Å 93.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 2.50 29.88 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.88-2.50) 96.9 (29.88-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.1	Depositor
R, R_{free}	0.191 , 0.258 0.195 , 0.256	Depositor DCC
R_{free} test set	1226 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtrriage
Anisotropy	0.471	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5361	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.54 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.7189e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	0/1777	0.90	5/2397 (0.2%)
1	B	0.88	2/1785 (0.1%)	0.93	6/2408 (0.2%)
1	C	0.73	0/1738	0.88	7/2349 (0.3%)
All	All	0.81	2/5300 (0.0%)	0.90	18/7154 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	9	LEU	CG-CD1	-9.62	1.16	1.51
1	B	9	LEU	CG-CD2	-9.01	1.18	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	9	LEU	CD1-CG-CD2	-10.05	80.34	110.50
1	B	9	LEU	CB-CG-CD2	9.19	126.62	111.00
1	A	10	ASP	CB-CG-OD2	7.07	124.66	118.30
1	B	9	LEU	CB-CG-CD1	6.88	122.70	111.00
1	C	76	ASP	CB-CG-OD2	6.86	124.48	118.30
1	B	17	GLN	CG-CD-OE1	6.35	134.30	121.60
1	A	178	ASP	CB-CG-OD2	6.11	123.80	118.30
1	C	195	ASP	CB-CG-OD2	6.07	123.76	118.30
1	C	129	ASP	CB-CG-OD2	6.03	123.72	118.30
1	C	66	ASP	CB-CG-OD2	5.82	123.54	118.30
1	C	154	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	A	167	ASP	CB-CG-OD2	5.70	123.43	118.30
1	B	178	ASP	CB-CG-OD2	5.63	123.36	118.30
1	B	198	ASP	CB-CG-OD2	5.56	123.31	118.30
1	C	13	ASP	CB-CG-OD2	5.32	123.09	118.30
1	C	216	ASP	CB-CG-OD2	5.18	122.96	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	ASP	CB-CG-OD2	5.11	122.90	118.30
1	A	71	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1739	0	1668	33	1
1	B	1747	0	1672	26	1
1	C	1700	0	1616	20	1
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	71	0	0	9	0
3	B	72	0	0	8	1
3	C	26	0	0	1	0
All	All	5361	0	4956	79	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:GLN:NE2	3:A:524:HOH:O	1.81	1.09
1:C:43:TYR:OH	1:C:47:TYR:HE1	1.39	1.03
1:A:104:GLU:OE1	3:A:529:HOH:O	1.82	0.97
1:B:43:TYR:OH	1:B:47:TYR:CE1	2.20	0.94
1:A:43:TYR:OH	1:A:47:TYR:CE1	2.26	0.87
1:B:43:TYR:OH	1:B:47:TYR:HE1	1.58	0.84
1:C:43:TYR:CZ	1:C:47:TYR:HE1	1.97	0.83
1:C:43:TYR:OH	1:C:47:TYR:CE1	2.21	0.82
1:C:43:TYR:CZ	1:C:47:TYR:CE1	2.70	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:TYR:CZ	1:B:47:TYR:CE1	2.72	0.77
1:A:43:TYR:CZ	1:A:47:TYR:CE1	2.75	0.75
1:A:43:TYR:OH	1:A:47:TYR:HE1	1.68	0.74
1:A:7:ASN:N	3:A:506:HOH:O	2.21	0.73
1:B:18:ASN:OD1	3:B:564:HOH:O	2.06	0.72
1:A:72:LYS:HG3	3:B:566:HOH:O	1.89	0.71
1:C:76:ASP:HA	1:C:79:MSE:HE3	1.73	0.69
1:A:21:MSE:HE3	1:A:148:ILE:CD1	2.25	0.65
1:C:43:TYR:HH	1:C:47:TYR:HE1	0.66	0.64
1:B:7:ASN:N	3:B:526:HOH:O	2.30	0.63
1:B:43:TYR:CZ	1:B:47:TYR:HE1	2.14	0.62
1:B:21:MSE:HE3	1:B:148:ILE:CD1	2.29	0.62
1:A:17:GLN:HG2	3:A:560:HOH:O	2.01	0.60
1:A:47:TYR:HE2	1:A:141:TYR:HH	1.49	0.60
1:B:75:LEU:CD2	1:B:79:MSE:SE	3.01	0.58
1:C:7:ASN:N	3:C:426:HOH:O	2.35	0.58
1:A:43:TYR:CZ	1:A:47:TYR:HE1	2.19	0.58
1:C:47:TYR:HE2	1:C:141:TYR:HH	1.53	0.57
1:A:57:TYR:HB3	1:A:138:LEU:HD21	1.89	0.54
1:B:57:TYR:HB3	1:B:138:LEU:HD21	1.89	0.53
1:B:76:ASP:HA	1:B:79:MSE:HE2	1.91	0.53
1:A:17:GLN:CG	3:A:524:HOH:O	2.58	0.52
1:B:43:TYR:CZ	1:B:47:TYR:CD1	2.97	0.52
1:B:124:ARG:HD2	3:B:551:HOH:O	2.10	0.52
1:A:7:ASN:ND2	3:A:570:HOH:O	2.25	0.51
1:A:43:TYR:CZ	1:A:47:TYR:CD1	2.98	0.51
1:A:77:ASN:HB3	3:A:530:HOH:O	2.11	0.51
1:B:75:LEU:HD23	1:B:79:MSE:HG3	1.93	0.51
1:B:7:ASN:ND2	3:B:559:HOH:O	2.35	0.50
1:C:44:ALA:HB1	1:C:96:VAL:HG22	1.94	0.50
1:B:142:GLU:OE2	1:B:185:GLU:OE1	2.30	0.49
1:A:75:LEU:CD2	1:A:79:MSE:SE	3.10	0.49
1:C:174:HIS:HD2	1:C:178:ASP:OD2	1.96	0.48
1:A:72:LYS:CG	3:B:566:HOH:O	2.56	0.48
1:B:22:LEU:HD12	1:B:147:ARG:HD2	1.96	0.47
1:C:43:TYR:CZ	1:C:47:TYR:CD1	3.02	0.47
1:A:142:GLU:OE2	1:A:185:GLU:OE1	2.34	0.46
1:A:22:LEU:HD12	1:A:147:ARG:HD2	1.96	0.46
1:A:75:LEU:HD23	1:A:79:MSE:HG3	1.98	0.46
1:A:76:ASP:HA	1:A:79:MSE:HE2	1.98	0.46
1:A:188:LEU:O	1:A:192:LEU:HD23	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147:ARG:HG2	3:B:569:HOH:O	2.15	0.46
1:B:82:GLU:HG2	3:B:566:HOH:O	2.16	0.45
1:B:47:TYR:HE2	1:B:141:TYR:HH	1.64	0.45
1:C:96:VAL:HG11	1:C:106:LEU:HD13	1.99	0.45
1:A:53:ALA:O	1:A:54:PHE:C	2.55	0.45
1:B:188:LEU:O	1:B:192:LEU:HD23	2.16	0.45
1:A:129:ASP:OD1	1:A:129:ASP:N	2.49	0.45
1:A:47:TYR:HE2	1:A:141:TYR:OH	1.99	0.45
1:C:193:LEU:HD12	1:C:196:ASP:OD1	2.16	0.45
1:A:160:PHE:HE1	3:A:565:HOH:O	2.01	0.44
1:B:22:LEU:HA	1:B:22:LEU:HD23	1.90	0.43
1:C:54:PHE:HA	1:C:57:TYR:CD1	2.54	0.43
1:A:17:GLN:HG3	3:A:524:HOH:O	2.18	0.43
1:C:203:ALA:O	1:C:207:VAL:HG23	2.18	0.42
1:A:9:LEU:HD12	1:A:186:LYS:HG2	2.00	0.42
1:B:53:ALA:O	1:B:54:PHE:C	2.57	0.42
1:C:30:TRP:HE3	1:C:35:LEU:HD12	1.84	0.42
1:A:22:LEU:HA	1:A:22:LEU:HD23	1.89	0.42
1:C:21:MSE:HE3	1:C:148:ILE:CD1	2.50	0.42
1:A:43:TYR:HA	1:A:218:PHE:CE2	2.55	0.41
1:B:43:TYR:HA	1:B:218:PHE:CE2	2.55	0.41
1:B:40:LEU:HD12	1:B:160:PHE:CE2	2.55	0.41
1:B:54:PHE:HA	1:B:57:TYR:CD1	2.56	0.41
1:A:40:LEU:HD12	1:A:160:PHE:CE2	2.56	0.41
1:A:30:TRP:C	1:A:30:TRP:CD1	2.93	0.41
1:C:12:LEU:HD21	1:C:139:TYR:CD2	2.56	0.41
1:C:163:SER:O	1:C:165:PRO:HD3	2.20	0.41
1:C:191:MSE:HE3	1:C:191:MSE:HB2	1.99	0.40
1:B:40:LEU:HD12	1:B:160:PHE:CD2	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:17:GLN:NE2	3:B:514:HOH:O[3_654]	1.44	0.76
1:A:104:GLU:OE2	1:C:85:TYR:O[7_544]	1.65	0.55

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/231 (91%)	208 (99%)	3 (1%)	0	100	100
1	B	212/231 (92%)	208 (98%)	4 (2%)	0	100	100
1	C	211/231 (91%)	205 (97%)	5 (2%)	1 (0%)	29	48
All	All	634/693 (92%)	621 (98%)	12 (2%)	1 (0%)	47	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	161	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/196 (93%)	175 (96%)	7 (4%)	33	58
1	B	183/196 (93%)	176 (96%)	7 (4%)	33	58
1	C	173/196 (88%)	164 (95%)	9 (5%)	23	44
All	All	538/588 (92%)	515 (96%)	23 (4%)	29	53

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	40	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	47	TYR
1	A	104	GLU
1	A	129	ASP
1	A	191	MSE
1	A	209	GLN
1	B	7	ASN
1	B	40	LEU
1	B	47	TYR
1	B	104	GLU
1	B	129	ASP
1	B	191	MSE
1	B	209	GLN
1	C	7	ASN
1	C	40	LEU
1	C	47	TYR
1	C	69	GLU
1	C	75	LEU
1	C	116	LYS
1	C	173	GLU
1	C	195	ASP
1	C	210	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	C	7	ASN
1	C	174	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/231 (90%)	-0.09	4 (1%) 66 69	19, 30, 47, 59	0
1	B	210/231 (90%)	-0.20	2 (0%) 82 84	18, 30, 47, 62	0
1	C	209/231 (90%)	0.18	9 (4%) 35 38	18, 36, 56, 75	0
All	All	628/693 (90%)	-0.04	15 (2%) 59 62	18, 31, 52, 75	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	163	SER	4.4
1	C	195	ASP	4.1
1	B	163	SER	2.9
1	C	177	ALA	2.6
1	A	195	ASP	2.6
1	C	194	LYS	2.5
1	C	198	ASP	2.4
1	A	85	TYR	2.4
1	C	196	ASP	2.4
1	A	32	LYS	2.3
1	C	182	ALA	2.2
1	C	8	PHE	2.1
1	B	220	ASP	2.1
1	C	113	GLU	2.1
1	A	129	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FE	A	401	1/1	0.92	0.09	38,38,38,38	0
2	FE	B	403	1/1	0.95	0.12	40,40,40,40	0
2	FE	C	405	1/1	0.97	0.05	57,57,57,57	0
2	FE	B	404	1/1	0.97	0.07	34,34,34,34	0
2	FE	A	402	1/1	0.98	0.04	33,33,33,33	0
2	FE	C	406	1/1	0.99	0.04	68,68,68,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.