Full wwPDB X-ray Structure Validation Report (i)

Aug 18, 2022 – 01:27 AM EDT

PDB ID	:	4RDK
Title	:	Crystal structure of Norovirus Boxer P domain in complex with Lewis b tetrasaccharide
Authors	:	Hao, N.; Chen, Y.; Xia, M.; Liu, W.; Tan, M.; Jiang, X.; Li, X.
Deposited on	:	2014-09-19
Resolution	:	1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	2%, 91%, • 5%
1	B	308	3%, 90%, 6% 5%
2	C	4	100%
2	D	4	100%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Capsid.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	2270	1448	380	429	13	0	0	0
1	B	294	2282	1454	383	432	13	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	219	GLY	-	expression tag	UNP Q8BCA3
A	220	PRO	-	expression tag	UNP Q8BCA3
A	221	LEU	-	expression tag	UNP Q8BCA3
A	222	GLY	-	expression tag	UNP Q8BCA3
A	223	SER	-	expression tag	UNP Q8BCA3
A	224	PRO	-	expression tag	UNP Q8BCA3
A	225	GLU	-	expression tag	UNP Q8BCA3
A	226	PHE	-	expression tag	UNP Q8BCA3
B	219	GLY	-	expression tag	UNP Q8BCA3
B	220	PRO	-	expression tag	UNP Q8BCA3
B	221	LEU	-	expression tag	UNP Q8BCA3
B	222	GLY	-	expression tag	UNP Q8BCA3
B	223	SER	-	expression tag	UNP Q8BCA3
B	224	PRO	-	expression tag	UNP Q8BCA3
B	225	GLU	-	expression tag	UNP Q8BCA3
B	226	PHE	-	expression tag	UNP Q8BCA3

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	4	46	26	1	19	0	0	0
2	D	4	46	26	1	19	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	352	352	352	0	0
3	B	342	342	342	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Capsid

Chain A:



• Molecule 1: Capsid

Chain B:



• Molecule 2: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain C:



• Molecule 2: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-alpha-D-glucopyranose

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α, β, γ	139.87Å 139.87Å 64.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.30 – 1.63 31.29 – 1.63	Depositor EDS
% Data completeness (in resolution range)	100.0 (31.30-1.63) 100.0 (31.29-1.63)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.24 (at 1.63Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.131 , 0.167 0.127 , 0.162	Depositor DCC
R_{free} test set	4527 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	21.7	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 45.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	5338	wwPDB-VP
Average B, all atoms (Å^2)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NDG, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.33	0/2335	0.52	0/3192
1	B	0.30	0/2348	0.50	0/3211
All	All	0.31	0/4683	0.51	0/6403

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2270	0	2182	7	0
1	B	2282	0	2192	12	0
2	C	46	0	39	0	0
2	D	46	0	39	0	0
3	A	352	0	0	0	0
3	B	342	0	0	1	0
All	All	5338	0	4452	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:506:GLN:HG2	3:B:1013:HOH:O	1.94	0.68
1:B:255:MET:HA	1:B:506:GLN:HG3	1.77	0.66
1:A:259:ASP:OD2	1:A:259:ASP:C	2.38	0.61
1:B:230:LYS:HD2	1:B:462:GLU:O	2.07	0.55
1:B:451:LEU:HD21	1:B:489:MET:HE2	1.92	0.52
1:B:439:PHE:CG	1:B:440:PRO:HD2	2.47	0.49
1:B:261:ASN:HA	1:B:425:PRO:HG2	1.94	0.48
1:B:254:MET:O	1:B:506:GLN:HG3	2.14	0.48
1:A:489:MET:HG2	1:A:495:MET:HB3	1.98	0.46
1:B:469:ASP:HB3	1:B:522:ARG:HB2	1.97	0.46
1:B:451:LEU:HD21	1:B:489:MET:CE	2.45	0.45
1:A:439:PHE:CG	1:A:440:PRO:HD2	2.51	0.45
1:A:498:VAL:HG23	1:A:526:LEU:HB2	1.97	0.45
1:B:359:TYR:O	1:B:360:ASP:HB2	2.18	0.43
1:A:320:PRO:HA	1:A:416:ALA:HB1	2.01	0.42
1:B:256:VAL:H	1:B:506:GLN:NE2	2.18	0.41
1:A:474:HIS:CD2	1:A:486:GLU:HG2	2.56	0.40
1:B:439:PHE:CD1	1:B:440:PRO:HD2	2.56	0.40
1:A:473:LEU:HD11	1:A:489:MET:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	287/308 (93%)	278 (97%)	9 (3%)	0	100	100
1	B	290/308 (94%)	282 (97%)	8 (3%)	0	100	100
All	All	577/616 (94%)	560 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/264 (96%)	250 (99%)	3 (1%)	71	52
1	B	255/264 (97%)	254 (100%)	1 (0%)	91	84
All	All	508/528 (96%)	504 (99%)	4 (1%)	81	69

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	259	ASP
1	A	418	GLN
1	A	441	THR
1	B	441	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	GLN
1	B	474	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NDG	C	1	2	15,15,15	2.08	7 (46%)	21,21,21	1.12	2 (9%)
2	GAL	C	2	2	11,11,12	2.28	3 (27%)	15,15,17	0.97	1 (6%)
2	FUC	C	3	2	10,10,11	1.94	3 (30%)	14,14,16	0.55	0
2	FUC	C	4	2	10,10,11	2.36	2 (20%)	14,14,16	0.81	0
2	NDG	D	1	2	15,15,15	2.08	7 (46%)	21,21,21	1.15	3 (14%)
2	GAL	D	2	2	11,11,12	2.38	3 (27%)	15,15,17	0.86	0
2	FUC	D	3	2	10,10,11	2.04	3 (30%)	14,14,16	0.74	0
2	FUC	D	4	2	10,10,11	2.47	2 (20%)	14,14,16	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDG	C	1	2	-	0/6/26/26	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
2	FUC	C	4	2	-	-	0/1/1/1
2	NDG	D	1	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	FUC	D	3	2	-	-	0/1/1/1
2	FUC	D	4	2	-	-	0/1/1/1

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GAL	C2-C3	-6.18	1.43	1.52
2	D	4	FUC	C2-C3	-5.86	1.43	1.52
2	C	2	GAL	C2-C3	-5.82	1.43	1.52
2	C	4	FUC	C2-C3	-5.61	1.44	1.52
2	D	4	FUC	O5-C1	4.06	1.50	1.43
2	C	4	FUC	O5-C1	3.75	1.49	1.43
2	D	1	NDG	C4-C3	-3.65	1.43	1.52
2	C	1	NDG	C4-C3	-3.54	1.43	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GAL	O5-C1	3.52	1.49	1.43
2	C	2	GAL	O5-C1	3.41	1.49	1.43
2	D	3	FUC	C4-C3	-3.37	1.43	1.52
2	C	3	FUC	C4-C3	-3.30	1.43	1.52
2	D	1	NDG	C3-C2	-3.20	1.47	1.53
2	D	1	NDG	O3-C3	3.11	1.50	1.43
2	D	3	FUC	O5-C5	3.10	1.50	1.43
2	C	1	NDG	O3-C3	3.09	1.50	1.43
2	C	1	NDG	C3-C2	-3.08	1.47	1.53
2	D	1	NDG	C2-N2	2.78	1.50	1.45
2	C	3	FUC	O5-C5	2.74	1.49	1.43
2	D	1	NDG	C7-N2	2.73	1.43	1.34
2	C	1	NDG	O4-C4	2.72	1.49	1.43
2	C	1	NDG	C2-N2	2.70	1.50	1.45
2	C	1	NDG	C7-N2	2.68	1.43	1.34
2	D	1	NDG	O4-C4	2.57	1.49	1.43
2	D	3	FUC	C2-C3	-2.43	1.48	1.52
2	C	3	FUC	C2-C3	-2.39	1.49	1.52
2	C	1	NDG	C1-C2	-2.23	1.50	1.52
2	D	2	GAL	O2-C2	2.19	1.48	1.43
2	D	1	NDG	C1-C2	-2.08	1.50	1.52
2	C	2	GAL	O2-C2	2.07	1.47	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	1	NDG	O5-C1-C2	2.48	112.01	109.52
2	C	2	GAL	C1-C2-C3	2.25	112.44	109.67
2	D	1	NDG	C8-C7-N2	2.06	119.58	116.10
2	C	1	NDG	C8-C7-N2	2.03	119.54	116.10
2	D	1	NDG	C1-C2-C3	2.02	113.30	110.54
2	D	1	NDG	C1-C2-N2	-2.02	108.39	110.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain C



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain D



5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	293/308 (95%)	-0.41	5 (1%)	70	68	16, 21, 42, 70	0
1	B	294/308 (95%)	-0.30	10 (3%)	45	41	16, 23, 42, 70	0
All	All	587/616 (95%)	-0.35	15 (2%)	56	53	16, 22, 42, 70	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	317	PHE	6.1
1	A	504	GLY	5.6
1	B	480	THR	4.7
1	B	230	LYS	4.4
1	B	500	ASN	4.4
1	B	317	PHE	4.3
1	B	479	ASP	4.1
1	B	482	ARG	3.6
1	A	505	PRO	3.2
1	A	506	GLN	2.6
1	B	467	ARG	2.6
1	B	481	HIS	2.5
1	B	231	PRO	2.3
1	A	300	GLY	2.1
1	B	260	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	FUC	D	4	10/11	0.87	0.11	24,27,30,30	0
2	NDG	D	1	15/15	0.92	0.17	28,36,38,43	0
2	FUC	C	4	10/11	0.92	0.07	21,22,24,24	0
2	NDG	C	1	15/15	0.93	0.13	22,29,38,38	0
2	FUC	D	3	10/11	0.95	0.08	28,29,31,31	0
2	GAL	D	2	11/12	0.95	0.10	24,26,28,33	0
2	FUC	C	3	10/11	0.96	0.08	23,25,26,28	0
2	GAL	C	2	11/12	0.96	0.06	21,23,25,28	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.