



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2024 – 10:24 am GMT

PDB ID : 8RKE
Title : Crystal structure of the complete N-terminal region of human ZP2 (hZP2-N1N2N3)
Authors : Fahrenkamp, D.; de Sanctis, D.; Jovine, L.
Deposited on : 2023-12-25
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: grey;"></div> </div>
1	B	335	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 76%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: grey;"></div> </div>
1	C	335	<div style="display: flex; align-items: center;"> <div style="width: 17%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 73%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: grey;"></div> </div>
2	D	6	<div style="display: flex; align-items: center;"> <div style="width: 50%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 50%; height: 10px; background-color: yellow;"></div> </div>
3	E	7	<div style="display: flex; align-items: center;"> <div style="width: 43%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 57%; height: 10px; background-color: yellow;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMA	D	3	-	-	-	X
2	MAN	D	4	-	-	-	X
2	MAN	D	5	-	-	-	X
2	MAN	D	6	-	-	-	X
4	NAG	A	401	-	-	-	X
4	NAG	B	401	-	-	-	X
4	NAG	C	401	-	-	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Zona pellucida sperm-binding protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	289	2256	1442	375	416	23	0	0	0
1	B	279	2185	1398	362	402	23	0	0	0
1	C	277	2175	1392	362	399	22	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	366	LEU	-	expression tag	UNP Q05996
A	367	GLU	-	expression tag	UNP Q05996
A	368	HIS	-	expression tag	UNP Q05996
A	369	HIS	-	expression tag	UNP Q05996
A	370	HIS	-	expression tag	UNP Q05996
A	371	HIS	-	expression tag	UNP Q05996
A	372	HIS	-	expression tag	UNP Q05996
A	373	HIS	-	expression tag	UNP Q05996
B	366	LEU	-	expression tag	UNP Q05996
B	367	GLU	-	expression tag	UNP Q05996
B	368	HIS	-	expression tag	UNP Q05996
B	369	HIS	-	expression tag	UNP Q05996
B	370	HIS	-	expression tag	UNP Q05996
B	371	HIS	-	expression tag	UNP Q05996
B	372	HIS	-	expression tag	UNP Q05996
B	373	HIS	-	expression tag	UNP Q05996
C	366	LEU	-	expression tag	UNP Q05996
C	367	GLU	-	expression tag	UNP Q05996
C	368	HIS	-	expression tag	UNP Q05996
C	369	HIS	-	expression tag	UNP Q05996
C	370	HIS	-	expression tag	UNP Q05996
C	371	HIS	-	expression tag	UNP Q05996
C	372	HIS	-	expression tag	UNP Q05996

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	373	HIS	-	expression tag	UNP Q05996

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	6	72	40	2	30	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	7	83	46	2	35	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0
4	C	1	14	8	1	5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	17	17	17	0	0
5	B	12	12	12	0	0
5	C	2	2	2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Zona pellucida sperm-binding protein 2




- Molecule 1: Zona pellucida sperm-binding protein 2




- Molecule 1: Zona pellucida sperm-binding protein 2





- Molecule 2: α -D-mannopyranose-(1-3)- α -D-mannopyranose-(1-6)-[α -D-mannopyranose-(1-3)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain D: 50% 50%



- Molecule 3: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] α -D-mannopyranose-(1-6)-[α -D-mannopyranose-(1-3)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain E: 43% 57%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	147.68Å 97.77Å 104.33Å 90.00° 94.80° 90.00°	Depositor
Resolution (Å)	21.92 – 2.70 21.92 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.4 (21.92-2.70) 99.5 (21.92-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.71Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158+SVN	Depositor
R, R_{free}	0.229 , 0.258 0.228 , 0.257	Depositor DCC
R_{free} test set	1576 reflections (3.89%)	wwPDB-VP
Wilson B-factor (Å ²)	85.0	Xtrriage
Anisotropy	0.430	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 61.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6886	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/2309	0.56	0/3130
1	B	0.29	0/2234	0.56	0/3025
1	C	0.29	0/2225	0.57	0/3014
All	All	0.29	0/6768	0.56	0/9169

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	0	2227	22	0
1	B	2185	0	2167	15	0
1	C	2175	0	2158	20	0
2	D	72	0	61	0	0
3	E	83	0	70	0	0
4	A	42	0	39	1	0
4	B	28	0	26	1	0
4	C	14	0	13	1	0
5	A	17	0	0	2	0
5	B	12	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	2	0	0	0	0
All	All	6886	0	6761	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:276:THR:HG22	1:B:313:LYS:HG2	1.71	0.73
1:A:229:HIS:ND1	5:A:501:HOH:O	2.29	0.65
1:C:221:PRO:HG2	1:C:224:ALA:HB2	1.84	0.58
1:A:221:PRO:HG2	1:A:224:ALA:HB2	1.83	0.58
1:C:352:MET:HE2	1:C:354:ILE:HG21	1.86	0.58
1:A:276:THR:HG22	1:A:313:LYS:HG2	1.87	0.57
1:C:183:GLY:HA2	1:C:199:LEU:HD21	1.87	0.57
1:C:352:MET:HE2	1:C:354:ILE:CG2	2.35	0.56
1:C:160:MET:HG2	1:C:261:CYS:HB2	1.88	0.56
1:C:302:ASN:O	1:C:320:LEU:HD11	2.05	0.55
1:C:108:ARG:NE	1:C:115:GLN:OE1	2.28	0.55
1:A:86:PRO:HB2	4:A:401:NAG:H83	1.89	0.54
1:A:187:GLU:HB3	1:A:243:LYS:HB3	1.88	0.54
1:C:107:THR:HG22	1:C:116:MET:HB2	1.90	0.54
4:C:401:NAG:H83	4:C:401:NAG:H3	1.90	0.53
1:C:328:LYS:C	1:C:364:VAL:HG23	2.29	0.53
1:A:107:THR:HG22	1:A:116:MET:HB2	1.91	0.52
1:A:82:GLY:O	1:A:119:ARG:NH2	2.42	0.52
1:B:56:ASP:OD1	1:B:57:GLU:N	2.40	0.52
1:C:56:ASP:OD1	1:C:57:GLU:N	2.38	0.52
1:B:53:VAL:HG21	1:B:136:PHE:CD1	2.47	0.50
1:C:300:HIS:ND1	1:C:305:ASP:OD1	2.44	0.49
1:A:56:ASP:OD1	1:A:57:GLU:N	2.42	0.49
1:A:262:ALA:HB3	1:A:352:MET:HE1	1.93	0.49
1:B:162:PHE:HB3	1:B:164:LEU:HD11	1.95	0.49
1:B:205:GLU:HG3	4:B:402:NAG:H81	1.95	0.49
1:C:285:LYS:O	1:C:296:VAL:HG23	2.11	0.49
1:B:262:ALA:HB3	1:B:352:MET:HE1	1.95	0.48
1:C:167:VAL:HA	1:C:216:MET:SD	2.53	0.48
1:A:100:ARG:NH2	5:A:503:HOH:O	2.46	0.47
1:A:160:MET:HG2	1:A:261:CYS:CB	2.45	0.46
1:A:287:VAL:HG13	1:A:339:LEU:HD11	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:186:ILE:HD11	1:B:242:LEU:HB3	1.98	0.46
1:C:102:THR:O	1:C:106:CYS:HB2	2.16	0.46
1:A:139:PRO:HG3	1:A:167:VAL:HG23	1.98	0.45
1:C:184:TRP:H	1:C:199:LEU:HD11	1.82	0.45
1:B:164:LEU:O	1:B:216:MET:N	2.45	0.45
1:C:161:SER:HA	1:C:218:PHE:O	2.17	0.44
1:A:184:TRP:H	1:A:199:LEU:HD11	1.82	0.44
1:B:107:THR:HG22	1:B:116:MET:HB2	1.98	0.44
1:A:160:MET:HG3	1:A:222:PHE:CE1	2.53	0.44
1:A:162:PHE:HB3	1:A:164:LEU:HD11	2.00	0.44
1:B:166:ARG:NH1	1:B:211:ILE:HG22	2.33	0.43
1:B:59:GLU:HG2	1:B:102:THR:HG22	2.01	0.43
1:C:237:LEU:HD13	1:C:262:ALA:HB2	2.01	0.43
1:A:302:ASN:O	1:A:320:LEU:HD11	2.18	0.43
1:C:261:CYS:O	1:C:352:MET:HE1	2.19	0.42
1:B:346:ARG:N	1:B:347:PRO:HD2	2.34	0.42
1:A:53:VAL:HG21	1:A:136:PHE:CD1	2.55	0.42
1:B:262:ALA:H	1:B:352:MET:HE1	1.85	0.42
1:C:237:LEU:CD1	1:C:262:ALA:HB2	2.50	0.41
1:A:102:THR:O	1:A:106:CYS:HB2	2.21	0.41
1:A:204:LYS:O	1:A:204:LYS:HD2	2.20	0.41
1:B:302:ASN:O	1:B:320:LEU:HD11	2.21	0.41
1:C:155:CYS:SG	1:C:261:CYS:O	2.79	0.41
1:A:46:ASN:HB2	1:A:47:PRO:HD3	2.02	0.40
1:A:285:LYS:O	1:A:296:VAL:HG23	2.20	0.40
1:B:287:VAL:HG13	1:B:339:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/335 (83%)	273 (98%)	6 (2%)	0	100	100
1	B	269/335 (80%)	262 (97%)	7 (3%)	0	100	100
1	C	267/335 (80%)	259 (97%)	8 (3%)	0	100	100
All	All	815/1005 (81%)	794 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/298 (87%)	258 (100%)	1 (0%)	91	97
1	B	252/298 (85%)	252 (100%)	0	100	100
1	C	250/298 (84%)	250 (100%)	0	100	100
All	All	761/894 (85%)	760 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	1	1,2	14,14,15	0.42	0	17,19,21	0.42	0
2	NAG	D	2	2	14,14,15	0.21	0	17,19,21	0.57	0
2	BMA	D	3	2	11,11,12	0.58	0	15,15,17	0.75	0
2	MAN	D	4	2	11,11,12	0.68	0	15,15,17	1.06	2 (13%)
2	MAN	D	5	2	11,11,12	0.82	1 (9%)	15,15,17	1.37	2 (13%)
2	MAN	D	6	2	11,11,12	0.74	0	15,15,17	0.95	2 (13%)
3	NAG	E	1	1,3	14,14,15	0.46	0	17,19,21	0.46	0
3	NAG	E	2	3	14,14,15	0.18	0	17,19,21	0.52	0
3	BMA	E	3	3	11,11,12	0.54	0	15,15,17	0.71	0
3	MAN	E	4	3	11,11,12	0.84	0	15,15,17	1.05	2 (13%)
3	MAN	E	5	3	11,11,12	0.74	0	15,15,17	1.00	2 (13%)
3	MAN	E	6	3	11,11,12	0.93	1 (9%)	15,15,17	1.21	2 (13%)
3	MAN	E	7	3	11,11,12	0.70	0	15,15,17	0.95	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	BMA	D	3	2	-	2/2/19/22	0/1/1/1
2	MAN	D	4	2	-	0/2/19/22	0/1/1/1
2	MAN	D	5	2	-	0/2/19/22	0/1/1/1
2	MAN	D	6	2	-	0/2/19/22	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	1/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	E	4	3	-	1/2/19/22	0/1/1/1
3	MAN	E	5	3	-	1/2/19/22	0/1/1/1
3	MAN	E	6	3	-	2/2/19/22	0/1/1/1
3	MAN	E	7	3	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	6	MAN	C1-C2	2.21	1.57	1.52
2	D	5	MAN	C1-C2	2.13	1.57	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	5	MAN	C1-O5-C5	4.39	118.14	112.19
3	E	6	MAN	C1-O5-C5	3.22	116.55	112.19
3	E	4	MAN	C1-O5-C5	2.78	115.95	112.19
2	D	4	MAN	C1-O5-C5	2.48	115.55	112.19
2	D	4	MAN	O2-C2-C3	-2.38	105.38	110.14
3	E	5	MAN	C1-O5-C5	2.30	115.31	112.19
3	E	5	MAN	O2-C2-C3	-2.21	105.71	110.14
2	D	6	MAN	C1-O5-C5	2.20	115.17	112.19
2	D	5	MAN	O2-C2-C3	-2.19	105.75	110.14
3	E	6	MAN	O2-C2-C3	-2.15	105.83	110.14
2	D	6	MAN	O2-C2-C3	-2.14	105.86	110.14
3	E	7	MAN	C1-O5-C5	2.11	115.06	112.19
3	E	4	MAN	O2-C2-C3	-2.05	106.04	110.14
3	E	7	MAN	O2-C2-C3	-2.04	106.05	110.14

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	3	BMA	O5-C5-C6-O6
2	D	3	BMA	C4-C5-C6-O6
3	E	6	MAN	C4-C5-C6-O6
3	E	6	MAN	O5-C5-C6-O6
3	E	2	NAG	C3-C2-N2-C7
2	D	2	NAG	C1-C2-N2-C7
3	E	4	MAN	C4-C5-C6-O6
2	D	2	NAG	C3-C2-N2-C7

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	E	5	MAN	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	401	1	14,14,15	0.58	0	17,19,21	0.59	0
4	NAG	A	402	1	14,14,15	0.46	0	17,19,21	0.43	0
4	NAG	C	401	1	14,14,15	0.37	0	17,19,21	1.30	2 (11%)
4	NAG	B	401	1	14,14,15	0.23	0	17,19,21	0.54	0
4	NAG	B	402	1	14,14,15	0.27	0	17,19,21	0.55	0
4	NAG	A	403	1	14,14,15	0.24	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	401	1	-	1/6/23/26	0/1/1/1
4	NAG	A	402	1	-	1/6/23/26	0/1/1/1
4	NAG	C	401	1	-	5/6/23/26	0/1/1/1
4	NAG	B	401	1	-	4/6/23/26	0/1/1/1
4	NAG	B	402	1	-	1/6/23/26	0/1/1/1
4	NAG	A	403	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	401	NAG	C2-N2-C7	4.34	129.09	122.90
4	C	401	NAG	C1-C2-N2	2.47	114.70	110.49

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	401	NAG	O5-C5-C6-O6
4	A	403	NAG	O5-C5-C6-O6
4	A	403	NAG	C4-C5-C6-O6
4	C	401	NAG	C4-C5-C6-O6
4	B	401	NAG	C4-C5-C6-O6
4	C	401	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	C	401	NAG	O7-C7-N2-C2
4	B	401	NAG	O5-C5-C6-O6
4	A	403	NAG	C1-C2-N2-C7
4	A	402	NAG	O5-C5-C6-O6
4	A	401	NAG	C3-C2-N2-C7
4	B	401	NAG	C3-C2-N2-C7
4	B	401	NAG	C1-C2-N2-C7
4	A	403	NAG	C3-C2-N2-C7
4	B	402	NAG	C3-C2-N2-C7
4	C	401	NAG	C3-C2-N2-C7

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	401	NAG	1	0
4	C	401	NAG	1	0
4	B	402	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	289/335 (86%)	0.02	10 (3%) 44 44	52, 82, 142, 190	0
1	B	279/335 (83%)	0.12	12 (4%) 35 33	53, 90, 155, 206	0
1	C	277/335 (82%)	1.16	58 (20%) 1 0	71, 151, 274, 339	0
All	All	845/1005 (84%)	0.43	80 (9%) 8 6	52, 102, 231, 339	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	359	LEU	11.0
1	C	363	PRO	8.0
1	C	360	CYS	7.9
1	C	355	TYR	7.5
1	C	330	LEU	7.2
1	C	338	SER	6.6
1	C	214	HIS	6.5
1	C	357	GLU	6.5
1	C	332	HIS	6.2
1	B	182	MET	6.1
1	C	331	LEU	6.0
1	C	361	GLU	5.9
1	C	337	ALA	5.6
1	C	301	ASP	5.5
1	C	334	PHE	5.4
1	C	111	HIS	5.3
1	C	166	ARG	5.0
1	C	271	THR	4.9
1	C	211	ILE	4.7
1	C	300	HIS	4.5
1	C	356	PRO	4.4
1	C	73	TRP	4.3
1	C	364	VAL	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	121	MET	4.1
1	C	358	CYS	4.1
1	B	131	ALA	4.0
1	B	132	VAL	3.9
1	C	131	ALA	3.9
1	C	362	SER	3.8
1	C	193	ARG	3.7
1	C	121	MET	3.7
1	A	106	CYS	3.7
1	B	214	HIS	3.7
1	C	298	GLN	3.6
1	C	112	GLY	3.6
1	C	308	ALA	3.4
1	C	323	THR	3.3
1	A	130	GLY	3.3
1	C	346	ARG	3.3
1	C	270	ALA	3.2
1	A	192	ALA	3.1
1	C	58	ARG	3.0
1	B	365	SER	3.0
1	C	213	ASN	2.9
1	C	319	THR	2.9
1	B	74	HIS	2.8
1	C	273	MET	2.8
1	C	333	GLN	2.7
1	C	215	ARG	2.7
1	C	154	ILE	2.6
1	C	276	THR	2.6
1	C	192	ALA	2.6
1	C	317	SER	2.5
1	A	83	LEU	2.5
1	C	328	LYS	2.5
1	A	131	ALA	2.5
1	B	117	THR	2.5
1	C	316	PHE	2.5
1	C	83	LEU	2.4
1	A	46	ASN	2.4
1	A	301	ASP	2.4
1	A	117	THR	2.3
1	C	191	GLY	2.3
1	C	336	LEU	2.2
1	C	327	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	309	THR	2.2
1	C	335	TYR	2.2
1	B	120	VAL	2.2
1	C	110	VAL	2.2
1	B	215	ARG	2.2
1	B	346	ARG	2.2
1	A	67	SER	2.2
1	C	113	GLY	2.1
1	A	121	MET	2.1
1	C	310	ASN	2.1
1	C	157	LYS	2.1
1	B	363	PRO	2.0
1	C	352	MET	2.0
1	C	289	PHE	2.0
1	C	249	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	D	6	11/12	0.51	0.69	223,242,247,249	0
2	MAN	D	4	11/12	0.61	0.47	187,201,215,216	0
2	MAN	D	5	11/12	0.74	0.75	180,203,207,209	0
2	BMA	D	3	11/12	0.74	0.52	203,215,223,234	0
3	MAN	E	7	11/12	0.79	0.25	138,150,161,161	0
2	NAG	D	1	14/15	0.82	0.19	120,130,141,145	0
3	MAN	E	5	11/12	0.84	0.24	102,104,113,115	0
3	BMA	E	3	11/12	0.85	0.27	109,131,147,155	0
3	MAN	E	6	11/12	0.86	0.32	80,97,102,110	0
2	NAG	D	2	14/15	0.89	0.31	148,163,191,202	0
3	NAG	E	2	14/15	0.90	0.24	99,112,126,133	0
3	MAN	E	4	11/12	0.93	0.32	94,103,113,122	0
3	NAG	E	1	14/15	0.94	0.12	73,82,105,106	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	401	14/15	0.58	0.57	159,197,203,205	0
4	NAG	A	401	14/15	0.74	0.52	133,162,179,184	0
4	NAG	C	401	14/15	0.75	0.43	157,174,184,184	0
4	NAG	B	402	14/15	0.79	0.27	101,126,147,148	0
4	NAG	A	403	14/15	0.81	0.23	103,124,138,147	0
4	NAG	A	402	14/15	0.86	0.39	99,135,142,146	0

6.5 Other polymers [i](#)

There are no such residues in this entry.