



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2024 – 10:58 PM EST

PDB ID : 3RO2
Title : Structures of the LGN/NuMA complex
Authors : Shang, Y.; Wei, Z.
Deposited on : 2011-04-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	
2	B	28	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2792 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called G-protein-signaling modulator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	328	2485	1557	444	477	7	0	2	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	GLY	-	expression tag	UNP Q8VDU0
A	14	SER	-	expression tag	UNP Q8VDU0

- Molecule 2 is a protein called peptide of Nuclear mitotic apparatus protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	22	186	116	29	39	2	0	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	96	Total O 96 96	0	0
4	B	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: G-protein-signaling modulator 2



- Molecule 2: peptide of Nuclear mitotic apparatus protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	91.31Å 91.31Å 178.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.14 – 2.30 36.14 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.1 (36.14-2.30) 97.1 (36.14-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.24 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.215 , 0.271 0.246 , 0.301	Depositor DCC
R_{free} test set	1012 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtrriage
Anisotropy	0.013	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2792	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/2537	0.52	1/3432 (0.0%)
2	B	0.57	0/190	0.55	0/257
All	All	0.43	0/2727	0.53	1/3689 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	223	LEU	CA-CB-CG	6.54	130.35	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2485	0	2318	23	0
2	B	186	0	161	4	0
3	A	18	0	24	4	0
4	A	96	0	0	0	0
4	B	7	0	0	0	0
All	All	2792	0	2503	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:235:ARG:HD2	1:A:273:VAL:HG11	1.52	0.89
1:A:63:ASN:HB3	2:B:2004:LEU:HD21	1.71	0.72
1:A:83:THR:O	1:A:87:THR:HG23	2.05	0.57
1:A:131:LYS:HD2	1:A:184:LEU:HD11	1.86	0.56
1:A:37:VAL:HG23	1:A:64:ALA:HB1	1.88	0.56
1:A:291:TYR:O	1:A:295:ILE:HD12	2.06	0.55
1:A:238:TYR:HB2	1:A:261:THR:HG22	1.88	0.54
1:A:143:ASN:HB3	3:A:3:GOL:H32	1.92	0.51
1:A:257:TYR:O	1:A:261:THR:HG23	2.12	0.50
1:A:163:PHE:HB2	1:A:164:PRO:CD	2.44	0.48
1:A:40:PHE:HB3	1:A:61:LEU:HG	1.96	0.48
1:A:177:LEU:HD21	3:A:1:GOL:H32	1.96	0.47
1:A:21:ALA:HB1	2:B:2010:ILE:HD13	1.95	0.47
1:A:163:PHE:HB2	1:A:164:PRO:HD3	1.95	0.47
1:A:258:TYR:HB3	1:A:281:LEU:HG	1.98	0.44
1:A:303:ALA:HA	1:A:306:GLN:OE1	2.19	0.43
1:A:182:LEU:HD13	1:A:197:ALA:HB3	2.00	0.42
1:A:147:ALA:HB2	3:A:3:GOL:H12	2.00	0.42
1:A:61:LEU:HD13	1:A:77:TYR:CE1	2.55	0.41
1:A:238:TYR:CB	1:A:261:THR:HG22	2.50	0.41
1:A:63:ASN:ND2	3:A:2:GOL:O3	2.53	0.41
1:A:63:ASN:HB3	2:B:2004:LEU:CD2	2.45	0.40
1:A:66:PHE:CE1	2:B:2002:GLU:HG2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/338 (96%)	322 (99%)	3 (1%)	1 (0%)	41	50
2	B	20/28 (71%)	18 (90%)	1 (5%)	1 (5%)	2	1
All	All	346/366 (94%)	340 (98%)	4 (1%)	2 (1%)	25	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	1997	CYS
1	A	330	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/265 (88%)	227 (97%)	7 (3%)	41	57
2	B	20/26 (77%)	18 (90%)	2 (10%)	7	9
All	All	254/291 (87%)	245 (96%)	9 (4%)	35	50

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	VAL
1	A	55	SER
1	A	122	LEU
1	A	201	LEU
1	A	221	ARG
1	A	223	LEU
1	A	276	GLN
2	B	1996	THR
2	B	2006	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	181	ASN
1	A	210	ASN
1	A	220	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	2	-	5,5,5	0.35	0	5,5,5	0.57	0
3	GOL	A	1	-	5,5,5	0.36	0	5,5,5	0.57	0
3	GOL	A	3	-	5,5,5	0.37	0	5,5,5	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	2	-	-	2/4/4/4	-
3	GOL	A	1	-	-	4/4/4/4	-
3	GOL	A	3	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1	GOL	O1-C1-C2-C3
3	A	1	GOL	C1-C2-C3-O3
3	A	3	GOL	O1-C1-C2-C3
3	A	3	GOL	C1-C2-C3-O3
3	A	1	GOL	O1-C1-C2-O2
3	A	3	GOL	O2-C2-C3-O3
3	A	2	GOL	C1-C2-C3-O3
3	A	1	GOL	O2-C2-C3-O3
3	A	2	GOL	O2-C2-C3-O3
3	A	3	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	GOL	1	0
3	A	1	GOL	1	0
3	A	3	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/338 (97%)	0.46	39 (11%) 4 6	25, 42, 97, 114	0
2	B	22/28 (78%)	1.62	6 (27%) 0 0	60, 94, 109, 129	0
All	All	350/366 (95%)	0.54	45 (12%) 3 5	25, 45, 100, 129	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1996	THR	6.9
1	A	308	LEU	6.6
1	A	337	PHE	5.3
1	A	328	LEU	5.0
1	A	268	LEU	3.9
1	A	324	ALA	3.6
1	A	153	GLY	3.5
1	A	154	CYS	3.4
2	B	1992	PHE	3.3
1	A	163	PHE	3.3
1	A	321	LEU	3.2
2	B	2007	TRP	3.1
2	B	1997	CYS	3.1
1	A	156	GLY	3.0
1	A	269	LYS	3.0
1	A	344	ILE	3.0
1	A	288	LEU	3.0
1	A	334	ALA	2.9
1	A	335	MET	2.9
1	A	152	PHE	2.8
1	A	157	PRO	2.7
1	A	310	ASP	2.7
2	B	2005	ASP	2.7
1	A	332	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	15	ALA	2.7
1	A	327	ALA	2.6
1	A	320	SER	2.6
1	A	270	ASP	2.5
1	A	313	GLY	2.4
1	A	158	GLN	2.4
1	A	329	GLY	2.3
1	A	287	LEU	2.3
1	A	336	HIS	2.3
1	A	271	ARG	2.3
1	A	222	LEU	2.3
1	A	309	LYS	2.2
1	A	314	GLU	2.2
1	A	323	ASN	2.1
1	A	319	TRP	2.1
1	A	333	GLN	2.1
1	A	303	ALA	2.1
1	A	311	ARG	2.1
1	A	293	LYS	2.0
1	A	307	GLU	2.0
2	B	1995	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	1	6/6	0.93	0.10	41,44,48,53	0
3	GOL	A	3	6/6	0.95	0.22	49,52,57,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	2	6/6	0.96	0.15	36,37,41,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.