



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2020 – 02:10 PM GMT

PDB ID : 6RPJ
Title : A Non-blocking anti-CTLA-4 Nanobody complexed with CTLA-4
Authors : Gao, H.; Zhou, A.; Zhang, F.
Deposited on : 2019-05-14
Resolution : 3.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1191 (3.30-3.22)
Clashscore	141614	1251 (3.30-3.22)
Ramachandran outliers	138981	1229 (3.30-3.22)
Sidechain outliers	138945	1228 (3.30-3.22)
RSRZ outliers	127900	1154 (3.30-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 91%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 40px;">91% • 9%</p>
1	C	129	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 89%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 40px;">89% • 9%</p>
1	E	129	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 4%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 40px;">88% • 8%</p>
1	G	129	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 86%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 40px;">86% • 9%</p>
2	B	142	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 40px;">84% • 12%</p>
2	D	142	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 40px;">85% • 12%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	142	
2	H	142	
3	I	2	
3	J	2	
3	K	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	G	201	-	-	-	X
5	NA	E	202	-	-	-	X
5	NA	G	202	-	-	-	X

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytotoxic T-lymphocyte protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	118	884	556	144	175	9	0	0	0
1	C	118	884	556	144	175	9	0	0	0
1	E	119	891	561	145	176	9	0	0	0
1	G	118	884	556	144	175	9	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	GLY	-	expression tag	UNP P16410
A	123	GLY	-	expression tag	UNP P16410
A	124	SER	-	expression tag	UNP P16410
A	125	HIS	-	expression tag	UNP P16410
A	126	HIS	-	expression tag	UNP P16410
A	127	HIS	-	expression tag	UNP P16410
A	128	HIS	-	expression tag	UNP P16410
A	129	HIS	-	expression tag	UNP P16410
A	130	HIS	-	expression tag	UNP P16410
C	122	GLY	-	expression tag	UNP P16410
C	123	GLY	-	expression tag	UNP P16410
C	124	SER	-	expression tag	UNP P16410
C	125	HIS	-	expression tag	UNP P16410
C	126	HIS	-	expression tag	UNP P16410
C	127	HIS	-	expression tag	UNP P16410
C	128	HIS	-	expression tag	UNP P16410
C	129	HIS	-	expression tag	UNP P16410
C	130	HIS	-	expression tag	UNP P16410
E	122	GLY	-	expression tag	UNP P16410
E	123	GLY	-	expression tag	UNP P16410
E	124	SER	-	expression tag	UNP P16410

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	125	HIS	-	expression tag	UNP P16410
E	126	HIS	-	expression tag	UNP P16410
E	127	HIS	-	expression tag	UNP P16410
E	128	HIS	-	expression tag	UNP P16410
E	129	HIS	-	expression tag	UNP P16410
E	130	HIS	-	expression tag	UNP P16410
G	122	GLY	-	expression tag	UNP P16410
G	123	GLY	-	expression tag	UNP P16410
G	124	SER	-	expression tag	UNP P16410
G	125	HIS	-	expression tag	UNP P16410
G	126	HIS	-	expression tag	UNP P16410
G	127	HIS	-	expression tag	UNP P16410
G	128	HIS	-	expression tag	UNP P16410
G	129	HIS	-	expression tag	UNP P16410
G	130	HIS	-	expression tag	UNP P16410

- Molecule 2 is a protein called A non-blocking CTLA-4 nanobody.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	125	Total 938	C 576	N 167	O 187	S 8	0	2	0
2	D	125	Total 935	C 574	N 167	O 186	S 8	0	1	0
2	F	125	Total 938	C 576	N 167	O 187	S 8	0	2	0
2	H	125	Total 938	C 576	N 167	O 187	S 8	0	2	0

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	I	2	Total 24	C 14	N 1	O 9	0	0	0
3	J	2	Total 24	C 14	N 1	O 9	0	0	0
3	K	2	Total 24	C 14	N 1	O 9	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	C	1	Total 14	C 8	N 1	O 5	0	0
4	C	1	Total 14	C 8	N 1	O 5	0	0
4	E	1	Total 14	C 8	N 1	O 5	0	0
4	G	1	Total 14	C 8	N 1	O 5	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
5	G	1	Total 1	Na 1	0	0
5	E	1	Total 1	Na 1	0	0

- Molecule 6 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	D	1	Total 1	O 1	0	0
6	H	1	Total 1	O 1	0	0



- Molecule 2: A non-blocking CTLA-4 nanobody



- Molecule 2: A non-blocking CTLA-4 nanobody



- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	309.98Å 75.71Å 83.05Å 90.00° 100.05° 90.00°	Depositor
Resolution (Å)	47.57 – 3.25 47.52 – 3.25	Depositor EDS
% Data completeness (in resolution range)	95.3 (47.57-3.25) 95.3 (47.52-3.25)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 3.25Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.249 , 0.262 0.248 , 0.263	Depositor DCC
R_{free} test set	1455 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	83.9	Xtrriage
Anisotropy	0.269	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7424	wwPDB-VP
Average B, all atoms (Å ²)	98.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/899	0.73	0/1225
1	C	0.67	0/899	0.72	0/1225
1	E	0.66	0/907	0.73	0/1237
1	G	0.66	0/899	0.72	0/1225
2	B	0.67	0/963	0.73	0/1302
2	D	0.67	0/957	0.73	0/1294
2	F	0.67	0/963	0.73	0/1302
2	H	0.68	0/963	0.73	0/1302
All	All	0.67	0/7450	0.73	0/10112

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	884	0	869	0	0
1	C	884	0	868	1	0
1	E	891	0	875	2	0
1	G	884	0	868	3	0
2	B	938	0	886	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	935	0	881	1	1
2	F	938	0	886	1	0
2	H	938	0	886	1	0
3	I	24	0	22	0	0
3	J	24	0	22	0	0
3	K	24	0	22	0	0
4	C	28	0	26	0	0
4	E	14	0	13	0	0
4	G	14	0	13	0	0
5	E	1	0	0	0	0
5	G	1	0	0	0	0
6	D	1	0	0	0	0
6	H	1	0	0	0	0
All	All	7424	0	7137	9	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:103:ARG:HD2	2:H:105:TRP:CE2	2.46	0.51
2:B:103:ARG:HD2	2:B:105:TRP:CE2	2.50	0.47
2:D:103:ARG:HD2	2:D:105:TRP:CE2	2.52	0.45
2:F:103:ARG:HD3	2:F:105:TRP:CE2	2.52	0.44
1:E:12:LEU:HD21	1:G:115:TYR:HB3	2.00	0.44
1:C:31:ALA:HB2	1:C:98:LEU:HD23	2.01	0.43
1:G:100:TYR:CG	1:G:101:PRO:HA	2.55	0.42
1:E:115:TYR:HB3	1:G:12:LEU:HD21	2.02	0.42

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:82:GLN:OE1	2:B:82:GLN:OE1[2_557]	2.07	0.13
2:D:82:GLN:OE1	2:D:82:GLN:OE1[2_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/129 (90%)	107 (92%)	9 (8%)	0	100	100
1	C	116/129 (90%)	107 (92%)	9 (8%)	0	100	100
1	E	117/129 (91%)	109 (93%)	8 (7%)	0	100	100
1	G	116/129 (90%)	107 (92%)	9 (8%)	0	100	100
2	B	125/142 (88%)	120 (96%)	5 (4%)	0	100	100
2	D	124/142 (87%)	119 (96%)	5 (4%)	0	100	100
2	F	125/142 (88%)	120 (96%)	5 (4%)	0	100	100
2	H	125/142 (88%)	121 (97%)	4 (3%)	0	100	100
All	All	964/1084 (89%)	910 (94%)	54 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/106 (92%)	97 (99%)	1 (1%)	76	85
1	C	98/106 (92%)	97 (99%)	1 (1%)	76	85
1	E	99/106 (93%)	96 (97%)	3 (3%)	41	67
1	G	98/106 (92%)	95 (97%)	3 (3%)	40	67
2	B	97/109 (89%)	95 (98%)	2 (2%)	53	75
2	D	96/109 (88%)	94 (98%)	2 (2%)	53	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	97/109 (89%)	95 (98%)	2 (2%)	53	75
2	H	97/109 (89%)	93 (96%)	4 (4%)	30	60
All	All	780/860 (91%)	762 (98%)	18 (2%)	50	73

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	MET
2	B	19	THR
2	B	78	THR
1	C	53	THR
2	D	19	THR
2	D	78	THR
1	E	30	LYS
1	E	53	THR
1	E	56	MET
2	F	19	THR
2	F	78	THR
1	G	30	LYS
1	G	53	THR
1	G	56	MET
2	H	19	THR
2	H	21	SER
2	H	72	ARG
2	H	78	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
2	B	84	ASN
1	C	7	GLN
2	D	84	ASN
1	E	7	GLN
2	F	84	ASN
1	G	7	GLN
2	H	84	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	I	1	1,3	14,14,15	0.30	0	17,19,21	1.11	2 (11%)
3	FUC	I	2	3	10,10,11	0.30	0	14,14,16	0.66	0
3	NAG	J	1	1,3	14,14,15	0.36	0	17,19,21	0.81	1 (5%)
3	FUC	J	2	3	10,10,11	0.32	0	14,14,16	0.65	0
3	NAG	K	1	1,3	14,14,15	0.33	0	17,19,21	0.79	1 (5%)
3	FUC	K	2	3	10,10,11	0.32	0	14,14,16	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	I	1	1,3	-	1/6/23/26	0/1/1/1
3	FUC	I	2	3	-	-	0/1/1/1
3	NAG	J	1	1,3	-	2/6/23/26	0/1/1/1
3	FUC	J	2	3	-	-	0/1/1/1
3	NAG	K	1	1,3	-	2/6/23/26	0/1/1/1
3	FUC	K	2	3	-	-	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	1	NAG	O5-C5-C6	2.65	111.35	107.20
3	J	1	NAG	C1-O5-C5	2.60	115.72	112.19
3	I	1	NAG	O5-C1-C2	-2.33	107.62	111.29
3	K	1	NAG	C1-O5-C5	2.15	115.10	112.19

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	K	1	NAG	O5-C5-C6-O6
3	K	1	NAG	C4-C5-C6-O6
3	I	1	NAG	O5-C5-C6-O6
3	J	1	NAG	C4-C5-C6-O6
3	J	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	202	1	14,14,15	0.32	0	17,19,21	0.77	0
4	NAG	C	201	1	14,14,15	0.28	0	17,19,21	1.20	1 (5%)
4	NAG	G	201	1	14,14,15	0.28	0	17,19,21	0.76	0
4	NAG	E	201	1	14,14,15	0.29	0	17,19,21	0.76	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	202	1	-	2/6/23/26	0/1/1/1
4	NAG	C	201	1	-	3/6/23/26	0/1/1/1
4	NAG	G	201	1	-	2/6/23/26	0/1/1/1
4	NAG	E	201	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	201	NAG	O5-C1-C2	-2.43	107.45	111.29
4	E	201	NAG	C1-O5-C5	2.01	114.91	112.19

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	202	NAG	O5-C5-C6-O6
4	C	202	NAG	C4-C5-C6-O6
4	G	201	NAG	C4-C5-C6-O6
4	C	201	NAG	C8-C7-N2-C2
4	C	201	NAG	O7-C7-N2-C2
4	G	201	NAG	O5-C5-C6-O6
4	C	201	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/129 (91%)	0.85	5 (4%) 36 33	100, 142, 188, 206	0
1	C	118/129 (91%)	0.42	2 (1%) 70 67	69, 108, 144, 183	0
1	E	119/129 (92%)	0.10	1 (0%) 86 86	59, 84, 112, 135	1 (0%)
1	G	118/129 (91%)	0.16	1 (0%) 86 86	61, 86, 111, 133	1 (0%)
2	B	125/142 (88%)	0.18	4 (3%) 47 45	64, 84, 115, 134	1 (0%)
2	D	125/142 (88%)	0.25	1 (0%) 86 86	62, 82, 111, 140	0
2	F	125/142 (88%)	0.37	4 (3%) 47 45	57, 94, 125, 138	2 (1%)
2	H	125/142 (88%)	0.46	9 (7%) 15 15	62, 104, 143, 151	1 (0%)
All	All	973/1084 (89%)	0.35	27 (2%) 53 50	57, 94, 149, 206	6 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	74	ASN	4.1
2	H	21	SER	3.7
2	H	8	GLY	3.6
2	F	21	SER	3.3
2	H	13	GLN	2.9
2	D	1	GLN	2.9
2	H	7	SER	2.8
1	A	73	SER	2.8
2	H	9	GLY	2.8
2	H	94	TYR	2.6
1	A	26	ALA	2.5
1	G	120	GLU	2.5
2	B	28	ALA	2.4
1	C	73	SER	2.3
2	H	74	ASN	2.2
2	F	41	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	18	ILE	2.2
2	B	41	PRO	2.2
1	A	59	GLU	2.2
2	B	42	GLY	2.1
2	H	95	TYR	2.1
1	A	98	LEU	2.1
2	F	1	GLN	2.1
1	E	121	PRO	2.1
1	A	18	ILE	2.1
2	B	1	GLN	2.0
2	H	41	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	I	1	14/15	0.80	0.33	112,115,117,117	0
3	NAG	K	1	14/15	0.84	0.18	81,84,90,93	0
3	NAG	J	1	14/15	0.85	0.20	80,83,88,90	0
3	FUC	J	2	10/11	0.86	0.29	93,95,96,97	0
3	FUC	K	2	10/11	0.86	0.41	97,99,101,101	0
3	FUC	I	2	10/11	0.91	0.35	116,117,117,118	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	201	14/15	0.67	0.31	90,92,93,94	0
5	NA	G	202	1/1	0.71	0.47	51,51,51,51	0
4	NAG	C	202	14/15	0.73	0.37	98,103,105,106	0
4	NAG	G	201	14/15	0.77	0.42	96,103,105,106	0
5	NA	E	202	1/1	0.79	0.46	36,36,36,36	0
4	NAG	E	201	14/15	0.83	0.33	86,92,94,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.