

Full wwPDB X-ray Structure Validation Report (i)

Feb 21, 2024 – 09:40 PM EST

PDB ID : 4RRF

Title : Editing domain of threonyl-tRNA synthetase from Methanococcus jannaschii

with L-Ser3AA

Authors: Ahmad, S.; Yerabham, A.S.K.; Kamarthapu, V.; Sankaranarayanan, R.

Deposited on : 2014-11-06

Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar \ resolution} \\ (\#{\rm Entries, \ resolution \ range(\AA)}) \end{array}$
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	141	87%	10%	
1	В	141	87%	10%	•
1	С	141	89%	9%	
1	D	141	80%	16%	
1	Е	141	87%	13%	



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of chain		
1	F	141	% 	18%	



2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 7212 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Threonine–tRNA ligase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace	
1	A	138	Total	С	N	О	S	0	0	0	
1	Λ	130	1091	696	180	210	5	0	U	U	
1	В	137	Total	С	N	О	S	5	0	0	
1	Ъ	137	1082	692	176	209	5	9	U	U	
1	С	139	Total	С	N	О	S	10	10 0	0	0
1		159	1096	699	181	211	5		U	0	
1	D	139	Total	С	N	О	S	5	0	0	
1	D	159	1096	699	181	211	5	9	U	U	
1	Е	140	Total	С	N	О	S	4	0	0	
1	l L	140	1109	707	183	214	5	4	U	U	
1	F	140	Total	С	N	О	S	5	0	0	
1	I.	140	1109	707	183	214	5	0	U	U	

• Molecule 2 is SERINE-3'-AMINOADENOSINE (three-letter code: A3S) (formula: $C_{13}H_{19}N_7O_5$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	С	N	О	0	0
2	Λ	1	25	13	7	5	0	
2	В	1	Total	С	N	О	0	0
	D	1	25	13	7	5	U	U
2	С	1	Total	С	N	Ο	0	0
	O	1	25	13	7	5	U	
2	D	1	Total	С	N	Ο	0	0
	D	1	25	13	7	5	U	
2	E	1	Total	С	N	О	0	0
	Ľ	1	25	13	7	5	0	
2	F	1	Total	С	N	Ο	0	0
	1'	1	25	13	7	5		

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

• Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Me	ol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	:	С	1	Total Mg 1 1	0	0
4	:	D	1	Total Mg 1 1	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	94	Total O 94 94	0	0
5	В	83	Total O 83 83	0	0
5	С	76	Total O 76 76	0	0
5	D	77	Total O 77 77	0	0
5	E	86	Total O 86 86	0	0
5	F	60	Total O 60 60	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Threonine–tRNA ligase



• Molecule 1: Threonine–tRNA ligase







4 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	159.59Å 52.81Å 98.31Å	Depositor	
a, b, c, α , β , γ	90.00° 104.04° 90.00°	Depositor	
Resolution (Å)	25.00 - 1.70	Depositor	
rtesolution (A)	24.65 - 1.70	EDS	
% Data completeness	90.3 (25.00-1.70)	Depositor	
(in resolution range)	90.3 (24.65-1.70)	EDS	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	1.91 (at 1.70Å)	Xtriage	
Refinement program	REFMAC 5.6.0117	Depositor	
D D.	0.194 , 0.239	Depositor	
R, R_{free}	0.194 , 0.239	DCC	
R_{free} test set	3993 reflections (5.01%)	wwPDB-VP	
Wilson B-factor (Å ²)	23.7	Xtriage	
Anisotropy	0.144	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 42.9	EDS	
L-test for twinning ²	$ < L > = 0.46, < L^2> = 0.29$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
F_o, F_c correlation	0.96	EDS	
Total number of atoms	7212	wwPDB-VP	
Average B, all atoms (Å ²)	27.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A3S, CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
IVIOI	Moi Chain		$RMSZ \mid \# Z > 5$		# Z > 5	
1	A	0.82	0/1109	0.91	0/1494	
1	В	0.78	1/1100 (0.1%)	0.85	0/1482	
1	С	0.80	1/1114 (0.1%)	0.86	0/1501	
1	D	0.79	0/1114	0.89	2/1501 (0.1%)	
1	Е	0.78	0/1127	0.84	0/1517	
1	F	0.74	1/1127 (0.1%)	0.82	0/1517	
All	All	0.79	3/6691 (0.0%)	0.86	2/9012 (0.0%)	

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	В	120	TRP	CD2-CE2	5.16	1.47	1.41
1	С	120	TRP	CD2-CE2	5.14	1.47	1.41
1	F	120	TRP	CD2-CE2	5.06	1.47	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	D	36	LEU	CB-CG-CD2	6.58	122.18	111.00
1	D	72	LEU	CB-CG-CD1	5.13	119.73	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1091	0	1109	15	0
1	В	1082	0	1097	10	0
1	С	1096	0	1111	9	0
1	D	1096	0	1111	21	0
1	Ε	1109	0	1128	12	0
1	F	1109	0	1128	19	0
2	A	25	0	19	0	0
2	В	25	0	19	0	0
2	С	25	0	19	0	0
2	D	25	0	19	0	0
2	Ε	25	0	19	0	0
2	F	25	0	19	0	0
3	A	1	0	0	1	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
5	A	94	0	0	4	0
5	В	83	0	0	3	0
5	С	76	0	0	2	0
5	D	77	0	0	4	0
5	Ε	86	0	0	1	0
5	F	60	0	0	3	0
All	All	7212	0	6798	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:16:GLU:HG2	1:B:35:LYS:HG3	1.43	0.97
1:A:115:ARG:NH1	3:A:502:CL:CL	2.46	0.85
1:A:49:GLU:HG2	5:A:676:HOH:O	1.83	0.77
1:E:64:GLU:HA	1:E:67:LYS:NZ	1.99	0.77
1:D:16:GLU:HG2	1:D:35:LYS:HD2	1.68	0.75
1:B:35:LYS:HD2	5:B:381:HOH:O	1.86	0.73
1:F:6:LYS:HD3	1:F:139:LYS:HD3	1.71	0.73
1:D:7:MET:SD	1:D:9:LEU:HD21	2.28	0.73
1:F:55:GLU:O	1:F:59:ILE:HG12	1.90	0.71
1:D:112:ASN:HB3	5:D:335:HOH:O	1.94	0.68
1:D:133:LEU:HD12	1:D:136:LEU:HD21	1.76	0.68



 $Continued\ from\ previous\ page...$

Atom 1		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)	
1:F:92:PRO:O	1:F:96:VAL:HG23	1.95	0.66	
1:C:19:ALA:HB3	1:C:32:LEU:HD22	1.78	0.65	
1:F:96:VAL:HG22	1:F:118:PHE:CD2	2.33	0.64	
1:F:96:VAL:HG22	1:F:118:PHE:CE2	2.36	0.61	
1:F:112:ASN:ND2	5:F:309:HOH:O	2.32	0.61	
1:A:122:LYS:NZ	5:A:607:HOH:O	2.32	0.60	
1:F:36:LEU:HD13	1:F:76:ASN:HB3	1.83	0.59	
1:E:67:LYS:NZ	1:E:67:LYS:HB2	2.17	0.59	
1:B:16:GLU:CG	1:B:35:LYS:HG3	2.28	0.57	
1:A:7:MET:HG2	1:A:9:LEU:CD2	2.36	0.56	
1:D:16:GLU:HG2	1:D:35:LYS:CD	2.36	0.55	
1:A:7:MET:HG2	1:A:9:LEU:HD21	1.88	0.55	
1:A:15:LEU:HD22	1:A:41:ALA:HB1	1.89	0.54	
1:C:9:LEU:HD12	1:C:9:LEU:N	2.23	0.54	
1:D:112:ASN:CG	1:E:110:GLY:O	2.46	0.54	
1:A:15:LEU:HD22	1:A:41:ALA:CB	2.39	0.52	
1:A:93:GLU:HG2	5:A:602:HOH:O	2.09	0.52	
1:D:5:MET:HE1	1:D:98:VAL:HG11	1.92	0.52	
1:E:64:GLU:HA	1:E:67:LYS:HZ2	1.71	0.51	
1:F:73:LYS:N	1:F:73:LYS:HD2	2.26	0.51	
1:D:54:PRO:HG2	5:D:339:HOH:O	2.11	0.50	
1:F:62:VAL:HG13	1:F:106:LEU:CD2	2.42	0.50	
1:A:48:ARG:O	1:A:51:GLU:HG3	2.11	0.50	
1:D:59:ILE:HD13	1:D:105:ILE:HD13	1.94	0.49	
1:D:111:TYR:OH	5:D:376:HOH:O	2.08	0.48	
1:A:131:HIS:HB2	1:A:132:PRO:HD2	1.96	0.48	
1:D:112:ASN:ND2	1:E:110:GLY:O	2.46	0.48	
1:A:126:ILE:HG22	1:D:126:ILE:HG22	1.94	0.48	
1:C:22:LYS:HB3	1:C:22:LYS:HE2	1.72	0.47	
1:A:116:ALA:HB1	1:A:122:LYS:HE3	1.96	0.47	
1:E:5:MET:HB3	1:E:140:ILE:HB	1.96	0.47	
1:C:5:MET:HB3	1:C:140:ILE:HB	1.96	0.47	
1:B:5:MET:HB3	1:B:140:ILE:HB	1.97	0.47	
1:A:115:ARG:NH1	5:A:662:HOH:O	2.36	0.46	
1:D:9:LEU:HD22	1:D:42:CYS:SG	2.55	0.46	
5:E:385:HOH:O	1:F:84:HIS:HD2	1.97	0.46	
1:D:133:LEU:CD1	1:D:136:LEU:HD21	2.43	0.46	
1:D:6:LYS:HD3	1:D:85:LEU:O	2.14	0.46	
1:D:7:MET:HG3	1:D:44:ILE:HG12	1.96	0.46	
1:F:96:VAL:CG2	1:F:118:PHE:CE2	2.98	0.45	
1:E:22:LYS:HG2	1:E:26:ALA:CB	2.45	0.45	



Continued from previous page...

A 4 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:A:7:MET:SD	1:A:9:LEU:HD21	2.55	0.45
1:B:112:ASN:ND2	5:B:352:HOH:O	2.49	0.45
1:B:135:GLU:O	1:C:84:HIS:HE1	1.99	0.45
1:B:16:GLU:HG2	1:B:35:LYS:CG	2.29	0.44
1:D:5:MET:HE1	1:D:98:VAL:CG1	2.47	0.44
1:E:22:LYS:HG2	1:E:26:ALA:HB3	1.98	0.44
1:C:84:HIS:HD2	5:C:341:HOH:O	2.01	0.44
1:D:55:GLU:O	1:D:59:ILE:HG12	2.18	0.44
1:F:15:LEU:HD12	1:F:41:ALA:HB1	2.00	0.43
1:F:28:GLU:HG3	5:F:360:HOH:O	2.17	0.43
1:B:7:MET:HG2	1:B:44:ILE:HG12	1.99	0.43
1:B:108:GLU:HG3	5:B:316:HOH:O	2.18	0.43
1:D:107:LYS:NZ	1:D:113:VAL:H	2.17	0.43
1:E:59:ILE:O	1:E:63:GLU:HG2	2.19	0.42
1:F:67:LYS:HB3	1:F:67:LYS:HE2	1.68	0.42
1:F:6:LYS:HA	1:F:6:LYS:HD2	1.86	0.42
1:F:70:ASN:ND2	5:F:322:HOH:O	2.50	0.42
1:D:77:ILE:O	1:D:113:VAL:HA	2.19	0.42
1:A:93:GLU:HG3	1:A:94:THR:N	2.34	0.42
1:E:6:LYS:HG3	1:E:47:GLU:OE2	2.20	0.41
1:F:66:GLU:OE1	1:F:109:ARG:NH1	2.44	0.41
1:E:62:VAL:HG11	1:E:105:ILE:HG22	2.02	0.41
1:F:117:PRO:HD2	1:F:122:LYS:HD3	2.03	0.41
1:C:35:LYS:C	1:C:35:LYS:HD2	2.40	0.41
1:B:48:ARG:HB3	1:B:88:ASP:OD2	2.21	0.41
1:E:99:LEU:HD23	1:E:99:LEU:HA	1.91	0.40
1:C:36:LEU:HA	1:C:36:LEU:HD12	1.84	0.40
1:C:84:HIS:CD2	5:C:341:HOH:O	2.74	0.40
1:D:138:ARG:CZ	5:D:360:HOH:O	2.69	0.40
1:F:15:LEU:HD12	1:F:41:ALA:CB	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the r	number of residu	ies for which	the backbone	conformation	was
analysed, and the total number of	residues.				

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	136/141 (96%)	133 (98%)	3 (2%)	0	100	100
1	В	135/141 (96%)	131 (97%)	4 (3%)	0	100	100
1	C	137/141 (97%)	134 (98%)	3 (2%)	0	100	100
1	D	137/141 (97%)	132 (96%)	5 (4%)	0	100	100
1	E	138/141 (98%)	134 (97%)	4 (3%)	0	100	100
1	F	138/141 (98%)	134 (97%)	4 (3%)	0	100	100
All	All	821/846 (97%)	798 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	120/123 (98%)	119 (99%)	1 (1%)	81	74
1	В	119/123 (97%)	117 (98%)	2 (2%)	60	46
1	С	120/123 (98%)	117 (98%)	3 (2%)	47	29
1	D	120/123 (98%)	116 (97%)	4 (3%)	38	19
1	E	122/123 (99%)	118 (97%)	4 (3%)	38	19
1	F	122/123 (99%)	116 (95%)	6 (5%)	25	9
All	All	723/738 (98%)	703 (97%)	20 (3%)	43	25

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	GLU
1	В	6	LYS
1	В	22	LYS
1	С	27	GLU



Continued from previous page...

Mol	Chain	Res	Type
1	С	31	ASN
1	С	108	GLU
1	D	36	LEU
1	D	55	GLU
1	D	72	LEU
1	D	104	SER
1	Е	9	LEU
1	Е	30	GLU
1	Е	55	GLU
1	Е	72	LEU
1	F	24	LYS
1	F	36	LEU
1	F	38	GLU
1	F	72	LEU
1	F	73	LYS
1	F	144	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	70	ASN
1	A	112	ASN
1	В	52	ASN
1	В	70	ASN
1	В	112	ASN
1	С	31	ASN
1	C C	52	ASN
1	С	84	HIS
1	С	112	ASN
1	D	71	GLN
1	D	84	HIS
1	Е	70	ASN
1	Е	71	GLN
1	Е	84	HIS
1	Е	112	ASN
1	F	70	ASN
1	F	75	ASN
1	F	84	HIS
1	F	112	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res		Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A3S	В	201	-	24,27,27	1.70	2 (8%)	20,39,39	1.88	3 (15%)
2	A3S	Е	201	-	24,27,27	1.67	2 (8%)	20,39,39	1.40	2 (10%)
2	A3S	С	201	-	24,27,27	1.71	5 (20%)	20,39,39	1.81	3 (15%)
2	A3S	F	201	-	24,27,27	1.94	4 (16%)	20,39,39	1.65	1 (5%)
2	A3S	D	201	-	24,27,27	1.74	2 (8%)	20,39,39	1.96	4 (20%)
2	A3S	A	501	-	24,27,27	1.99	5 (20%)	20,39,39	1.32	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A3S	В	201	-	-	2/12/32/32	0/3/3/3
2	A3S	Е	201	-	-	3/12/32/32	0/3/3/3



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A3S	С	201	-	-	3/12/32/32	0/3/3/3
2	A3S	F	201	-	-	2/12/32/32	0/3/3/3
2	A3S	D	201	-	-	2/12/32/32	0/3/3/3
2	A3S	A	501	-	-	3/12/32/32	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	F	201	A3S	O-C	6.90	1.37	1.23
2	A	501	A3S	O-C	6.44	1.36	1.23
2	Е	201	A3S	O-C	6.37	1.36	1.23
2	D	201	A3S	O-C	6.27	1.35	1.23
2	С	201	A3S	O-C	5.97	1.35	1.23
2	В	201	A3S	O-C	5.26	1.33	1.23
2	В	201	A3S	O4'-C1'	4.97	1.48	1.41
2	A	501	A3S	O4'-C1'	4.28	1.47	1.41
2	D	201	A3S	O4'-C1'	4.21	1.47	1.41
2	F	201	A3S	O4'-C1'	3.90	1.46	1.41
2	A	501	A3S	O4'-C4'	-3.48	1.37	1.45
2	С	201	A3S	O4'-C1'	3.23	1.45	1.41
2	A	501	A3S	C2'-C1'	-3.06	1.49	1.53
2	F	201	A3S	C2-N3	2.43	1.36	1.32
2	F	201	A3S	C4'-C3'	2.27	1.56	1.52
2	Е	201	A3S	O2'-C2'	2.26	1.48	1.43
2	A	501	A3S	C2-N3	2.16	1.35	1.32
2	С	201	A3S	C2'-C1'	-2.07	1.50	1.53
2	С	201	A3S	C3'-N8	2.06	1.49	1.45
2	С	201	A3S	O4'-C4'	2.02	1.49	1.45

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	D	201	A3S	N3-C2-N1	-7.06	117.64	128.68
2	В	201	A3S	N3-C2-N1	-6.74	118.15	128.68
2	С	201	A3S	N3-C2-N1	-5.76	119.68	128.68
2	F	201	A3S	N3-C2-N1	-5.37	120.28	128.68
2	Е	201	A3S	N3-C2-N1	-4.20	122.12	128.68
2	A	501	A3S	O4'-C4'-C5'	-3.29	102.09	109.21
2	В	201	A3S	C2-N1-C6	2.95	123.80	118.75
2	A	501	A3S	N3-C2-N1	-2.85	124.22	128.68
2	D	201	A3S	C2-N1-C6	2.71	123.39	118.75



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	201	A3S	O4'-C4'-C5'	2.60	114.82	109.21
2	В	201	A3S	O4'-C4'-C5'	2.58	114.78	109.21
2	D	201	A3S	O2'-C2'-C1'	-2.39	102.04	110.85
2	Е	201	A3S	CB-CA-C	-2.37	104.43	110.21
2	С	201	A3S	C2-N1-C6	2.34	122.76	118.75
2	D	201	A3S	CB-CA-C	-2.02	105.28	110.21

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	В	201	A3S	N-CA-CB-OG
2	В	201	A3S	C-CA-CB-OG
2	F	201	A3S	O-C-CA-CB
2	С	201	A3S	C-CA-CB-OG
2	A	501	A3S	O-C-CA-N
2	Е	201	A3S	N8-C-CA-N
2	F	201	A3S	N8-C-CA-CB
2	С	201	A3S	N-CA-CB-OG
2	D	201	A3S	N-CA-CB-OG
2	A	501	A3S	O-C-CA-CB
2	Е	201	A3S	O-C-CA-N
2	A	501	A3S	C4'-C3'-N8-C
2	D	201	A3S	C4'-C3'-N8-C
2	Е	201	A3S	C4'-C3'-N8-C
2	С	201	A3S	C4'-C3'-N8-C


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	138/141 (97%)	-0.07	1 (0%) 87 90	17, 24, 34, 41	0
1	В	137/141 (97%)	-0.05	2 (1%) 73 77	17, 24, 39, 48	1 (0%)
1	С	139/141 (98%)	-0.04	1 (0%) 87 90	17, 25, 37, 55	2 (1%)
1	D	139/141 (98%)	-0.07	0 100 100	17, 24, 37, 48	1 (0%)
1	Е	140/141 (99%)	-0.05	2 (1%) 75 79	17, 25, 38, 57	1 (0%)
1	F	140/141 (99%)	0.06	2 (1%) 75 79	20, 28, 40, 61	1 (0%)
All	All	833/846 (98%)	-0.03	8 (0%) 82 85	17, 25, 38, 61	6 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	141	VAL	3.2
1	С	143	LYS	3.1
1	Е	144	GLU	2.9
1	F	144	GLU	2.5
1	F	143	LYS	2.5
1	Е	143	LYS	2.1
1	В	78	VAL	2.0
1	A	10	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	A3S	F	201	25/25	0.92	0.10	20,26,41,54	0
2	A3S	Ε	201	25/25	0.93	0.09	20,24,35,56	0
2	A3S	D	201	25/25	0.93	0.11	19,25,39,53	0
2	A3S	В	201	25/25	0.94	0.09	18,24,42,56	0
2	A3S	С	201	25/25	0.95	0.08	16,24,41,56	0
2	A3S	A	501	25/25	0.95	0.08	19,21,27,28	0
4	MG	D	202	1/1	0.97	0.12	30,30,30,30	0
4	MG	С	202	1/1	0.98	0.12	34,34,34,34	0
3	CL	A	502	1/1	0.99	0.03	26,26,26,26	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around A3S E 201:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A3S D 201:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around A3S B 201:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around A3S C 201:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

