



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2023 – 06:11 AM EDT

PDB ID : 4RZW
Title : Crystal structure of BRAF (R509H) kinase domain bound to AZ628
Authors : Wu, Y.; Gavathiotis, E.
Deposited on : 2014-12-24
Resolution : 3.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	
1	B	281	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	251	2010	1275	358	365	12	0	0	0
1	B	252	2022	1284	359	366	13	0	0	0

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	509	HIS	ARG	engineered mutation	UNP P15056
A	543	ALA	ILE	engineered mutation	UNP P15056
A	544	SER	ILE	engineered mutation	UNP P15056
A	551	LYS	ILE	engineered mutation	UNP P15056
A	562	ARG	GLN	engineered mutation	UNP P15056
A	588	ASN	LEU	engineered mutation	UNP P15056
A	630	SER	LYS	engineered mutation	UNP P15056
A	667	GLU	PHE	engineered mutation	UNP P15056
A	673	SER	TYR	engineered mutation	UNP P15056
A	688	ARG	ALA	engineered mutation	UNP P15056
A	706	SER	LEU	engineered mutation	UNP P15056
A	709	ARG	GLN	engineered mutation	UNP P15056
A	713	GLU	SER	engineered mutation	UNP P15056
A	716	GLU	LEU	engineered mutation	UNP P15056
A	720	GLU	SER	engineered mutation	UNP P15056
A	722	SER	PRO	engineered mutation	UNP P15056
A	723	GLY	LYS	engineered mutation	UNP P15056
B	509	HIS	ARG	engineered mutation	UNP P15056
B	543	ALA	ILE	engineered mutation	UNP P15056
B	544	SER	ILE	engineered mutation	UNP P15056
B	551	LYS	ILE	engineered mutation	UNP P15056
B	562	ARG	GLN	engineered mutation	UNP P15056
B	588	ASN	LEU	engineered mutation	UNP P15056
B	630	SER	LYS	engineered mutation	UNP P15056
B	667	GLU	PHE	engineered mutation	UNP P15056

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	673	SER	TYR	engineered mutation	UNP P15056
B	688	ARG	ALA	engineered mutation	UNP P15056
B	706	SER	LEU	engineered mutation	UNP P15056
B	709	ARG	GLN	engineered mutation	UNP P15056
B	713	GLU	SER	engineered mutation	UNP P15056
B	716	GLU	LEU	engineered mutation	UNP P15056
B	720	GLU	SER	engineered mutation	UNP P15056
B	722	SER	PRO	engineered mutation	UNP P15056
B	723	GLY	LYS	engineered mutation	UNP P15056

- Molecule 2 is 3-(2-cyanopropan-2-yl)-N-{4-methyl-3-[(3-methyl-4-oxo-3,4-dihydroquinazolin-6-yl)amino]phenyl}benzamide (three-letter code: B1E) (formula: C₂₇H₂₅N₅O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			34	27	5	2		
2	B	1	Total	C	N	O	0	0
			34	27	5	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.08Å 114.82Å 56.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.76 – 3.49 47.76 – 3.49	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.76-3.49) 99.5 (47.76-3.49)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.55 (at 3.48Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1819)	Depositor
R, R_{free}	0.247 , 0.280 0.239 , 0.272	Depositor DCC
R_{free} test set	345 reflections (4.60%)	wwPDB-VP
Wilson B-factor (Å ²)	72.3	Xtrriage
Anisotropy	0.508	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 12.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4100	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: B1E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/2052	0.45	0/2764
1	B	0.22	0/2064	0.44	0/2780
All	All	0.22	0/4116	0.45	0/5544

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2010	0	2018	26	0
1	B	2022	0	2035	37	0
2	A	34	0	25	5	0
2	B	34	0	25	2	0
All	All	4100	0	4103	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:661:ASN:HD21	1:B:664:GLN:HG3	1.52	0.73
1:B:461:GLN:HB2	1:B:473:LYS:HB3	1.76	0.67
1:B:681:VAL:HG11	1:B:690:LYS:HD3	1.78	0.65
1:A:532:CYS:HB2	2:A:801:B1E:C9	2.31	0.61
1:B:627:MET:SD	1:B:632:PRO:HG3	2.42	0.60
1:B:580:ASN:N	1:B:580:ASN:OD1	2.35	0.59
1:B:490:PRO:HB2	1:B:495:LEU:HB2	1.85	0.58
1:B:532:CYS:HB2	2:B:801:B1E:N10	2.18	0.58
1:B:454:ASP:OD1	1:B:455:GLY:N	2.37	0.57
1:B:659:ILE:HD12	1:B:674:LEU:HD11	1.87	0.57
1:A:532:CYS:HB2	2:A:801:B1E:N10	2.19	0.57
1:A:540:HIS:HB3	1:A:548:PHE:HE1	1.70	0.56
1:B:651:THR:HG22	1:B:681:VAL:HA	1.89	0.55
1:B:705:PRO:HA	1:B:709:ARG:HH21	1.72	0.55
1:A:668:MET:HE2	1:A:674:LEU:HD22	1.90	0.54
1:A:540:HIS:HB3	1:A:548:PHE:CE1	2.42	0.53
1:A:594:ASP:N	2:A:801:B1E:O23	2.26	0.53
1:A:707:PHE:HA	1:A:710:ILE:HB	1.91	0.53
1:B:482:VAL:HG13	1:B:528:VAL:HG12	1.91	0.53
1:B:468:PHE:CG	1:B:494:GLN:HG3	2.45	0.52
1:B:540:HIS:HB3	1:B:548:PHE:HE2	1.74	0.51
1:A:578:LYS:HE3	1:A:580:ASN:HB3	1.92	0.51
1:B:617:ILE:HG22	1:B:620:MET:HE3	1.92	0.51
1:A:681:VAL:HG21	1:A:690:LYS:HD2	1.93	0.50
1:A:668:MET:HB3	1:A:674:LEU:HB2	1.94	0.50
1:A:695:GLU:HA	1:A:698:LYS:HE3	1.92	0.50
1:B:550:MET:HA	1:B:553:LEU:HD23	1.93	0.50
1:A:623:GLU:HG3	1:A:626:ARG:HH21	1.76	0.50
1:B:707:PHE:HA	1:B:710:ILE:HB	1.93	0.49
1:B:517:MET:HB2	1:B:528:VAL:HG23	1.94	0.49
1:A:619:TRP:HA	1:A:644:ILE:HG13	1.97	0.47
1:B:466:GLY:N	1:B:469:GLY:O	2.45	0.47
1:A:448:ASP:N	1:A:448:ASP:OD1	2.46	0.46
1:A:448:ASP:OD1	1:B:450:TRP:NE1	2.25	0.46
1:B:468:PHE:CD1	1:B:494:GLN:HG3	2.51	0.46
1:A:512:ASN:ND2	1:A:559:GLN:HB3	2.30	0.46
1:B:530:GLN:OE1	1:B:591:LYS:HE2	2.16	0.46
1:B:621:ALA:HB3	1:B:624:VAL:HG23	1.98	0.46
1:A:454:ASP:HB3	1:A:523:PRO:HD3	1.98	0.45
1:A:547:LYS:HE2	1:A:682:ARG:HH22	1.81	0.45
1:B:668:MET:HB3	1:B:674:LEU:HB2	1.99	0.45
1:B:644:ILE:O	1:B:648:GLU:HG3	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:660:ASN:OD1	1:B:660:ASN:N	2.46	0.44
1:B:537:LEU:HD21	1:B:649:LEU:HD21	1.99	0.43
1:B:496:GLN:HG3	1:B:497:ALA:N	2.32	0.43
1:A:540:HIS:O	1:A:544:SER:OG	2.37	0.43
1:B:532:CYS:HB3	1:B:583:PHE:CD1	2.54	0.43
1:B:510:HIS:HB3	1:B:513:ILE:HG12	2.00	0.42
1:B:664:GLN:HB3	1:B:668:MET:HE2	2.01	0.42
1:B:540:HIS:HB3	1:B:548:PHE:CE2	2.54	0.42
1:B:593:GLY:HA2	2:B:801:B1E:N34	2.35	0.42
1:A:578:LYS:HD3	1:A:616:SER:HB3	2.02	0.42
1:B:586:GLU:HB3	1:B:588:ASN:ND2	2.34	0.42
1:A:581:ASN:HB3	1:A:593:GLY:O	2.18	0.42
1:A:618:LEU:HD12	1:A:654:LEU:HD22	2.01	0.41
1:B:713:GLU:H	1:B:713:GLU:HG2	1.59	0.41
2:A:801:B1E:H14	2:A:801:B1E:H22	1.71	0.41
1:A:456:GLN:HB3	1:A:476:TRP:HE1	1.86	0.41
1:B:624:VAL:HA	1:B:632:PRO:HB2	2.03	0.41
1:B:690:LYS:HE3	1:B:690:LYS:HB3	1.91	0.41
1:A:481:ALA:HB2	1:A:531:TRP:HE3	1.86	0.41
1:A:532:CYS:H	2:A:801:B1E:C9	2.34	0.40
1:A:617:ILE:HG13	1:A:662:ARG:NH1	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/281 (87%)	234 (96%)	11 (4%)	0	100	100
1	B	246/281 (88%)	234 (95%)	11 (4%)	1 (0%)	34	72
All	All	491/562 (87%)	468 (95%)	22 (4%)	1 (0%)	47	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	459	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/247 (89%)	212 (96%)	8 (4%)	35	66
1	B	222/247 (90%)	210 (95%)	12 (5%)	22	55
All	All	442/494 (90%)	422 (96%)	20 (4%)	27	61

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	448	ASP
1	A	470	THR
1	A	486	ASN
1	A	522	LYS
1	A	547	LYS
1	A	616	SER
1	A	660	ASN
1	A	713	GLU
1	B	493	GLN
1	B	511	VAL
1	B	524	GLN
1	B	525	LEU
1	B	551	LYS
1	B	553	LEU
1	B	577	LEU
1	B	580	ASN
1	B	586	GLU
1	B	660	ASN
1	B	674	LEU
1	B	713	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	494	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B1E	B	801	-	35,37,37	3.31	11 (31%)	49,54,54	2.42	17 (34%)
2	B1E	A	801	-	35,37,37	3.45	13 (37%)	49,54,54	2.11	12 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B1E	B	801	-	-	4/18/21/21	0/4/4/4
2	B1E	A	801	-	-	5/18/21/21	0/4/4/4

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	B1E	C9-N10	15.11	1.47	1.29
2	B	801	B1E	C9-N10	14.82	1.47	1.29
2	A	801	B1E	C2-N10	5.55	1.49	1.40
2	B	801	B1E	C9-N8	5.51	1.46	1.35
2	B	801	B1E	C29-C25	5.20	1.57	1.53
2	A	801	B1E	C9-N8	5.15	1.45	1.35
2	B	801	B1E	C2-N10	5.11	1.49	1.40
2	A	801	B1E	C29-C25	5.01	1.57	1.53
2	A	801	B1E	C21-N20	4.75	1.48	1.35
2	B	801	B1E	C21-N20	4.48	1.47	1.35
2	A	801	B1E	C22-C21	3.89	1.58	1.50
2	B	801	B1E	C1-C7	3.34	1.54	1.47
2	A	801	B1E	C13-N12	3.32	1.49	1.39
2	B	801	B1E	C5-N12	3.29	1.48	1.40
2	A	801	B1E	C5-N12	3.27	1.48	1.40
2	A	801	B1E	C17-N20	3.22	1.48	1.41
2	A	801	B1E	C1-C7	2.82	1.53	1.47
2	B	801	B1E	C17-N20	2.69	1.47	1.41
2	B	801	B1E	C13-N12	2.60	1.47	1.39
2	A	801	B1E	O11-C7	-2.40	1.17	1.22
2	B	801	B1E	C22-C21	2.22	1.54	1.50
2	A	801	B1E	C1-C2	-2.10	1.37	1.40
2	A	801	B1E	C31-C29	2.04	1.57	1.54
2	B	801	B1E	O23-C21	-2.01	1.19	1.23

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	B1E	C24-C22-C21	-6.73	98.15	120.44
2	A	801	B1E	C28-C22-C21	6.18	140.63	120.62
2	A	801	B1E	C24-C22-C21	-6.10	100.23	120.44
2	B	801	B1E	C28-C22-C21	5.74	139.22	120.62
2	B	801	B1E	N8-C9-N10	-5.51	119.60	126.02
2	A	801	B1E	N8-C9-N10	-5.40	119.73	126.02
2	B	801	B1E	C2-N10-C9	4.61	121.16	116.62
2	B	801	B1E	C6-C5-N12	-4.40	105.43	120.32
2	A	801	B1E	C32-C29-C25	-3.73	104.76	111.52
2	A	801	B1E	C2-N10-C9	3.73	120.29	116.62
2	B	801	B1E	C4-C5-N12	3.47	132.27	120.64
2	B	801	B1E	C32-C29-C30	-3.42	101.77	107.25
2	B	801	B1E	C15-C14-C13	3.18	120.44	117.44
2	A	801	B1E	C6-C5-N12	-3.14	109.69	120.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	B1E	C24-C25-C29	3.04	125.25	120.52
2	A	801	B1E	C15-C14-C13	3.04	120.31	117.44
2	B	801	B1E	C5-N12-C13	-2.98	118.88	126.66
2	B	801	B1E	C28-C22-C24	2.86	122.62	119.24
2	B	801	B1E	C31-C29-C30	2.83	111.78	107.25
2	B	801	B1E	C3-C2-C1	2.75	122.02	119.16
2	B	801	B1E	C22-C24-C25	-2.66	117.46	121.07
2	A	801	B1E	C5-N12-C13	-2.65	119.73	126.66
2	B	801	B1E	C24-C25-C29	2.59	124.55	120.52
2	B	801	B1E	C1-C2-N10	-2.50	118.98	122.54
2	A	801	B1E	C4-C5-N12	2.39	128.65	120.64
2	A	801	B1E	O23-C21-C22	2.36	125.15	120.94
2	B	801	B1E	C22-C21-N20	-2.35	110.75	115.92
2	B	801	B1E	C4-C5-C6	2.22	122.29	119.65
2	A	801	B1E	C9-N8-C7	-2.08	119.37	122.11

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	B1E	C24-C25-C29-C32
2	B	801	B1E	C14-C13-N12-C5
2	A	801	B1E	C14-C13-N12-C5
2	A	801	B1E	C26-C25-C29-C32
2	B	801	B1E	C24-C25-C29-C32
2	A	801	B1E	C18-C13-N12-C5
2	B	801	B1E	C18-C13-N12-C5
2	A	801	B1E	C24-C25-C29-C30
2	B	801	B1E	C24-C25-C29-C30


There are no ring outliers.


2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	B1E	2	0
2	A	801	B1E	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/281 (89%)	0.31	7 (2%) 53 47	44, 63, 94, 105	0
1	B	252/281 (89%)	0.42	7 (2%) 53 47	46, 66, 100, 121	0
All	All	503/562 (89%)	0.36	14 (2%) 53 47	44, 65, 97, 121	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	466	GLY	3.5
1	A	525	LEU	3.2
1	B	659	ILE	3.1
1	B	487	VAL	2.8
1	B	620	MET	2.7
1	A	524	GLN	2.5
1	A	573	ILE	2.5
1	A	467	SER	2.2
1	B	672	GLY	2.2
1	A	528	VAL	2.2
1	A	522	LYS	2.1
1	B	660	ASN	2.1
1	B	650	MET	2.1
1	B	593	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	B1E	B	801	34/34	0.91	0.48	47,50,51,51	0
2	B1E	A	801	34/34	0.94	0.51	51,52,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.