



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 4, 2024 – 09:22 PM EST

PDB ID : 1S02
Title : EFFECTS OF ENGINEERED SALT BRIDGES ON THE STABILITY OF
SUBTILISIN BPN'
Authors : Erwin, C.R.; Barnett, B.L.; Oliver, J.D.; Sullivan, J.F.
Deposited on : 1991-02-20
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	275	81% 18%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUBTILISIN BPN'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	1938	1204	333	396	5	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	GLU	GLN	conflict	UNP P00782
A	271	GLU	GLN	conflict	UNP P00782

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	118	Total	O	0	0
			118	118		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: SUBTILISIN BPN⁷



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	39.30Å 72.90Å 75.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.90	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.169 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2063	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.17	4/1976 (0.2%)	1.51	15/2697 (0.6%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	112	GLU	CB-CG	8.10	1.67	1.52
1	A	34	GLY	N-CA	6.10	1.55	1.46
1	A	87	SER	CA-CB	5.47	1.61	1.52
1	A	258	GLY	N-CA	5.02	1.53	1.46

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	186	ARG	NE-CZ-NH2	-10.64	114.98	120.30
1	A	259	ASP	CB-CG-OD1	10.10	127.39	118.30
1	A	140	ASP	CB-CG-OD1	9.57	126.92	118.30
1	A	181	ASP	CB-CG-OD1	7.73	125.26	118.30
1	A	181	ASP	CB-CG-OD2	-6.32	112.61	118.30
1	A	73	ALA	N-CA-CB	-6.20	101.42	110.10
1	A	112	GLU	CA-CB-CG	-6.01	100.17	113.40
1	A	120	ASP	CB-CG-OD2	-5.78	113.10	118.30
1	A	73	ALA	N-CA-C	5.75	126.53	111.00
1	A	231	ALA	N-CA-CB	5.40	117.66	110.10
1	A	186	ARG	NE-CZ-NH1	5.39	123.00	120.30
1	A	112	GLU	N-CA-CB	5.24	120.03	110.60
1	A	31	ILE	O-C-N	5.21	131.04	122.70
1	A	85	ALA	CB-CA-C	5.06	117.69	110.10
1	A	275	GLN	CA-C-O	-5.04	109.50	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1938	0	1895	32	0
2	A	2	0	0	0	0
3	A	5	0	0	1	0
4	A	118	0	0	3	2
All	All	2063	0	1895	32	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:VAL:HG11	1:A:174:VAL:CG2	2.12	0.80
1:A:13:ALA:HB3	1:A:14:PRO:HD3	1.77	0.65
1:A:112:GLU:HG2	4:A:322:HOH:O	1.97	0.64
1:A:23:GLY:HA2	1:A:236:SER:HB3	1.83	0.59
1:A:126:LEU:C	1:A:126:LEU:HD12	2.23	0.59
1:A:185:GLN:OE1	4:A:378:HOH:O	2.16	0.58
1:A:183:SER:O	1:A:184:ASN:HB2	2.04	0.57
1:A:127:GLY:HA2	1:A:167:TYR:O	2.05	0.57
1:A:13:ALA:N	1:A:14:PRO:CD	2.68	0.56
1:A:64:HIS:HE1	1:A:125:SER:O	1.88	0.55
1:A:126:LEU:HD12	1:A:126:LEU:O	2.08	0.54
1:A:224:SER:HB3	1:A:225:PRO:HD3	1.89	0.53
1:A:43:LYS:HB3	1:A:75:LEU:HD21	1.91	0.53
1:A:224:SER:N	1:A:225:PRO:HD2	2.25	0.52
1:A:139:VAL:HG11	1:A:174:VAL:HG22	1.92	0.51
1:A:238:HIS:ND1	3:A:278:SO4:O1	2.35	0.46
1:A:205:ILE:HG13	1:A:222:MET:HB3	1.98	0.46
1:A:271:GLU:O	4:A:354:HOH:O	2.21	0.46
1:A:205:ILE:O	1:A:216:ALA:HA	2.17	0.45
1:A:31:ILE:HD12	1:A:122:ILE:HG23	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:ILE:HD12	1:A:122:ILE:CG2	2.48	0.44
1:A:36:ASP:OD2	1:A:210:PRO:HA	2.18	0.43
1:A:68:VAL:HG13	1:A:225:PRO:HG3	2.01	0.43
1:A:37:SER:HB3	1:A:58:PHE:HB3	2.00	0.43
1:A:29:ALA:HB2	1:A:119:MET:HG3	2.01	0.42
1:A:71:THR:CG2	1:A:225:PRO:HB2	2.49	0.42
1:A:62:ASN:O	1:A:63:SER:HB2	2.20	0.42
1:A:62:ASN:O	1:A:63:SER:CB	2.68	0.42
1:A:62:ASN:OD1	1:A:64:HIS:HB2	2.21	0.41
1:A:104:TYR:HD2	1:A:104:TYR:HA	1.62	0.41
1:A:25:ASN:HD22	1:A:25:ASN:HA	1.57	0.40
1:A:21:TYR:CZ	1:A:237:LYS:HG3	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:340:HOH:O	4:A:360:HOH:O[3_554]	2.14	0.06
4:A:316:HOH:O	4:A:395:HOH:O[1_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/275 (99%)	262 (96%)	11 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	205/205 (100%)	202 (98%)	3 (2%)	65 62

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	TYR
1	A	222	MET
1	A	240	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	59	GLN
1	A	206	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	278	-	4,4,4	0.74	0	6,6,6	0.74	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	278	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.