



# Full wwPDB X-ray Structure Validation Report i

Oct 15, 2023 – 04:10 AM EDT

PDB ID : 7S49  
Title : Crystal Structure of Inhibitor-bound Galactokinase  
Authors : Whitby, F.G.  
Deposited on : 2021-09-08  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 12257 atoms, of which 5990 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Galactokinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
1	A	391	Total	C 5941	H 1852	N 2975	O 536	S 561	17	133	2	0
1	B	391	Total	C 5920	H 1846	N 2961	O 535	S 562	16	133	1	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	252	ALA	LYS	engineered mutation	UNP P51570
A	253	ALA	GLU	engineered mutation	UNP P51570
B	252	ALA	LYS	engineered mutation	UNP P51570
B	253	ALA	GLU	engineered mutation	UNP P51570

- Molecule 2 is alpha-D-galactopyranose (three-letter code: GLA) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 24	C 6	H 12	O 6	4	0
2	B	1	Total 24	C 6	H 12	O 6	4	0

- Molecule 3 is (4R)-2-[(1,3-benzoxazol-2-yl)amino]-4-(4-chloro-1H-pyrazol-5-yl)-4,6,7,8-tetrahydroquinazolin-5(1H)-one (three-letter code: 4QI) (formula: C<sub>18</sub>H<sub>15</sub>ClN<sub>6</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 42	C 18	Cl 1	H 15	N 6	O 2	0	0
3	B	1	Total 42	C 18	Cl 1	H 15	N 6	O 2	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	B	1	Total Na 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	112	Total O 112 112	0	0
6	B	130	Total O 130 130	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Galactokinase



- Molecule 1: Galactokinase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	129.41Å 129.41Å 240.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.20 30.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.00-2.20) 99.3 (30.00-2.20)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.62 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
$R$ , $R_{free}$	0.158 , 0.226 0.168 , 0.234	Depositor DCC
$R_{free}$ test set	2000 reflections (3.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 39.7	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12257	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GLA, PO4, 4QI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/3020	0.92	2/4095 (0.0%)
1	B	0.77	0/3013	0.95	2/4086 (0.0%)
All	All	0.77	0/6033	0.94	4/8181 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	166	ARG	NE-CZ-NH2	-9.91	115.34	120.30
1	B	342	ARG	NE-CZ-NH2	-5.66	117.47	120.30
1	A	21	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	388	LYS	CB-CA-C	-5.15	100.10	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2966	2975	2963	27	0
1	B	2959	2961	2950	31	0
2	A	12	12	11	0	0
2	B	12	12	11	0	0
3	A	27	15	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	27	15	0	0	0
4	A	15	0	0	1	0
4	B	5	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	112	0	0	3	0
6	B	130	0	0	6	0
All	All	6267	5990	5935	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:GLN:HE22	1:A:273:GLU:HG2	1.49	0.76
1:B:199:LEU:HD12	1:B:210:LEU:HD23	1.77	0.67
1:A:54:PRO:HD2	1:A:199:LEU:O	1.95	0.66
4:A:403:PO4:O3	6:A:501:HOH:O	2.13	0.66
1:B:199:LEU:CD1	1:B:210:LEU:HD23	2.27	0.65
1:A:266:ALA:HB1	1:A:269:LEU:HD13	1.82	0.61
1:A:218[B]:LEU:HD21	1:A:355:LEU:HD11	1.84	0.60
1:A:4:LEU:HG	1:A:377:THR:HG22	1.82	0.60
1:B:260:LEU:O	1:B:264:GLU:HG2	2.02	0.59
1:B:63:LEU:HD11	1:B:129:VAL:HG22	1.84	0.59
1:B:8:GLN:HB3	6:B:575:HOH:O	2.03	0.58
1:A:208:THR:OG1	1:B:210:LEU:HD12	2.02	0.58
1:A:218[A]:LEU:HD11	1:A:300:TYR:CD2	2.38	0.58
1:A:29:GLU:OE1	6:A:502:HOH:O	2.17	0.57
1:A:204:ARG:NH1	1:A:257:GLU:OE2	2.37	0.57
1:B:369:GLN:NE2	1:B:376:ALA:HB3	2.19	0.57
1:A:163:ILE:H	1:A:163:ILE:HD12	1.70	0.56
1:A:77:THR:O	6:A:503:HOH:O	2.18	0.56
1:A:385:ASP:OD2	1:A:388:LYS:NZ	2.41	0.54
1:B:338:VAL:HA	1:B:356:LEU:HB3	1.91	0.52
1:B:190:SER:HB2	6:B:554:HOH:O	2.11	0.51
1:B:51:LEU:HD23	1:B:202:ASP:HA	1.92	0.51
1:B:218:LEU:HD11	1:B:355:LEU:HG	1.92	0.51
1:B:88:LEU:HD22	1:B:104:PRO:HD2	1.92	0.50
1:B:369:GLN:HE22	1:B:376:ALA:HB3	1.76	0.50
1:B:52:VAL:HG23	1:B:54:PRO:HD3	1.93	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:301:ARG:HA	6:B:505:HOH:O	2.13	0.48
1:A:73:VAL:O	1:A:89:GLN:HA	2.13	0.48
1:B:16:ALA:HB2	1:B:130:VAL:CG2	2.44	0.47
1:B:195:LYS:NZ	6:B:511:HOH:O	2.47	0.47
1:B:38:VAL:HG23	1:B:344:THR:HG21	1.95	0.47
1:B:291:ALA:HB1	1:B:303:PHE:CE2	2.49	0.47
1:A:105:ARG:NH1	3:A:402:4QI:O23	2.48	0.46
1:A:51:LEU:HD23	1:A:202:ASP:HA	1.96	0.46
1:B:309:GLU:OE1	6:B:501:HOH:O	2.21	0.45
1:A:43:GLU:HB2	1:A:342:ARG:NH2	2.32	0.45
1:A:98:SER:HB2	1:A:115:GLN:NE2	2.33	0.44
1:B:219:ALA:O	1:B:356:LEU:HD12	2.18	0.44
1:B:219:ALA:HB3	1:B:356:LEU:CD1	2.48	0.43
1:B:223:THR:HG23	1:B:378:PHE:CE1	2.53	0.43
1:A:98:SER:HB2	1:A:115:GLN:CD	2.39	0.43
1:A:160:SER:HA	6:B:511:HOH:O	2.16	0.43
1:A:238:VAL:O	1:A:242:GLN:HG3	2.18	0.43
1:A:42:GLY:O	1:A:52:VAL:HG12	2.18	0.43
1:A:18:ARG:HA	1:A:18:ARG:HD2	1.88	0.43
1:A:361:ALA:HB3	1:A:362:PRO:HD3	2.00	0.42
1:B:162:THR:CG2	1:B:165:ALA:H	2.32	0.42
1:A:218[A]:LEU:HD11	1:A:300:TYR:HD2	1.83	0.42
1:B:84:GLU:OE1	1:B:85:PRO:HA	2.19	0.42
1:A:159:ASP:OD1	1:A:159:ASP:C	2.58	0.42
1:A:82:ALA:HB3	1:A:86:GLN:HE21	1.85	0.41
1:A:223:THR:O	1:A:351:CYS:HA	2.19	0.41
1:B:150:TYR:CE1	1:B:166:ARG:CD	3.03	0.41
1:B:49:GLN:HG3	1:B:254:SER:CB	2.50	0.41
1:B:162:THR:HG23	1:B:164:ALA:N	2.35	0.41
1:B:380:LEU:HD23	1:B:380:LEU:HA	1.95	0.41
1:B:16:ALA:HB2	1:B:130:VAL:HG23	2.03	0.41
1:B:150:TYR:CE1	1:B:166:ARG:HD2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	391/392 (100%)	386 (99%)	5 (1%)	0	100 100
1	B	390/392 (100%)	377 (97%)	12 (3%)	1 (0%)	41 46
All	All	781/784 (100%)	763 (98%)	17 (2%)	1 (0%)	51 60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	214	SER

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	310/309 (100%)	299 (96%)	11 (4%)	36 46
1	B	309/309 (100%)	297 (96%)	12 (4%)	32 41
All	All	619/618 (100%)	596 (96%)	23 (4%)	34 43

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	74	SER
1	A	87	ARG
1	A	188	PHE
1	A	213	LEU
1	A	214	SER
1	A	227	VAL
1	A	230	SER
1	A	248	ARG
1	A	356	LEU
1	A	377	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	8	GLN
1	B	37	ARG
1	B	80	GLU
1	B	97	ARG
1	B	192	MET
1	B	217	LYS
1	B	261	GLU
1	B	286	ARG
1	B	301	ARG
1	B	318	TYR
1	B	355	LEU
1	B	356	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	86	GLN
1	A	172	GLN
1	B	8	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PO4	B	403	-	4,4,4	2.33	1 (25%)	6,6,6	0.26	0
2	GLA	B	401	-	12,12,12	3.05	5 (41%)	17,17,17	2.33	4 (23%)
3	4QI	B	402	-	26,31,31	2.03	8 (30%)	27,45,45	2.91	10 (37%)
4	PO4	A	404	-	4,4,4	0.92	0	6,6,6	0.45	0
2	GLA	A	401	-	12,12,12	3.01	7 (58%)	17,17,17	2.06	5 (29%)
3	4QI	A	402	-	26,31,31	1.90	6 (23%)	27,45,45	2.74	8 (29%)
4	PO4	A	405	-	4,4,4	1.31	1 (25%)	6,6,6	0.37	0
4	PO4	A	403	-	4,4,4	0.92	0	6,6,6	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4QI	B	402	-	-	2/2/42/42	0/5/5/5
2	GLA	A	401	-	-	1/2/22/22	0/1/1/1
3	4QI	A	402	-	-	2/2/42/42	0/5/5/5
2	GLA	B	401	-	-	1/2/22/22	0/1/1/1

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	GLA	O5-C1	-6.01	1.27	1.42
2	A	401	GLA	O5-C1	-5.78	1.28	1.42
2	B	401	GLA	O5-C5	-4.99	1.32	1.44
2	B	401	GLA	O3-C3	-4.92	1.31	1.43
2	A	401	GLA	O5-C5	-4.78	1.32	1.44
3	A	402	4QI	C11-N16	4.76	1.44	1.36
2	A	401	GLA	O3-C3	-4.65	1.32	1.43
3	B	402	4QI	C11-N16	4.59	1.44	1.36
4	B	403	PO4	P-O1	4.53	1.61	1.50
3	A	402	4QI	C15-C14	-4.17	1.30	1.36
3	B	402	4QI	C15-N16	3.88	1.43	1.37
3	B	402	4QI	C19-C14	3.73	1.52	1.45

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	4QI	C15-N16	3.41	1.43	1.37
2	B	401	GLA	C4-C3	3.31	1.60	1.52
2	A	401	GLA	O1-C1	3.18	1.49	1.39
3	B	402	4QI	C15-C14	-3.12	1.32	1.36
3	A	402	4QI	O23-C19	-2.91	1.17	1.23
4	A	405	PO4	P-O1	2.54	1.56	1.50
3	B	402	4QI	C11-N10	2.52	1.40	1.36
3	B	402	4QI	C22-C15	2.49	1.53	1.49
3	B	402	4QI	C1-N9	-2.41	1.31	1.38
2	A	401	GLA	C4-C3	2.40	1.58	1.52
2	B	401	GLA	O1-C1	2.28	1.46	1.39
2	A	401	GLA	C3-C2	2.22	1.58	1.52
3	B	402	4QI	O23-C19	-2.18	1.18	1.23
2	A	401	GLA	C1-C2	2.10	1.57	1.52
3	A	402	4QI	C5-C4	2.09	1.43	1.38
3	A	402	4QI	C22-C15	2.07	1.53	1.49

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	4QI	C13-C14-C19	8.27	127.96	116.84
3	A	402	4QI	C24-C13-C14	7.18	122.53	113.62
2	B	401	GLA	C1-O5-C5	6.97	126.82	113.66
3	B	402	4QI	N10-C11-N12	6.84	129.01	117.79
3	A	402	4QI	N10-C11-N12	6.56	128.56	117.79
3	B	402	4QI	C24-C13-C14	6.25	121.37	113.62
3	A	402	4QI	C13-C14-C19	6.03	124.94	116.84
2	A	401	GLA	C1-O5-C5	5.02	123.14	113.66
3	A	402	4QI	C14-C15-N16	-4.62	115.63	120.55
2	B	401	GLA	C4-C3-C2	-4.22	103.45	110.82
3	A	402	4QI	O23-C19-C14	-4.21	117.05	121.30
2	A	401	GLA	C1-C2-C3	3.87	118.34	110.31
3	B	402	4QI	C22-C15-C14	3.63	127.83	123.49
3	B	402	4QI	C19-C14-C15	-3.47	116.42	119.51
3	B	402	4QI	C8-N10-C11	-3.32	119.12	126.49
2	A	401	GLA	C4-C3-C2	-3.14	105.33	110.82
3	B	402	4QI	C14-C15-N16	-2.91	117.45	120.55
3	A	402	4QI	C19-C14-C15	-2.77	117.04	119.51
3	B	402	4QI	O23-C19-C20	-2.70	116.43	120.86
2	A	401	GLA	O5-C5-C6	2.60	112.90	106.44
2	B	401	GLA	O5-C5-C6	2.47	112.58	106.44
3	A	402	4QI	C22-C15-C14	2.25	126.18	123.49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	GLA	C1-C2-C3	2.15	114.77	110.31
3	A	402	4QI	C20-C19-C14	2.13	121.13	117.96
3	B	402	4QI	C13-C14-C15	-2.12	113.57	119.13
2	A	401	GLA	O2-C2-C3	-2.10	105.49	110.35
3	B	402	4QI	C20-C19-C14	2.04	120.99	117.96

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	GLA	O5-C5-C6-O6
2	B	401	GLA	O5-C5-C6-O6
3	A	402	4QI	N12-C11-N10-C8
3	A	402	4QI	N16-C11-N10-C8
3	B	402	4QI	N12-C11-N10-C8
3	B	402	4QI	N16-C11-N10-C8

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	4QI	1	0
4	A	403	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	391/392 (99%)	-0.05	19 (4%) 29 28	35, 53, 81, 104	0
1	B	391/392 (99%)	0.07	30 (7%) 13 12	33, 57, 101, 128	0
All	All	782/784 (99%)	0.01	49 (6%) 20 19	33, 55, 93, 128	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	344	THR	4.7
1	A	80	GLU	4.4
1	A	231	LEU	4.3
1	B	344	THR	4.2
1	B	345	GLY	3.9
1	A	230	SER	3.9
1	B	370	GLU	3.9
1	B	214	SER	3.9
1	B	301	ARG	3.8
1	A	345	GLY	3.8
1	A	248	ARG	3.8
1	B	42	GLY	3.6
1	B	229	HIS	3.6
1	A	138	GLY	3.5
1	B	231	LEU	3.4
1	A	233	SER	3.3
1	A	351	CYS	3.2
1	B	2	ALA	3.2
1	B	343	MET	3.1
1	B	39	ASN	3.1
1	A	300	TYR	3.0
1	B	342	ARG	3.0
1	B	357	GLU	3.0
1	B	217	LYS	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	366	ARG	3.0
1	B	230	SER	2.9
1	A	241	ARG	2.9
1	B	41	ILE	2.8
1	B	352	THR	2.8
1	A	39	ASN	2.7
1	A	38	VAL	2.7
1	B	353	VAL	2.7
1	B	335	VAL	2.7
1	A	140	SER	2.6
1	B	310	SER	2.6
1	A	245	GLU	2.6
1	B	336	PRO	2.6
1	B	372	TYR	2.6
1	B	43	GLU	2.6
1	B	371	HIS	2.5
1	B	80	GLU	2.5
1	A	336	PRO	2.5
1	A	139	LEU	2.4
1	B	367	HIS	2.4
1	B	40	LEU	2.2
1	A	346	GLY	2.2
1	A	37	ARG	2.1
1	B	334	ALA	2.0
1	B	354	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å²)	Q<0.9
4	PO4	A	404	5/5	0.89	0.28	82,92,111,120	0
4	PO4	A	405	5/5	0.89	0.17	87,92,98,115	0
3	4QI	B	402	27/27	0.93	0.12	47,54,62,73	0
3	4QI	A	402	27/27	0.94	0.12	45,57,65,68	0
5	NA	B	404	1/1	0.96	0.10	65,65,65,65	0
4	PO4	B	403	5/5	0.97	0.16	60,70,76,76	0
2	GLA	A	401	12/12	0.97	0.27	20,50,59,60	4
2	GLA	B	401	12/12	0.98	0.19	20,46,49,52	4
5	NA	A	406	1/1	0.98	0.03	54,54,54,54	0
4	PO4	A	403	5/5	0.98	0.30	53,59,64,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.