

Full wwPDB X-ray Structure Validation Report (i)

Jan 13, 2024 – 07:01 pm GMT

PDB ID : 6SBI

Title: X-ray structure of murine Fumarylacetoacetate hydrolase domain containing

protein 1 (FAHD1) in complex with inhibitor oxalate

Authors: Rupp, B.; Naschberger, A.; Weiss, A.K.H.

Deposited on : 2019-07-21

Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	265	67%	14%	•	18%		
1	В	265	67%	13%		18%		
1	С	265	70%	9%	•	19%		
1	D	265	68%	13%		18%		



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6785 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Acylpyruvase FAHD1, mitochondrial.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	٨	216	Total	С	N	Ο	S	0	0	0
1	A	210	1680	1078	283	308	11	U	0	U
1	В	216	Total	С	N	О	S	0	0	0
1	Ъ	210	1680	1078	283	308	11	U	U	0
1	C	214	Total	С	N	О	S	0	1	0
1		214	1673	1074	282	306	11	U	1	0
1	D	216	Total	С	N	О	S	0		0
1	ש	210	1680	1078	283	308	11	U	U	U

There are 152 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-37	MET	-	initiating methionine	UNP Q8R0F8
A	-36	HIS	-	expression tag	UNP Q8R0F8
A	-35	HIS	-	expression tag	UNP Q8R0F8
A	-34	HIS	-	expression tag	UNP Q8R0F8
A	-33	HIS	-	expression tag	UNP Q8R0F8
A	-32	HIS	-	expression tag	UNP Q8R0F8
A	-31	HIS	-	expression tag	UNP Q8R0F8
A	-30	SER	-	expression tag	UNP Q8R0F8
A	-29	SER	-	expression tag	UNP Q8R0F8
A	-28	GLY	-	expression tag	UNP Q8R0F8
A	-27	LEU	-	expression tag	UNP Q8R0F8
A	-26	VAL	-	expression tag	UNP Q8R0F8
A	-25	PRO	-	expression tag	UNP Q8R0F8
A	-24	ARG	-	expression tag	UNP Q8R0F8
A	-23	GLY	-	expression tag	UNP Q8R0F8
A	-22	SER	-	expression tag	UNP Q8R0F8
A	-21	GLY	-	expression tag	UNP Q8R0F8
A	-20	MET	-	expression tag	UNP Q8R0F8
A	-19	LYS	-	expression tag	UNP Q8R0F8
A	-18	GLU	-	expression tag	UNP Q8R0F8
A	-17	THR	-	expression tag	UNP Q8R0F8



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	-16	ALA	-	expression tag	UNP Q8R0F8
A	-15	ALA	_	expression tag	UNP Q8R0F8
A	-14	ALA	-	expression tag	UNP Q8R0F8
A	-13	LYS	_	expression tag	UNP Q8R0F8
A	-12	PHE	-	expression tag	UNP Q8R0F8
A	-11	GLU	_	expression tag	UNP Q8R0F8
A	-10	ARG	-	expression tag	UNP Q8R0F8
A	-9	GLN	-	expression tag	UNP Q8R0F8
A	-8	HIS	-	expression tag	UNP Q8R0F8
A	-7	MET	-	expression tag	UNP Q8R0F8
A	-6	ASP	-	expression tag	UNP Q8R0F8
A	-5	SER	-	expression tag	UNP Q8R0F8
A	-4	PRO	-	expression tag	UNP Q8R0F8
A	-3	ASP	_	expression tag	UNP Q8R0F8
A	-2	LEU	-	expression tag	UNP Q8R0F8
A	-1	GLY	-	expression tag	UNP Q8R0F8
A	0	THR	-	expression tag	UNP Q8R0F8
В	-37	MET	-	initiating methionine	UNP Q8R0F8
В	-36	HIS	-	expression tag	UNP Q8R0F8
В	-35	HIS	-	expression tag	UNP Q8R0F8
В	-34	HIS	-	expression tag	UNP Q8R0F8
В	-33	HIS	-	expression tag	UNP Q8R0F8
В	-32	HIS	-	expression tag	UNP Q8R0F8
В	-31	HIS	-	expression tag	UNP Q8R0F8
В	-30	SER	-	expression tag	UNP Q8R0F8
В	-29	SER	_	expression tag	UNP Q8R0F8
В	-28	GLY	-	expression tag	UNP Q8R0F8
В	-27	LEU	-	expression tag	UNP Q8R0F8
В	-26	VAL	_	expression tag	UNP Q8R0F8
В	-25	PRO	-	expression tag	UNP Q8R0F8
В	-24	ARG	-	expression tag	UNP Q8R0F8
В	-23	GLY	_	expression tag	UNP Q8R0F8
В	-22	SER	-	expression tag	UNP Q8R0F8
В	-21	GLY	_	expression tag	UNP Q8R0F8
В	-20	MET	-	expression tag	UNP Q8R0F8
В	-19	LYS	-	expression tag	UNP Q8R0F8
В	-18	GLU	-	expression tag	UNP Q8R0F8
В	-17	THR	-	expression tag	UNP Q8R0F8
В	-16	ALA		expression tag	UNP Q8R0F8
В	-15	ALA		expression tag	UNP Q8R0F8
В	-14	ALA	-	expression tag	UNP Q8R0F8
В	-13	LYS		expression tag	UNP Q8R0F8



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	-12	PHE	-	expression tag	UNP Q8R0F8
В	-11	GLU	-	expression tag	UNP Q8R0F8
В	-10	ARG	-	expression tag	UNP Q8R0F8
В	-9	GLN	_	expression tag	UNP Q8R0F8
В	-8	HIS	-	expression tag	UNP Q8R0F8
В	-7	MET	-	expression tag	UNP Q8R0F8
В	-6	ASP	-	expression tag	UNP Q8R0F8
В	-5	SER	-	expression tag	UNP Q8R0F8
В	-4	PRO	-	expression tag	UNP Q8R0F8
В	-3	ASP	-	expression tag	UNP Q8R0F8
В	-2	LEU	-	expression tag	UNP Q8R0F8
В	-1	GLY	-	expression tag	UNP Q8R0F8
В	0	THR	-	expression tag	UNP Q8R0F8
С	-37	MET	-	initiating methionine	UNP Q8R0F8
С	-36	HIS	-	expression tag	UNP Q8R0F8
С	-35	HIS	-	expression tag	UNP Q8R0F8
С	-34	HIS	-	expression tag	UNP Q8R0F8
С	-33	HIS	-	expression tag	UNP Q8R0F8
С	-32	HIS	-	expression tag	UNP Q8R0F8
С	-31	HIS	-	expression tag	UNP Q8R0F8
С	-30	SER	-	expression tag	UNP Q8R0F8
С	-29	SER	-	expression tag	UNP Q8R0F8
С	-28	GLY	-	expression tag	UNP Q8R0F8
С	-27	LEU	-	expression tag	UNP Q8R0F8
С	-26	VAL	_	expression tag	UNP Q8R0F8
С	-25	PRO	-	expression tag	UNP Q8R0F8
С	-24	ARG	-	expression tag	UNP Q8R0F8
С	-23	GLY	-	expression tag	UNP Q8R0F8
С	-22	SER	-	expression tag	UNP Q8R0F8
С	-21	GLY	-	expression tag	UNP Q8R0F8
С	-20	MET	-	expression tag	UNP Q8R0F8
С	-19	LYS	-	expression tag	UNP Q8R0F8
С	-18	GLU	-	expression tag	UNP Q8R0F8
С	-17	THR	-	expression tag	UNP Q8R0F8
С	-16	ALA	-	expression tag	UNP Q8R0F8
С	-15	ALA	-	expression tag	UNP Q8R0F8
С	-14	ALA	-	expression tag	UNP Q8R0F8
С	-13	LYS	-	expression tag	UNP Q8R0F8
С	-12	PHE	-	expression tag	UNP Q8R0F8
С	-11	GLU	-	expression tag	UNP Q8R0F8
С	-10	ARG	-	expression tag	UNP Q8R0F8
С	-9	GLN	-	expression tag	UNP Q8R0F8



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	-8	HIS	-	expression tag	UNP Q8R0F8
С	-7	MET	-	expression tag	UNP Q8R0F8
С	-6	ASP	-	expression tag	UNP Q8R0F8
С	-5	SER	-	expression tag	UNP Q8R0F8
С	-4	PRO	-	expression tag	UNP Q8R0F8
С	-3	ASP	-	expression tag	UNP Q8R0F8
С	-2	LEU	=	expression tag	UNP Q8R0F8
С	-1	GLY	-	expression tag	UNP Q8R0F8
С	0	THR	-	expression tag	UNP Q8R0F8
D	-37	MET	-	initiating methionine	UNP Q8R0F8
D	-36	HIS	-	expression tag	UNP Q8R0F8
D	-35	HIS	-	expression tag	UNP Q8R0F8
D	-34	HIS	-	expression tag	UNP Q8R0F8
D	-33	HIS	-	expression tag	UNP Q8R0F8
D	-32	HIS	-	expression tag	UNP Q8R0F8
D	-31	HIS	-	expression tag	UNP Q8R0F8
D	-30	SER	-	expression tag	UNP Q8R0F8
D	-29	SER	-	expression tag	UNP Q8R0F8
D	-28	GLY	-	expression tag	UNP Q8R0F8
D	-27	LEU	-	expression tag	UNP Q8R0F8
D	-26	VAL	-	expression tag	UNP Q8R0F8
D	-25	PRO	-	expression tag	UNP Q8R0F8
D	-24	ARG	1	expression tag	UNP Q8R0F8
D	-23	GLY	-	expression tag	UNP Q8R0F8
D	-22	SER	-	expression tag	UNP Q8R0F8
D	-21	GLY	-	expression tag	UNP Q8R0F8
D	-20	MET	-	expression tag	UNP Q8R0F8
D	-19	LYS	-	expression tag	UNP Q8R0F8
D	-18	GLU	1	expression tag	UNP Q8R0F8
D	-17	THR	-	expression tag	UNP Q8R0F8
D	-16	ALA	-	expression tag	UNP Q8R0F8
D	-15	ALA	1	expression tag	UNP Q8R0F8
D	-14	ALA	-	expression tag	UNP Q8R0F8
D	-13	LYS	-	expression tag	UNP Q8R0F8
D	-12	PHE	-	expression tag	UNP Q8R0F8
D	-11	GLU		expression tag	UNP Q8R0F8
D	-10	ARG	-	expression tag	UNP Q8R0F8
D	-9	GLN		expression tag	UNP Q8R0F8
D	-8	HIS		expression tag	UNP Q8R0F8
D	-7	MET	-	expression tag	UNP Q8R0F8
D	-6	ASP	-	expression tag	UNP Q8R0F8
D	-5	SER	-	expression tag	UNP Q8R0F8



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-4	PRO	-	expression tag	UNP Q8R0F8
D	-3	ASP	-	expression tag	UNP Q8R0F8
D	-2	LEU	-	expression tag	UNP Q8R0F8
D	-1	GLY	-	expression tag	UNP Q8R0F8
D	0	THR	-	expression tag	UNP Q8R0F8

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	В	1	Total Mg 1 1	0	0
2	С	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0

• Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0
3	С	1	Total K 1 1	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total Cl 3 3	0	0
4	В	2	Total Cl 2 2	0	0
4	С	3	Total Cl 3 3	0	0
4	D	1	Total Cl 1 1	0	0

• Molecule 5 is OXALATE ION (three-letter code: OXL) (formula: C_2O_4) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 2 4	0	0
5	В	1	Total C O 6 2 4	0	0
5	С	1	Total C O 6 2 4	0	0
5	D	1	Total C O 6 2 4	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	8	Total O 8 8	0	0
6	В	12	Total O 12 12	0	0
6	С	6	Total O 6 6	0	0
6	D	7	Total O 7 7	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acylpyruvase FAHD1, mitochondrial









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	52.60Å 103.35Å 86.60Å	Donositon
a, b, c, α , β , γ	90.00° 90.21° 90.00°	Depositor
Resolution (Å)	44.37 - 2.70	Depositor
Resolution (A)	44.37 - 2.70	EDS
% Data completeness	62.4 (44.37-2.70)	Depositor
(in resolution range)	$62.4 \ (44.37 - 2.70)$	EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.29 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
υ .	0.187 , 0.234	Depositor
R, R_{free}	0.192 , 0.237	DCC
R_{free} test set	756 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.30 , -11.1	EDS
L-test for twinning ²	$< L > = 0.46, < L^2> = 0.28$	Xtriage
Estimated twinning fraction	0.206 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6785	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.50% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG, OXL, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.67	0/1719	0.86	0/2328
1	В	0.68	0/1719	0.84	0/2328
1	С	0.67	0/1712	0.86	0/2317
1	D	0.68	0/1719	0.84	0/2328
All	All	0.68	0/6869	0.85	0/9301

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1680	0	1716	19	0
1	В	1680	0	1716	15	0
1	С	1673	0	1709	12	0
1	D	1680	0	1716	14	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	1	0	0	0	0
4	A	3	0	0	1	0
4	В	2	0	0	1	0
4	С	3	0	0	1	0
4	D	1	0	0	1	0
5	A	6	0	0	0	0
5	В	6	0	0	0	0
5	С	6	0	0	0	0
5	D	6	0	0	0	0
6	A	8	0	0	0	0
6	В	12	0	0	0	0
6	С	6	0	0	1	0
6	D	7	0	0	0	0
All	All	6785	0	6857	60	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:ARG:NH2	1:A:18:GLU:OE1	2.37	0.58
1:B:30:TYR:O	1:B:34:VAL:HG23	2.05	0.57
1:A:30:TYR:O	1:A:34:VAL:HG23	2.07	0.54
1:C:30:TYR:O	1:C:34:VAL:HG23	2.08	0.54
1:D:30:TYR:O	1:D:34:VAL:HG23	2.07	0.54
1:A:44:GLU:OE1	1:A:174:TYR:OH	2.27	0.53
1:D:32:ASP:HB2	1:D:197:LYS:HB3	1.92	0.52
1:A:34:VAL:HG12	1:A:39:SER:O	2.10	0.52
1:A:32:ASP:HB2	1:A:197:LYS:HB3	1.91	0.51
1:B:31:ALA:O	1:B:35:LYS:HG3	2.11	0.51
1:B:32:ASP:HB2	1:B:197:LYS:HB3	1.93	0.50
1:B:34:VAL:HG12	1:B:39:SER:O	2.12	0.49
1:B:164:LYS:HB3	4:B:303:CL:CL	2.50	0.49
1:C:50:LYS:HD3	1:C:132:CYS:HB2	1.94	0.49
1:A:23:ILE:HD11	1:A:55:TYR:CZ	2.48	0.49
1:D:110:ASP:HB2	4:D:302:CL:CL	2.49	0.49
1:C:32:ASP:HB2	1:C:197:LYS:HB3	1.94	0.48
1:D:23:ILE:HD11	1:D:55:TYR:CZ	2.49	0.48
1:D:34:VAL:HG12	1:D:39:SER:O	2.13	0.47
1:B:23:ILE:HD11	1:B:55:TYR:CZ	2.49	0.47



Continued from previous page...

Continuea from pre		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:32:ASP:OD2	1:C:197:LYS:HE3	2.15	0.47
1:C:23:ILE:HD11	1:C:55:TYR:CZ	2.50	0.46
1:A:15:ARG:HD2	1:A:18:GLU:OE2	2.14	0.46
1:A:44:GLU:OE2	1:A:44:GLU:HA	2.16	0.46
1:B:64:MET:HE2	1:B:223:LYS:O	2.16	0.46
1:C:75:VAL:CG2	1:C:102:LEU:HD11	2.46	0.46
1:A:75:VAL:CG2	1:A:102:LEU:HD11	2.46	0.46
1:A:42:LEU:HD12	1:A:42:LEU:H	1.80	0.46
1:B:75:VAL:CG2	1:B:102:LEU:HD11	2.45	0.46
1:C:34:VAL:HG12	1:C:39:SER:O	2.16	0.45
1:C:44:GLU:HA	1:C:44:GLU:OE2	2.18	0.44
1:C:62:VAL:HG22	1:C:133:PRO:HG3	1.99	0.44
1:C:35:LYS:HE2	1:C:35:LYS:HB2	1.69	0.44
1:D:75:VAL:CG2	1:D:102:LEU:HD11	2.47	0.44
1:A:28:ARG:HD2	1:A:42:LEU:HD11	1.99	0.43
1:B:50:LYS:HD3	1:B:132:CYS:HB2	1.99	0.43
1:A:143:ILE:HA	1:A:144:PRO:HD2	1.92	0.43
1:C:58:GLU:HA	6:C:403:HOH:O	2.18	0.43
1:A:50:LYS:HD3	1:A:132:CYS:HB2	2.01	0.43
1:C:110:ASP:HB2	4:C:303:CL:CL	2.56	0.43
1:B:38:ARG:CZ	1:B:38:ARG:HB2	2.49	0.43
1:D:172:ILE:HB	1:D:173:PRO:HD3	2.00	0.42
1:B:15:ARG:HD2	1:B:18:GLU:OE2	2.18	0.42
1:B:44:GLU:OE1	1:B:44:GLU:HA	2.20	0.42
1:A:62:VAL:HG22	1:A:133:PRO:HG3	2.02	0.42
1:D:50:LYS:HD3	1:D:132:CYS:HB2	2.01	0.42
1:D:44:GLU:HA	1:D:44:GLU:OE1	2.20	0.41
1:D:143:ILE:HA	1:D:144:PRO:HD2	1.96	0.41
1:A:23:ILE:HD11	1:A:55:TYR:CE1	2.55	0.41
1:A:74:GLU:O	1:A:104:LEU:HA	2.20	0.41
1:B:74:GLU:O	1:B:104:LEU:HA	2.21	0.41
1:D:23:ILE:HD11	1:D:55:TYR:CE1	2.56	0.41
1:A:24:VAL:O	1:A:191:ILE:HA	2.20	0.41
1:A:108:ALA:HB1	4:A:304:CL:CL	2.58	0.41
1:B:151:LEU:O	1:B:162:GLU:HA	2.21	0.41
1:B:23:ILE:HD11	1:B:55:TYR:CE1	2.56	0.41
1:D:24:VAL:O	1:D:191:ILE:HA	2.21	0.41
1:A:172:ILE:HB	1:A:173:PRO:HD3	2.02	0.40
1:D:84:ARG:NH1	1:D:184:THR:HG21	2.37	0.40
1:D:124:LEU:H	1:D:124:LEU:HD12	1.86	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	214/265~(81%)	204 (95%)	10 (5%)	0	100	100
1	В	214/265 (81%)	204 (95%)	10 (5%)	0	100	100
1	С	213/265 (80%)	202 (95%)	11 (5%)	0	100	100
1	D	$214/265 \ (81\%)$	203 (95%)	11 (5%)	0	100	100
All	All	855/1060 (81%)	813 (95%)	42 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	centiles
1	A	185/226~(82%)	171 (92%)	14 (8%)	13	30
1	В	185/226 (82%)	168 (91%)	17 (9%)	9	21
1	С	184/226 (81%)	167 (91%)	17 (9%)	9	21
1	D	185/226 (82%)	167 (90%)	18 (10%)	8	19
All	All	739/904 (82%)	673 (91%)	66 (9%)	10	22

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	26	VAL



Continued from previous page...

	Continued from previous page					
Mol	Chain	Res	Type			
1	A	32	ASP			
1	A	42	LEU			
1	A	43	SER			
1	A A	80	LEU			
1	A	113	GLU			
1	A	164	LYS			
1	A A	177	SER			
1	A A	187	GLU			
1	A	197	LYS			
1	A A	203	LYS			
1	A	223	LYS			
1	A	224	ARG			
1	В	10	THR			
1	В	26	VAL			
1	В	32	ASP			
1	В	37	MET			
1	В	62	VAL			
1	В	63	LEU			
1	В	80	LEU			
1	В	113	GLU			
1	В	141	GLU			
1	В	164	LYS			
1	В	177	SER			
1	В	195	THR			
1	В	197	LYS			
1	В	202	ILE			
1	В	203	LYS			
1	В	221	LYS			
1	В	225	SER			
1		13	LEU			
1	C C C C C C C C	26	VAL			
1	С	32	ASP LYS			
1	С	35	LYS			
1	С	39	SER LEU			
1	С	80	LEU			
1	С	86	GLU GLU			
1	С	113	GLU			
1	С	164	LYS			
1	С	177	SER			
1	С	195	THR			
1	С	197	LYS			
1	С	203	LYS			



Continued from previous page...

Mol	Chain	Res	Type
1	C C C	219	ARG
1	С	221[A]	LYS
1	С	221[B]	LYS
1	С	224	ARG
1	D	10	THR
1	D	26	VAL
1	D	32	ASP
1	D	35	LYS
1	D	64	MET
1	D	69	ARG
1	D	80	LEU
1	D	83	LYS
1	D	84	ARG
1	D	113	GLU
1	D	177	SER
1	D	195	THR
1	D	197	LYS
1	D	202	ILE
1	D	203	LYS
1	D	216	VAL
1	D	224	ARG
1	D	225	SER

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such side chains are listed below:

Mol	Chain	Res	Type
1	A	205	ASN
1	В	205	ASN
1	D	205	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 15 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Trino	Chain	Res	Link	Bond lengths			Bond angles		
	туре				Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	OXL	С	306	2	5,5,5	1.42	2 (40%)	6,6,6	1.21	0
5	OXL	В	304	2	5,5,5	1.41	2 (40%)	6,6,6	1.24	0
5	OXL	D	303	2	5,5,5	1.28	1 (20%)	6,6,6	1.27	0
5	OXL	A	306	2	5,5,5	1.58	2 (40%)	6,6,6	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OXL	С	306	2	-	0/4/4/4	-
5	OXL	В	304	2	-	0/4/4/4	-
5	OXL	D	303	2	-	0/4/4/4	-
5	OXL	A	306	2	-	0/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
5	A	306	OXL	O3-C1	-2.79	1.22	1.30
5	С	306	OXL	O4-C2	-2.30	1.23	1.30
5	В	304	OXL	O3-C1	-2.19	1.24	1.30
5	В	304	OXL	O4-C2	-2.19	1.24	1.30
5	D	303	OXL	O4-C2	-2.09	1.24	1.30
5	A	306	OXL	O4-C2	-2.08	1.24	1.30
5	С	306	OXL	O3-C1	-2.05	1.24	1.30



There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(\AA^2)$	Q<0.9
1	A	216/265 (81%)	-0.69	1 (0%) 91 92	14, 32, 69, 103	0
1	В	216/265 (81%)	-0.74	0 100 100	12, 28, 62, 87	0
1	С	$214/265 \ (80\%)$	-0.72	0 100 100	10, 30, 59, 84	0
1	D	$216/265\ (81\%)$	-0.70	1 (0%) 91 92	13, 30, 63, 102	0
All	All	862/1060 (81%)	-0.71	2 (0%) 95 96	10, 30, 63, 103	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	38	ARG	2.6
1	A	38	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
4	CL	В	303	1/1	0.94	0.15	74,74,74,74	0
4	CL	A	305	1/1	0.96	0.08	33,33,33,33	0
4	CL	С	303	1/1	0.97	0.08	54,54,54,54	0
4	CL	С	304	1/1	0.97	0.14	37,37,37,37	0
2	MG	С	301	1/1	0.98	0.04	28,28,28,28	0
3	K	С	302	1/1	0.98	0.06	41,41,41,41	0
5	OXL	A	306	6/6	0.98	0.09	24,33,39,42	0
5	OXL	В	304	6/6	0.98	0.09	29,35,36,38	0
5	OXL	D	303	6/6	0.98	0.14	34,43,46,46	0
4	CL	В	302	1/1	0.99	0.05	31,31,31,31	0
2	MG	D	301	1/1	0.99	0.04	24,24,24,24	0
3	K	A	302	1/1	0.99	0.06	27,27,27,27	0
2	MG	В	301	1/1	0.99	0.07	22,22,22,22	0
4	CL	С	305	1/1	0.99	0.05	22,22,22,22	0
4	CL	D	302	1/1	0.99	0.05	40,40,40,40	0
4	CL	A	303	1/1	0.99	0.05	39,39,39,39	0
4	CL	A	304	1/1	0.99	0.04	42,42,42,42	0
5	OXL	С	306	6/6	0.99	0.08	35,42,51,52	0
2	MG	A	301	1/1	0.99	0.05	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MG C 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around OXL D 303: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



















6.5 Other polymers (i)

There are no such residues in this entry.

