



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2022 – 01:09 PM EDT

PDB ID : 5SEC
Title : Crystal Structure of human phosphodiesterase 10 in complex with 2-[2-(1,4-diphenylimidazol-2-yl)ethyl]-3-methylquinoxaline
Authors : Joseph, C.; Groebke-Zbinden, K.; Benz, J.; Schlatter, D.; Rudolph, M.G.
Deposited on : 2022-01-21
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.31.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	 84% 7% 9%
1	B	343	 83% 8% 8%
1	C	343	 85% 5% 9%
1	D	343	 83% 8% 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	2552	1631	435	462	24	0	2	0
1	B	315	2563	1637	438	464	24	0	1	0
1	C	313	2549	1629	435	461	24	0	1	0
1	D	312	2532	1619	432	457	24	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233
B	447	GLY	-	expression tag	UNP Q9Y233
B	448	SER	-	expression tag	UNP Q9Y233
C	447	GLY	-	expression tag	UNP Q9Y233
C	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0
2	D	1	Total 1	Zn 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

- Molecule 4 is 2-[2-(1,4-diphenyl-1H-imidazol-2-yl)ethyl]-3-methylquinoxaline (three-letter code: IEX) (formula: C₂₆H₂₂N₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N 30 26 4	0	0
4	B	1	Total C N 30 26 4	0	0
4	C	1	Total C N 30 26 4	0	0
4	D	1	Total C N 30 26 4	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	105	Total 105	O 105	0	0
5	B	125	Total 125	O 125	0	0
5	C	124	Total 124	O 124	0	0
5	D	61	Total 61	O 61	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	135.75Å 135.75Å 235.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.80 – 2.03 43.80 – 2.03	Depositor EDS
% Data completeness (in resolution range)	95.1 (43.80-2.03) 95.1 (43.80-2.03)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.03Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.181 , 0.225 0.189 , 0.229	Depositor DCC
R_{free} test set	5228 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	35.6	Xtrriage
Anisotropy	0.174	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10739	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IEX, ZN, MG, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.88	1/2609 (0.0%)	0.94	2/3529 (0.1%)
1	B	0.83	0/2617	0.92	1/3540 (0.0%)
1	C	0.83	2/2603 (0.1%)	0.93	3/3521 (0.1%)
1	D	0.85	0/2583	0.89	0/3495
All	All	0.85	3/10412 (0.0%)	0.92	6/14085 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	592	GLU	CD-OE2	7.75	1.34	1.25
1	C	624	GLU	CD-OE1	-5.27	1.19	1.25
1	A	471	GLU	CD-OE1	5.27	1.31	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	642	ARG	NE-CZ-NH2	-5.99	117.31	120.30
1	B	627	ARG	NE-CZ-NH1	5.79	123.19	120.30
1	C	510	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	C	719	LYS	CB-CA-C	-5.69	99.02	110.40
1	A	496	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	C	642	ARG	NE-CZ-NH2	-5.46	117.57	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2552	0	2529	19	0
1	B	2563	0	2534	19	0
1	C	2549	0	2524	10	0
1	D	2532	0	2507	20	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	30	0	0	0	0
4	B	30	0	0	0	0
4	C	30	0	0	1	0
4	D	30	0	0	0	0
5	A	105	0	0	0	0
5	B	125	0	0	3	0
5	C	124	0	0	2	0
5	D	61	0	0	2	0
All	All	10739	0	10094	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:767:ARG:NH2	1:B:769:GLU:OE1	1.98	0.95
1:C:714:MET:HE2	5:C:1001:HOH:O	1.69	0.90
1:A:497:SER:O	1:A:553:ARG:HD2	1.72	0.89
1:D:714:MET:HE2	5:D:949:HOH:O	1.76	0.84
1:D:461:GLN:HE21	1:D:500:THR:HG21	1.51	0.75
1:C:714:MET:CE	5:C:1001:HOH:O	2.33	0.74
1:B:509:CME:HB2	5:B:1010:HOH:O	1.91	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:461:GLN:HE22	1:A:500:THR:HG21	1.65	0.62
1:B:727:LEU:HD12	1:B:759:LEU:HD11	1.82	0.60
1:A:497:SER:O	1:A:553:ARG:CD	2.49	0.60
1:C:716:ARG:O	1:C:719:LYS:HG3	2.02	0.60
1:C:700:GLY:HA3	1:C:714:MET:O	2.06	0.55
1:A:735:ILE:HB	1:A:736:PRO:HD3	1.88	0.54
1:B:546:THR:HG21	1:D:767:ARG:HH21	1.71	0.54
1:B:497:SER:O	1:B:553:ARG:HD2	2.07	0.53
1:B:700:GLY:HA3	1:B:714:MET:O	2.09	0.53
1:B:727:LEU:HD12	1:B:759:LEU:CD1	2.39	0.52
1:A:461:GLN:NE2	1:A:461:GLN:HA	2.24	0.52
1:B:467:ARG:O	1:B:471:GLU:HG3	2.09	0.52
1:D:461:GLN:NE2	1:D:500:THR:HG21	2.22	0.52
1:D:497:SER:O	1:D:553:ARG:HD2	2.09	0.52
1:D:627:ARG:O	1:D:631:ILE:HG12	2.10	0.51
1:A:662:ARG:HG2	1:A:662:ARG:NH1	2.25	0.51
1:B:727:LEU:CD1	1:B:759:LEU:HD11	2.40	0.51
1:C:562:CYS:HB3	1:C:565:LEU:HD12	1.93	0.50
1:A:730:TYR:HA	1:A:734:ALA:HB3	1.93	0.50
1:D:700:GLY:HA3	1:D:714:MET:O	2.13	0.48
1:D:727:LEU:C	1:D:727:LEU:HD23	2.33	0.48
1:A:662:ARG:HG2	1:A:662:ARG:HH11	1.79	0.48
1:A:662:ARG:CG	1:A:662:ARG:HH11	2.26	0.48
1:D:647:GLU:O	1:D:651:THR:HB	2.13	0.48
1:B:770:GLU:HG3	1:B:771:THR:N	2.29	0.47
1:B:545:HIS:HD2	5:B:959:HOH:O	1.97	0.47
1:C:542:GLN:NE2	1:C:542:GLN:HA	2.30	0.47
1:A:544:ASN:HB3	1:A:547:LEU:HD13	1.98	0.46
1:C:727:LEU:HD12	1:C:759:LEU:HD11	1.98	0.46
1:D:547:LEU:N	1:D:547:LEU:HD12	2.31	0.46
1:A:749:GLU:N	1:A:750:PRO:CD	2.79	0.45
1:B:497:SER:OG	1:B:557:LEU:HD11	2.17	0.45
1:D:493:MET:SD	1:D:535:HIS:HA	2.57	0.45
1:A:461:GLN:HE22	1:A:500:THR:CG2	2.30	0.44
1:A:461:GLN:NE2	1:A:461:GLN:CA	2.80	0.44
1:A:492:TYR:CZ	1:A:496:ARG:HD2	2.53	0.43
1:A:497:SER:OG	1:A:557:LEU:HD11	2.18	0.43
1:C:693:TYR:OH	4:C:803:IEX:N18	2.51	0.43
1:A:461:GLN:NE2	1:A:500:THR:HG21	2.31	0.43
1:D:497:SER:O	1:D:553:ARG:CD	2.67	0.42
1:C:638:TYR:OH	1:C:642:ARG:HD3	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:477:PHE:HB3	1:B:535:HIS:CE1	2.54	0.42
1:D:492:TYR:CZ	1:D:496:ARG:HD2	2.55	0.42
1:B:770:GLU:CG	1:B:771:THR:N	2.82	0.42
1:B:642:ARG:NH2	5:B:909:HOH:O	2.53	0.41
1:D:714:MET:CE	5:D:949:HOH:O	2.51	0.41
1:A:542:GLN:NE2	1:A:542:GLN:HA	2.35	0.41
1:D:533:VAL:HG13	1:D:673:CYS:HB3	2.01	0.41
1:C:565:LEU:O	1:C:595:HIS:ND1	2.44	0.41
1:D:457:GLN:HA	1:D:460:MET:CE	2.51	0.41
1:B:546:THR:HG21	1:D:767:ARG:NH2	2.35	0.41
1:B:551:LEU:HD23	1:B:551:LEU:HA	1.82	0.41
1:B:562:CYS:HB3	1:B:599:THR:OG1	2.20	0.41
1:B:611:PHE:HB3	1:B:614:LEU:HD22	2.03	0.41
1:D:479:ILE:HD11	1:D:528:LYS:HG3	2.03	0.41
1:D:752:LEU:HD21	1:D:756:ARG:NH1	2.35	0.41
1:A:523:PRO:HD2	1:A:695:GLU:HG2	2.03	0.40
1:A:722:VAL:N	1:A:723:PRO:CD	2.83	0.40
1:D:497:SER:HA	1:D:542:GLN:HE22	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/343 (91%)	303 (97%)	9 (3%)	0	100	100
1	B	313/343 (91%)	305 (97%)	8 (3%)	0	100	100
1	C	311/343 (91%)	306 (98%)	5 (2%)	0	100	100
1	D	309/343 (90%)	301 (97%)	8 (3%)	0	100	100
All	All	1245/1372 (91%)	1215 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/305 (93%)	282 (100%)	1 (0%)	91	93
1	B	283/305 (93%)	279 (99%)	4 (1%)	67	70
1	C	282/305 (92%)	277 (98%)	5 (2%)	59	61
1	D	280/305 (92%)	278 (99%)	2 (1%)	84	87
All	All	1128/1220 (92%)	1116 (99%)	12 (1%)	73	77

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	681	LEU
1	B	617	SER
1	B	638	TYR
1	B	660	SER
1	B	749	GLU
1	C	504	GLU
1	C	616	SER
1	C	617	SER
1	C	638	TYR
1	C	749	GLU
1	D	617	SER
1	D	651	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	461	GLN
1	A	484	ASN
1	A	542	GLN
1	A	604	GLN
1	A	743	GLN
1	A	761	GLN
1	B	484	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	542	GLN
1	B	604	GLN
1	C	542	GLN
1	C	604	GLN
1	C	743	GLN
1	D	461	GLN
1	D	484	ASN
1	D	542	GLN
1	D	621	GLN
1	D	731	ASN
1	D	743	GLN
1	D	761	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	C	509	1	8,9,10	0.55	0	5,9,11	0.73	0
1	CME	B	509	1	8,9,10	0.57	0	5,9,11	0.65	0
1	CME	D	509	1	8,9,10	0.67	0	5,9,11	0.40	0
1	CME	A	509	1	8,9,10	0.69	0	5,9,11	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	C	509	1	-	2/5/8/10	-
1	CME	B	509	1	-	0/5/8/10	-
1	CME	D	509	1	-	1/5/8/10	-
1	CME	A	509	1	-	2/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	509	CME	CE-SD-SG-CB
1	A	509	CME	SD-CE-CZ-OH
1	A	509	CME	CZ-CE-SD-SG
1	C	509	CME	CZ-CE-SD-SG
1	D	509	CME	CZ-CE-SD-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	509	CME	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	IEX	A	803	-	32,34,34	1.03	3 (9%)	37,47,47	0.68	1 (2%)
4	IEX	C	803	-	32,34,34	0.81	1 (3%)	37,47,47	0.64	0
4	IEX	B	803	-	32,34,34	0.60	1 (3%)	37,47,47	0.57	0
4	IEX	D	803	-	32,34,34	0.50	0	37,47,47	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IEX	A	803	-	-	1/13/13/13	0/5/5/5
4	IEX	C	803	-	-	1/13/13/13	0/5/5/5
4	IEX	B	803	-	-	1/13/13/13	0/5/5/5
4	IEX	D	803	-	-	0/13/13/13	0/5/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	803	IEX	C16-N15	-3.29	1.35	1.39
4	A	803	IEX	C16-N15	-2.38	1.36	1.39
4	A	803	IEX	C14-N15	2.31	1.40	1.37
4	A	803	IEX	C19-C17	-2.12	1.45	1.48
4	B	803	IEX	C16-N15	-2.03	1.37	1.39

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	803	IEX	C17-C16-N15	-2.44	105.38	107.88

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	803	IEX	C12-C13-C14-N15
4	C	803	IEX	C12-C13-C14-N15
4	A	803	IEX	C12-C13-C14-N15



There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	803	IEX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/343 (90%)	-0.29	1 (0%) 94 93	28, 40, 67, 98	0
1	B	314/343 (91%)	-0.24	1 (0%) 94 93	27, 38, 65, 106	0
1	C	312/343 (90%)	-0.50	2 (0%) 89 89	28, 40, 62, 92	0
1	D	311/343 (90%)	-0.19	6 (1%) 66 66	35, 50, 73, 93	0
All	All	1249/1372 (91%)	-0.30	10 (0%) 86 85	27, 42, 69, 106	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	458	GLY	4.0
1	C	770	GLU	3.1
1	B	771	THR	2.6
1	D	654	LEU	2.5
1	C	768	GLY	2.3
1	D	603	LEU	2.3
1	D	457	GLN	2.2
1	D	656	LEU	2.2
1	D	657	ASN	2.1
1	D	768	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	C	509	10/11	0.91	0.11	42,47,80,83	0
1	CME	B	509	10/11	0.92	0.14	42,53,91,98	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	509	10/11	0.92	0.12	42,57,90,90	0
1	CME	D	509	10/11	0.93	0.14	46,60,104,106	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	D	802	1/1	0.91	0.03	38,38,38,38	0
4	IEX	D	803	30/30	0.94	0.10	36,46,52,53	0
4	IEX	B	803	30/30	0.95	0.10	31,34,38,39	0
4	IEX	C	803	30/30	0.97	0.10	27,37,43,44	0
4	IEX	A	803	30/30	0.97	0.10	27,36,43,44	0
3	MG	A	802	1/1	0.98	0.05	29,29,29,29	0
3	MG	C	802	1/1	0.99	0.09	29,29,29,29	0
2	ZN	D	801	1/1	0.99	0.06	42,42,42,42	0
2	ZN	B	801	1/1	1.00	0.10	32,32,32,32	0
3	MG	B	802	1/1	1.00	0.11	27,27,27,27	0
2	ZN	C	801	1/1	1.00	0.07	34,34,34,34	0
2	ZN	A	801	1/1	1.00	0.10	35,35,35,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around IEX D 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IEX B 803:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IEX C 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around IEX A 803:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.