



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 27, 2024 – 02:35 PM EST

PDB ID : 8SNF  
Title : Crystal structure of metformin hydrolase (MfmAB) from *Pseudomonas mendocina* sp. MET-2 with Ni<sup>2+</sup> bound  
Authors : Tassoulas, L.J.; Rankin, J.A.; Elias, M.H.; Wackett, L.P.  
Deposited on : 2023-04-27  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	 3% 82% 9% • 6%
1	B	348	 5% 81% 11% • 6%
1	C	348	 5% 81% 11% • 6%
1	E	348	 3% 82% 11% • 6%
2	D	364	 2% 80% 12% •• 6%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	364	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '82%', a yellow segment labeled '12%', and a small grey segment at the end labeled '5%'.</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 15792 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called metformin hydrolase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	326	2507	1579	439	470	19	0	0	0
1	B	326	2507	1579	439	470	19	0	0	0
1	C	326	2507	1579	439	470	19	0	0	0
1	E	326	2507	1579	439	470	19	0	0	0

- Molecule 2 is a protein called metformin hydrolase subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	347	2715	1711	476	505	23	0	0	0
2	F	347	2715	1711	476	505	23	0	0	0

- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	2	Total	Ni	0	0
			2	2		
3	F	2	Total	Ni	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	55	Total	O	0	0
			55	55		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	B	27	Total O 27 27	0	0
4	C	24	Total O 24 24	0	0
4	D	91	Total O 91 91	0	0
4	E	57	Total O 57 57	0	0
4	F	76	Total O 76 76	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: metformin hydrolase subunit B




- Molecule 1: metformin hydrolase subunit B




- Molecule 1: metformin hydrolase subunit B





- Molecule 1: metformin hydrolase subunit B







• Molecule 2: metformin hydrolase subunit A



• Molecule 2: metformin hydrolase subunit A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.40Å 96.40Å 96.70Å 115.50° 106.20° 101.10°	Depositor
Resolution (Å)	19.86 – 2.30 19.86 – 2.30	Depositor EDS
% Data completeness (in resolution range)	91.6 (19.86-2.30) 91.8 (19.86-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.99 (at 2.30Å)	Xtrriage
Refinement program	REFMAC 5.8.0411	Depositor
R, $R_{free}$	0.172 , 0.219 0.171 , 0.212	Depositor DCC
$R_{free}$ test set	5056 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.3	Xtrriage
Anisotropy	0.122	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 58.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	15792	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2558	0.80	4/3459 (0.1%)
1	B	0.36	0/2558	0.66	0/3459
1	C	0.37	0/2558	0.69	0/3459
1	E	0.43	0/2558	0.76	0/3459
2	D	0.40	0/2787	0.79	3/3777 (0.1%)
2	F	0.40	0/2787	0.73	0/3777
All	All	0.40	0/15806	0.74	7/21390 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	E	0	3
2	D	0	2
2	F	0	4
All	All	0	14

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	283	ARG	NE-CZ-NH1	8.47	124.54	120.30
2	D	158	ASP	CB-CA-C	-7.85	94.70	110.40
1	A	283	ARG	NE-CZ-NH2	-7.25	116.68	120.30
1	A	271	THR	CB-CA-C	-5.69	96.24	111.60
2	D	318	ASP	CB-CA-C	5.49	121.37	110.40

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	115	ARG	Sidechain
1	A	137	ARG	Sidechain
1	A	243	ARG	Sidechain
1	A	291	ARG	Sidechain
1	A	32	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2507	0	2475	22	0
1	B	2507	0	2475	31	0
1	C	2507	0	2475	28	0
1	E	2507	0	2475	24	0
2	D	2715	0	2600	35	0
2	F	2715	0	2600	31	0
3	D	2	0	0	0	0
3	F	2	0	0	0	0
4	A	55	0	0	0	0
4	B	27	0	0	0	0
4	C	24	0	0	1	0
4	D	91	0	0	1	0
4	E	57	0	0	1	0
4	F	76	0	0	0	0
All	All	15792	0	15100	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:206:MET:HE1	1:E:81:TYR:HA	1.50	0.92
2:D:94:LEU:HD23	2:D:336:THR:CG2	2.04	0.88

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:94:LEU:HD23	2:F:336:THR:HG23	1.59	0.84
1:B:113:HIS:ND1	1:B:142:THR:HG21	1.95	0.81
2:D:94:LEU:HD23	2:D:336:THR:HG23	1.64	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/348 (92%)	311 (97%)	10 (3%)	1 (0%)	41	50
1	B	322/348 (92%)	311 (97%)	10 (3%)	1 (0%)	41	50
1	C	322/348 (92%)	307 (95%)	12 (4%)	3 (1%)	17	20
1	E	322/348 (92%)	311 (97%)	10 (3%)	1 (0%)	41	50
2	D	345/364 (95%)	323 (94%)	21 (6%)	1 (0%)	41	50
2	F	345/364 (95%)	323 (94%)	22 (6%)	0	100	100
All	All	1978/2120 (93%)	1886 (95%)	85 (4%)	7 (0%)	34	42

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	103	SER
1	B	107	VAL
1	C	107	VAL
1	E	233	ILE
2	D	83	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/279 (95%)	256 (97%)	9 (3%)	37	51
1	B	265/279 (95%)	255 (96%)	10 (4%)	33	47
1	C	265/279 (95%)	249 (94%)	16 (6%)	19	26
1	E	265/279 (95%)	257 (97%)	8 (3%)	41	57
2	D	287/302 (95%)	275 (96%)	12 (4%)	30	42
2	F	287/302 (95%)	278 (97%)	9 (3%)	40	55
All	All	1634/1720 (95%)	1570 (96%)	64 (4%)	32	46

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	118	VAL
2	F	224	ARG
1	C	136	ASP
1	C	118	ARG
2	F	255	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	13	GLN
2	D	304	GLN
2	F	357	GLN
1	E	79	GLN
1	C	165	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/348 (93%)	-0.21	10 (3%) 49 56	54, 72, 103, 150	0
1	B	326/348 (93%)	0.05	17 (5%) 27 34	62, 82, 115, 181	0
1	C	326/348 (93%)	0.04	18 (5%) 25 31	63, 84, 117, 174	0
1	E	326/348 (93%)	-0.29	9 (2%) 53 60	53, 70, 101, 139	0
2	D	347/364 (95%)	-0.22	8 (2%) 60 67	54, 70, 95, 149	0
2	F	347/364 (95%)	-0.20	9 (2%) 56 63	55, 72, 100, 140	0
All	All	1998/2120 (94%)	-0.14	71 (3%) 42 49	53, 75, 108, 181	0

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	106	LYS	6.0
2	D	356	LYS	5.8
1	C	7	TYR	5.6
2	F	355	GLY	5.2
1	C	306	PHE	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NI	D	401	1/1	0.96	0.03	112,112,112,112	0
3	NI	F	402	1/1	0.96	0.05	125,125,125,125	0
3	NI	F	401	1/1	0.98	0.02	95,95,95,95	0
3	NI	D	402	1/1	0.98	0.04	87,87,87,87	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around NI F 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.