

wwPDB X-ray Structure Validation Summary Report (i)

Aug 15, 2023 – 11:38 PM EDT

PDB ID	:	1T36
Title	:	Crystal structure of E. coli carbamoyl phosphate synthetase small subunit
		mutant C248D complexed with uridine 5'-monophosphate
Authors	:	Thoden, J.B.; Huang, X.; Raushel, F.M.; Holden, H.M.
Deposited on	:	2004-04-24
Resolution	:	2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
	$(\# {\rm Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of ch	ain	
1	Δ	1073	2%	200/	70/
1		1075	62%	50%	/% •
1	С	1073	58%	32%	7% ••
1	Е	1073	65%	27%	6% ••
1	C	1079	4%		_
	G	1073	52% 4%	37%	9% ••
2	В	382	54%	35%	8% ••



Continued from previous page...

Mol	Chain	Length	Quality of chain					
2	D	382	3% 52%	38%	8% ••			
2	F	382	<u>53%</u>	37%	8% ••			
2	Н	382	9%	46%	11% ••			

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PO4	Е	1078	-	Х	-	-
5	PO4	G	1078	-	Х	-	-



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 48757 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	1058	Total	С	Ν	Ο	\mathbf{S}	0	Q	0
1	Π	1050	8212	5155	1436	1575	46	0	3	0
1	С	1058	Total	С	Ν	Ο	\mathbf{S}	0	8	0
1	U	1058	8197	5146	1426	1579	46	0	0	
1	F	1058	Total	С	Ν	Ο	\mathbf{S}	0	Б	0
1		1058	8182	5137	1425	1575	45	0	5	0
1	1 C	1058	Total	С	Ν	Ο	S	0	8	0
I G	1058	8206	5152	1432	1577	45		8	U	

• Molecule 1 is a protein called Carbamoyl-phosphate synthase large chain.

• Molecule 2 is a protein called Carbamoyl-phosphate synthase small chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
0	р	270	Total	С	Ν	0	\mathbf{S}	0	0	0
	D	579	2897	1826	508	554	9	0	0	U
0	П	270	Total	С	Ν	0	S	0	1	0
	D	519	2904	1830	511	554	9	0	L	0
0	Б	270	Total	С	Ν	0	S	0	0	0
	Г	579	2897	1826	508	554	9	0	0	0
0	о п	II 970	Total	С	Ν	0	S	0	0	0
	379	2897	1826	508	554	9		U	0	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	248	ASP	CYS	engineered mutation	UNP P00907
D	248	ASP	CYS	engineered mutation	UNP P00907
F	248	ASP	CYS	engineered mutation	UNP P00907
Н	248	ASP	CYS	engineered mutation	UNP P00907

• Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	3	Total Mn 3 3	0	0
3	С	3	Total Mn 3 3	0	0
3	Ε	3	Total Mn 3 3	0	0
3	G	3	Total Mn 3 3	0	0

• Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	4	Total K 4 4	0	0
4	В	1	Total K 1 1	0	0
4	С	4	Total K 4 4	0	0
4	D	1	Total K 1 1	0	0
4	Ε	5	Total K 5 5	0	0
4	F	1	Total K 1 1	0	0
4	G	5	Total K 5 5	0	0
4	Н	1	Total K 1 1	0	0

• Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
5	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
5	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
5	Е	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
5	G	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	5	Total Cl 5 5	0	0
6	С	6	Total Cl 6 6	0	0
6	D	1	Total Cl 1 1	0	0
6	Е	6	Total Cl 6 6	0	0
6	F	1	Total Cl 1 1	0	0
6	G	6	Total Cl 6 6	0	0
6	Н	2	Total Cl 2 2	0	0



• Molecule 7 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
7	٨	1	Total	С	Ν	Ο	Р	0	0		
(A	1	27	10	5	10	2	0	0		
7	Λ	1	Total	С	Ν	Ο	Р	0	0		
(A	1	27	10	5	10	2	0	0		
7	С	1	Total	С	Ν	Ο	Р	0	0		
1	U	1	27	10	5	10	2	0	0		
7	С	1	Total	С	Ν	Ο	Р	0	0		
1	U	U		T	27	10	5	10	2	0	0
7	F	1	Total	С	Ν	Ο	Р	0	0		
1	Ľ	Ľ	Ľ	1	27	10	5	10	2	0	0
7	F	1	Total	С	Ν	Ο	Р	0	0		
1	Ľ	1	27	10	5	10	2	0	0		
7	С	1	Total	С	Ν	Ο	Р	0	0		
1	G		27	10	5	10	2	0	0		
7	С	1	Total	С	Ν	Ο	Р	0	0		
1	G		27	10	5	10	2	0	0		

• Molecule 8 is L-ornithine (three-letter code: ORN) (formula: $C_5H_{12}N_2O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	1	Total C N O	0	0
8	С	1	9322TotalCNO	0	0
		1	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		0
8	Ε	1	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	0	0
8	G	1	$\begin{array}{cccc} \text{Total} \text{C} \text{N} \text{O} \\ 9 5 2 2 \end{array}$	0	0

 $\bullet\,$ Molecule 9 is TETRAETHYLAMMONIUM ION (three-letter code: NET) (formula: C_8H_{20}N).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	1	Total C N 9 8 1	0	0
9	С	1	Total C N 9 8 1	0	0
9	Е	1	Total C N 9 8 1	0	0
9	G	1	$\begin{array}{c cc} Total & C & N \\ 9 & 8 & 1 \end{array}$	0	0

• Molecule 10 is URIDINE-5'-MONOPHOSPHATE (three-letter code: U) (formula: $C_9H_{13}N_2O_9P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
10	Δ	1	Total	С	Ν	0	Р	0	0
10	A	L	21	9	2	9	1	0	0
10	C	1	Total	С	Ν	0	Р	0	0
10	U	L	21	9	2	9	1	0	0
10	F	1	Total	С	Ν	0	Р	0	0
10	Ľ	L	21	9	2	9	1	0	0
10	С	1	Total	С	Ν	Ο	Р	0	0
10	G	L	21	9	2	9	1	0	0

• Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	А	851	Total O 851 851	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	В	161	Total O 161 161	0	0
11	С	819	Total O 819 819	0	0
11	D	221	Total O 221 221	0	0
11	Е	832	Total O 832 832	0	0
11	F	200	Total O 200 200	0	0
11	G	705	Total O 705 705	0	0
11	Н	118	Total O 118 118	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Carbamoyl-phosphate synthase large chain



• Molecule 1: Carbamoyl-phosphate synthase large chain



• Molecule 1: Carbamoyl-phosphate synthase large chain















• Molecule 2: Carbamoyl-phosphate synthase small chain



• Molecule 2: Carbamoyl-phosphate synthase small chain









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	152.50Å 164.90Å 333.10Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(A)	30.00 - 2.10	Depositor
Resolution (A)	29.70 - 2.10	EDS
% Data completeness	90.0 (30.00-2.10)	Depositor
(in resolution range)	88.6 (29.70-2.10)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$< I/\sigma(I) > 1$	$0.88 (at 2.10 \text{\AA})$	Xtriage
Refinement program	TNT	Depositor
B B.	0.176 , 0.209	Depositor
II, II free	0.175 , 0.176	DCC
R_{free} test set	42730 reflections $(9.94%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	28.2	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.29, 120.0	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	48757	wwPDB-VP
Average B, all atoms $(Å^2)$	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, ADP, PO4, ORN, NET, K, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	B	ond lengths	Bond angles		
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	1.05	68/8374~(0.8%)	1.50	137/11315~(1.2%)	
1	С	1.06	81/8355~(1.0%)	1.48	127/11293~(1.1%)	
1	Е	1.06	74/8328~(0.9%)	1.50	128/11257~(1.1%)	
1	G	1.05	74/8368~(0.9%)	1.50	148/11308~(1.3%)	
2	В	0.94	19/2959~(0.6%)	1.46	43/4019~(1.1%)	
2	D	0.97	17/2970~(0.6%)	1.48	45/4033~(1.1%)	
2	F	0.97	19/2959~(0.6%)	1.45	41/4019~(1.0%)	
2	Н	0.94	20/2959~(0.7%)	1.42	37/4019~(0.9%)	
All	All	1.03	372/45272~(0.8%)	1.49	706/61263~(1.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	1	0
2	D	1	0
2	F	1	0
All	All	3	0

The worst 5 of 372 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	372	GLU	CD-OE2	8.39	1.34	1.25
2	Н	166	GLU	CD-OE2	8.34	1.34	1.25
2	D	145	GLU	CD-OE2	8.28	1.34	1.25
1	G	1009[A]	GLU	CD-OE2	8.21	1.34	1.25
1	G	1009[B]	GLU	CD-OE2	8.21	1.34	1.25

The worst 5 of 706 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	Е	514	ARG	NE-CZ-NH2	-15.12	112.74	120.30
1	А	944	ARG	NE-CZ-NH1	14.25	127.42	120.30
1	С	514	ARG	NE-CZ-NH2	-14.00	113.30	120.30
1	А	104	ARG	NE-CZ-NH1	13.43	127.02	120.30
1	А	652	ARG	NE-CZ-NH1	13.13	126.87	120.30

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	В	154	ASN	CA
2	D	154	ASN	CA
2	F	154	ASN	CA

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	8212	0	8255	231	0
1	С	8197	0	8225	254	0
1	Е	8182	0	8216	209	0
1	G	8206	0	8247	340	0
2	В	2897	0	2860	147	0
2	D	2904	0	2869	133	0
2	F	2897	0	2860	128	0
2	Н	2897	0	2860	213	0
3	А	3	0	0	0	0
3	С	3	0	0	0	0
3	Е	3	0	0	0	0
3	G	3	0	0	0	0
4	А	4	0	0	0	0
4	В	1	0	0	0	0
4	С	4	0	0	0	0
4	D	1	0	0	0	0
4	Е	5	0	0	0	0
4	F	1	0	0	0	0
4	G	5	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Н	1	0	0	0	0
5	А	5	0	0	0	0
5	С	10	0	0	1	0
5	Е	5	0	0	0	0
5	G	5	0	0	0	0
6	А	5	0	0	1	0
6	С	6	0	0	0	0
6	D	1	0	0	0	0
6	Е	6	0	0	1	0
6	F	1	0	0	0	0
6	G	6	0	0	2	0
6	Н	2	0	0	0	0
7	А	54	0	24	2	0
7	С	54	0	24	0	0
7	Е	54	0	24	3	0
7	G	54	0	24	0	0
8	А	9	0	11	0	0
8	С	9	0	11	1	0
8	Е	9	0	11	2	0
8	G	9	0	11	1	0
9	А	9	0	20	1	0
9	С	9	0	20	1	0
9	Е	9	0	20	2	0
9	G	9	0	20	2	0
10	А	21	0	11	1	0
10	С	21	0	11	2	0
10	Е	21	0	11	2	0
10	G	21	0	11	1	0
11	А	851	0	0	29	1
11	В	161	0	0	5	0
11	С	819	0	0	23	1
11	D	221	0	0	5	0
11	Е	832	0	0	24	0
11	F	200	0	0	2	0
11	G	705	0	0	28	0
11	Н	118	0	0	5	0
All	All	48757	0	44656	1643	1

Continued from previous nage

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 1643 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:27:VAL:HG22	2:H:131:CYS:HB2	1.13	1.10
2:D:227:ASP:HA	2:D:230:LYS:HD2	1.25	1.07
1:G:784:GLN:NE2	1:G:784:GLN:H	1.53	1.05
2:H:133:ILE:HD12	2:H:143:ALA:HB2	1.41	1.03
1:A:38:ARG:HG3	1:A:38:ARG:HH11	1.25	1.00

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:A:1848:HOH:O	11:C:1912:HOH:O[4_555]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	1061/1073~(99%)	1015 (96%)	44 (4%)	2~(0%)	47	49
1	С	1060/1073~(99%)	1005 (95%)	54 (5%)	1 (0%)	51	54
1	Е	1057/1073~(98%)	1013 (96%)	42 (4%)	2~(0%)	47	49
1	G	1061/1073~(99%)	996 (94%)	59~(6%)	6 (1%)	25	21
2	В	377/382~(99%)	348 (92%)	26 (7%)	3 (1%)	19	15
2	D	378/382~(99%)	354 (94%)	23~(6%)	1 (0%)	41	41
2	F	377/382~(99%)	355 (94%)	21 (6%)	1 (0%)	41	41
2	Н	377/382~(99%)	346 (92%)	27 (7%)	4 (1%)	14	9
All	All	5748/5820~(99%)	5432 (94%)	296 (5%)	20 (0%)	41	41

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
2	В	154	ASN	
Continued on next nage				



Continued from previous page...

Mol	Chain	Res	Type
2	D	154	ASN
1	Е	738	PHE
1	G	485	ASN
1	G	975	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	874/878~(100%)	811 (93%)	63~(7%)	14 11
1	С	873/878~(99%)	809 (93%)	64 (7%)	14 11
1	Ε	870/878~(99%)	818 (94%)	52~(6%)	19 16
1	G	874/878~(100%)	800~(92%)	74 (8%)	10 7
2	В	308/310~(99%)	283~(92%)	25~(8%)	11 8
2	D	309/310~(100%)	283~(92%)	26~(8%)	11 7
2	F	308/310~(99%)	276~(90%)	32 (10%)	7 4
2	Н	308/310~(99%)	278 (90%)	30 (10%)	8 5
All	All	4724/4752~(99%)	4358 (92%)	366 (8%)	13 9

 $5~{\rm of}~366$ residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
2	F	4	SER
1	G	509	ARG
2	F	166	GLU
2	F	376	GLN
1	G	696	THR

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 78 such side chains are listed below:

Mol	Chain	Res	Type	
1	G	266	ASN	
Continued on next page				



Continued from previous page...

Mol	Chain	Res	Type
1	G	1071	GLN
1	G	679	GLN
1	G	936	ASN
2	Н	291	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 86 ligands modelled in this entry, 61 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Tink	Bo	Bond lengths			Bond angles		
	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	PO4	C	1088	-	4,4,4	2.51	3 (75%)	6,6,6	0.97	0	
7	ADP	C	1090	3	24,29,29	1.01	1 (4%)	29,45,45	1.17	4 (13%)	
7	ADP	А	1087	3	24,29,29	1.14	2 (8%)	29,45,45	1.05	3 (10%)	
7	ADP	А	1088	3	24,29,29	0.95	1 (4%)	29,45,45	1.23	2 (6%)	
7	ADP	G	1090	3	24,29,29	1.22	3 (12%)	29,45,45	1.15	4 (13%)	
9	NET	А	1090	-	8,8,8	0.72	0	10,10,10	0.65	0	
7	ADP	E	1090	3	24,29,29	1.21	3 (12%)	29,45,45	1.10	2 (6%)	
10	U	G	1093	-	22,22,22	0.94	1 (4%)	33,33,33	1.44	3 (9%)	



17	$\Gamma 36$
	- 00

Mal	Tuno	Chain	Dog	Link	Bo	ond leng	$_{\rm sths}$	Bond angles		
INIOI	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	U	А	1091	-	22,22,22	0.84	1 (4%)	33,33,33	0.81	0
10	U	С	1093	-	22,22,22	0.82	1 (4%)	33,33,33	0.90	0
7	ADP	G	1089	3	24,29,29	1.27	3 (12%)	29,45,45	1.21	4 (13%)
7	ADP	С	1089	3	24,29,29	1.21	2 (8%)	29,45,45	1.39	4 (13%)
5	PO4	Е	1078	3	4,4,4	2.51	3 (75%)	6,6,6	1.33	1 (16%)
8	ORN	А	1089	-	7,8,8	0.89	0	8,9,9	1.91	2 (25%)
8	ORN	С	1091	-	7,8,8	1.17	1 (14%)	8,9,9	1.21	1 (12%)
8	ORN	G	1091	-	7,8,8	0.84	0	8,9,9	1.26	1 (12%)
10	U	Е	1093	-	22,22,22	0.96	1 (4%)	33,33,33	1.60	5 (15%)
5	PO4	А	1078	3	4,4,4	1.96	1 (25%)	6,6,6	1.37	1 (16%)
7	ADP	Е	1089	3	24,29,29	1.16	2 (8%)	29,45,45	1.09	3 (10%)
9	NET	Е	1092	-	8,8,8	0.39	0	10,10,10	0.75	0
5	PO4	С	1078	3	4,4,4	2.29	3 (75%)	6,6,6	1.05	0
5	PO4	G	1078	3	4,4,4	2.31	3 (75%)	6,6,6	1.20	1 (16%)
9	NET	С	1092	-	8,8,8	0.49	0	10,10,10	0.76	0
9	NET	G	1092	-	8,8,8	0.71	0	10,10,10	0.46	0
8	ORN	Е	1091	-	7,8,8	1.07	1 (14%)	8,9,9	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
7	ADP	С	1090	3	-	3/12/32/32	0/3/3/3
7	ADP	А	1087	3	-	1/12/32/32	0/3/3/3
7	ADP	А	1088	3	-	4/12/32/32	0/3/3/3
7	ADP	G	1090	3	-	1/12/32/32	0/3/3/3
9	NET	А	1090	-	-	3/12/12/12	-
7	ADP	Е	1090	3	-	3/12/32/32	0/3/3/3
10	U	G	1093	-	-	6/10/26/26	0/2/2/2
10	U	А	1091	-	-	6/10/26/26	0/2/2/2
10	U	С	1093	-	-	5/10/26/26	0/2/2/2
7	ADP	G	1089	3	-	0/12/32/32	0/3/3/3
7	ADP	С	1089	3	-	2/12/32/32	0/3/3/3
8	ORN	А	1089	-	-	6/8/8/8	-
8	ORN	С	1091	-	-	6/8/8/8	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	ORN	G	1091	-	-	6/8/8/8	-
10	U	Е	1093	-	-	6/10/26/26	0/2/2/2
7	ADP	Е	1089	3	-	1/12/32/32	0/3/3/3
9	NET	Е	1092	-	-	0/12/12/12	-
9	NET	С	1092	-	-	0/12/12/12	-
9	NET	G	1092	-	-	0/12/12/12	-
8	ORN	Е	1091	-	-	5/8/8/8	-

Continued from previous page...

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
5	С	1088	PO4	P-O2	-3.46	1.44	1.54
5	Е	1078	PO4	P-O3	-3.27	1.44	1.54
7	Е	1090	ADP	O2'-C2'	3.27	1.50	1.43
7	С	1089	ADP	O2'-C2'	3.13	1.50	1.43
7	Е	1090	ADP	O3'-C3'	3.11	1.50	1.43

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
7	С	1089	ADP	C5-C6-N6	5.14	128.16	120.35
10	Е	1093	U	OP3-P-O5'	-4.97	93.51	106.73
10	G	1093	U	OP3-P-O5'	-4.36	95.14	106.73
7	А	1088	ADP	C5-C6-N6	4.02	126.47	120.35
8	А	1089	ORN	CB-CA-C	-3.39	102.24	110.30

There are no chirality outliers.

5 of 64 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	С	1090	ADP	PA-O3A-PB-O3B
7	Е	1090	ADP	PA-O3A-PB-O3B
8	А	1089	ORN	N-CA-CB-CG
8	А	1089	ORN	C-CA-CB-CG
8	С	1091	ORN	N-CA-CB-CG

There are no ring outliers.

16 monomers are involved in 22 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	А	1087	ADP	1	0
7	А	1088	ADP	1	0
9	А	1090	NET	1	0
7	Е	1090	ADP	2	0
10	G	1093	U	1	0
10	А	1091	U	1	0
10	С	1093	U	2	0
8	С	1091	ORN	1	0
8	G	1091	ORN	1	0
10	Е	1093	U	2	0
7	Е	1089	ADP	1	0
9	Е	1092	NET	2	0
5	С	1078	PO4	1	0
9	С	1092	NET	1	0
9	G	1092	NET	2	0
8	Е	1091	ORN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	А	1058/1073~(98%)	-0.41	23 (2%) 62 66	17, 33, 73, 100	0
1	С	1058/1073~(98%)	-0.28	25 (2%) 59 64	18, 35, 81, 100	0
1	Ε	1058/1073~(98%)	-0.35	20 (1%) 66 71	17, 31, 75, 100	0
1	G	1058/1073~(98%)	-0.11	40 (3%) 40 46	20, 42, 84, 100	0
2	В	379/382~(99%)	0.02	16 (4%) 36 42	21, 48, 87, 100	0
2	D	379/382~(99%)	-0.04	11 (2%) 51 57	21, 40, 78, 100	0
2	F	379/382~(99%)	0.17	24 (6%) 20 24	20, 47, 92, 100	0
2	Н	379/382~(99%)	0.47	36 (9%) 8 10	33, 63, 97, 100	0
All	All	5748/5820 (98%)	-0.17	195 (3%) 45 51	17, 38, 84, 100	0

The worst 5 of 195 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	1	MET	8.3
1	Е	1	MET	8.0
1	А	1	MET	7.4
1	G	1	MET	6.9
1	С	697	ALA	5.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
6	CL	Е	1084	1/1	0.82	0.15	74,74,74,74	0
6	CL	G	1084	1/1	0.83	0.22	72,72,72,72	0
6	CL	Н	385	1/1	0.84	0.07	82,82,82,82	0
6	CL	С	1083	1/1	0.91	0.16	77,77,77,77	0
6	CL	А	1083	1/1	0.91	0.10	66,66,66,66	0
6	CL	G	1086	1/1	0.92	0.07	54,54,54,54	0
5	PO4	С	1088	5/5	0.92	0.21	80,80,80,80	0
6	CL	Е	1087	1/1	0.94	0.06	54,54,54,54	0
10	U	G	1093	21/21	0.94	0.12	31,57,80,85	0
6	CL	Н	384	1/1	0.95	0.13	67,67,67,67	0
4	К	Е	1081	1/1	0.95	0.06	42,42,42,42	0
8	ORN	А	1089	9/9	0.95	0.18	24,29,36,37	0
10	U	С	1093	21/21	0.95	0.12	33,51,83,89	0
6	CL	С	1086	1/1	0.95	0.11	67,67,67,67	0
4	K	С	1082	1/1	0.96	0.06	44,44,44,44	0
8	ORN	G	1091	9/9	0.96	0.17	28,33,36,63	0
10	U	А	1091	21/21	0.96	0.10	28,58,80,81	0
6	CL	С	1084	1/1	0.96	0.08	43,43,43,43	0
10	U	Е	1093	21/21	0.96	0.10	37,54,80,89	0
6	CL	А	1086	1/1	0.96	0.06	56, 56, 56, 56	0
4	K	G	1081	1/1	0.97	0.06	46,46,46,46	0
4	K	Н	383	1/1	0.97	0.11	64,64,64,64	0
6	CL	G	1088	1/1	0.97	0.17	67,67,67,67	0
8	ORN	С	1091	9/9	0.97	0.19	20,29,35,39	0
8	ORN	Е	1091	9/9	0.97	0.13	16,25,37,40	0
6	CL	G	1085	1/1	0.98	0.06	46,46,46,46	0
4	K	Е	1083	1/1	0.98	0.05	58,58,58,58	0
6	CL	G	1087	1/1	0.98	0.09	81,81,81,81	0
6	CL	Е	1085	1/1	0.98	0.05	49,49,49,49	0
9	NET	А	1090	9/9	0.98	0.10	18,23,28,38	0
9	NET	Е	1092	9/9	0.98	0.18	15,23,27,28	0
4	K	А	1082	1/1	0.98	0.04	41,41,41,41	0
4	K	В	383	1/1	0.98	0.06	45,45,45,45	0
7	ADP	С	1090	27/27	0.98	0.08	26,42,61,77	0
7	ADP	G	1090	27/27	0.98	0.09	29,47,80,89	0
4	K	Е	1077	1/1	0.99	0.12	30,30,30,30	0
6	CL	С	1087	1/1	0.99	0.15	63,63,63,63	0



1	T36	
т	100	

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
6	CL	D	384	1/1	0.99	0.07	34,34,34,34	0
4	K	Е	1080	1/1	0.99	0.06	27,27,27,27	0
4	K	А	1077	1/1	0.99	0.08	27,27,27,27	0
6	CL	Е	1086	1/1	0.99	0.07	41,41,41,41	0
4	Κ	А	1080	1/1	0.99	0.08	29,29,29,29	0
6	CL	Е	1088	1/1	0.99	0.10	44,44,44,44	0
6	CL	G	1082	1/1	0.99	0.10	29,29,29,29	0
4	Κ	F	383	1/1	0.99	0.07	36,36,36,36	0
4	Κ	G	1077	1/1	0.99	0.10	38,38,38,38	0
4	Κ	G	1080	1/1	0.99	0.07	38,38,38,38	0
3	MN	С	1075	1/1	0.99	0.07	25,25,25,25	0
4	Κ	G	1083	1/1	0.99	0.05	52,52,52,52	0
3	MN	С	1079	1/1	0.99	0.05	46,46,46,46	0
5	PO4	А	1078	5/5	0.99	0.08	17,21,24,29	0
7	ADP	А	1087	27/27	0.99	0.09	16,22,31,38	0
7	ADP	А	1088	27/27	0.99	0.07	20,33,47,60	0
7	ADP	С	1089	27/27	0.99	0.10	16,23,31,44	0
5	PO4	С	1078	5/5	0.99	0.09	16,22,24,25	0
7	ADP	Е	1089	27/27	0.99	0.12	15,26,34,36	0
7	ADP	Е	1090	27/27	0.99	0.06	23,35,48,56	0
7	ADP	G	1089	27/27	0.99	0.11	17,27,35,41	0
4	Κ	С	1077	1/1	0.99	0.11	27,27,27,27	0
5	PO4	Е	1078	5/5	0.99	0.08	19,22,26,27	0
5	PO4	G	1078	5/5	0.99	0.07	17,24,31,35	0
6	CL	А	1081	1/1	0.99	0.08	28,28,28,28	0
4	Κ	С	1080	1/1	0.99	0.08	36,36,36,36	0
6	CL	А	1084	1/1	0.99	0.05	40,40,40,40	0
9	NET	С	1092	9/9	0.99	0.13	16,21,23,29	0
6	CL	А	1085	1/1	0.99	0.06	39,39,39,39	0
9	NET	G	1092	9/9	0.99	0.13	18,28,29,37	0
3	MN	G	1079	1/1	0.99	0.05	$47,\!47,\!47,\!47$	0
6	CL	С	1081	1/1	0.99	0.10	29,29,29,29	0
4	Κ	D	383	1/1	0.99	0.03	34,34,34,34	0
4	Κ	Е	1076	1/1	0.99	0.09	$25,\!25,\!25,\!25$	0
3	MN	Ε	1075	1/1	1.00	0.08	$27,\!27,\!27,\!27$	0
6	CL	F	384	1/1	1.00	0.06	32,32,32,32	0
3	MN	E	1079	1/1	1.00	0.05	41,41,41,41	0
3	MN	G	1074	1/1	1.00	0.07	31,31,31,31	0
3	MN	G	1075	1/1	1.00	0.08	28,28,28,28	0
3	MN	A	1079	1/1	1.00	0.05	$37,\!37,\!37,\!37$	0
6	CL	С	1085	1/1	1.00	0.07	39,39,39,39	0
4	K	A	1076	1/1	1.00	0.07	21,21,21,21	0

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	С	1074	1/1	1.00	0.08	27,27,27,27	0
3	MN	А	1074	1/1	1.00	0.07	25,25,25,25	0
6	CL	Е	1082	1/1	1.00	0.14	24,24,24,24	0
3	MN	А	1075	1/1	1.00	0.06	23,23,23,23	0
3	MN	Е	1074	1/1	1.00	0.09	27,27,27,27	0
4	K	G	1076	1/1	1.00	0.09	28,28,28,28	0
4	K	С	1076	1/1	1.00	0.07	22,22,22,22	0

Continued from previous page...

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

































6.5 Other polymers (i)

There are no such residues in this entry.

