

# Full wwPDB X-ray Structure Validation Report (i)

May 13, 2020 – 09:37 pm BST

PDB ID : 1T3G

Title: Crystal structure of the Toll/interleukin-1 receptor (TIR) domain of human

IL-1RAPL

Authors : Khan, J.A.; Tong, L.

Deposited on : 2004-04-26

Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{resolution range}( ext{Å}))$
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	159	72%	21%			
1	В	159	72%	20%	• 5%		



## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2725 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called X-linked interleukin-1 receptor accessory protein-like 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	152	Total 1271		N 212		Se 5	0	0	0
1	В	151	Total 1263		N 211		Se 5	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	437	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
A	477	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
A	499	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
A	519	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
A	559	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
В	437	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
В	477	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
В	499	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
В	519	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1
В	559	MSE	MET	MODIFIED RESIDUE	UNP Q9NZN1

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	96	Total O 96 96	0	0
2	В	95	Total O 95 95	0	0




# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: X-linked interleukin-1 receptor accessory protein-like 1



• Molecule 1: X-linked interleukin-1 receptor accessory protein-like 1





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	51.99Å 53.27Å 183.25Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.88 - 2.30	Depositor
% Data completeness	94.6 (28.88-2.30)	Depositor
(in resolution range)	34.0 (20.00 2.90)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
$R, R_{free}$	0.228 , $0.272$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2725	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP



## 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
WIGI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	0.33	0/1293	0.58	0/1734	
1	В	0.35	0/1285	0.60	1/1723 (0.1%)	
All	All	0.34	0/2578	0.59	$1/3457 \ (0.0\%)$	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	${f Res}$	Type	${f Atoms}$	${f Z}$	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
1	В	534	LEU	CA-CB-CG	5.88	128.82	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1271	0	1295	31	0
1	В	1263	0	1291	39	0
2	A	96	0	0	6	0
2	В	95	0	0	2	0
All	All	2725	0	2586	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \; (\mathring{\rm A})$	overlap (Å)
1:B:454:ILE:HD12	1:B:454:ILE:H	1.25	0.98
1:A:434:LEU:HB3	1:A:435:PRO:HD3	1.61	0.82
1:B:560:PRO:O	1:B:561:PHE:HB2	1.79	0.81
1:B:474:ILE:HD11	1:B:559:MSE:HE1	1.63	0.80
1:A:518:ILE:H	1:A:518:ILE:HD12	1.48	0.78
1:A:542:PRO:HA	1:A:545:ASN:ND2	1.99	0.77
1:A:426:GLU:HB2	2:A:150:HOH:O	1.88	0.72
1:B:474:ILE:CD1	1:B:559:MSE:HE1	2.22	0.70
1:A:547:LEU:C	1:A:548:ASN:HD22	1.99	0.66
1:B:454:ILE:O	1:B:456:THR:HG23	1.94	0.66
1:A:516:ARG:HA	1:A:516:ARG:HE	1.61	0.66
1:B:434:LEU:HB3	1:B:435:PRO:HD3	1.78	0.65
1:B:499:MSE:HA	1:B:504:GLU:HG3	1.77	0.65
1:B:454:ILE:HD12	1:B:454:ILE:N	2.05	0.65
1:A:515:LEU:HD21	1:A:521:TYR:HA	1.82	0.62
1:A:465:ARG:O	1:A:469:GLN:HG3	2.02	0.60
1:B:555:LEU:HD11	1:B:559:MSE:CE	2.32	0.60
1:A:516:ARG:NE	1:A:516:ARG:HA	2.16	0.59
1:A:516:ARG:NE	1:A:517:GLY:H	2.01	0.59
1:A:425:GLU:HG2	2:A:108:HOH:O	2.03	0.56
1:A:479:PRO:HG3	1:A:513:SER:HB3	1.88	0.56
1:B:526:ALA:O	1:B:530:THR:HG23	2.07	0.55
1:B:453:LEU:O	1:B:455:PRO:HD3	2.07	0.54
1:A:542:PRO:HA	1:A:545:ASN:HD21	1.71	0.54
1:A:513:SER:HA	2:A:62:HOH:O	2.09	0.52
1:A:518:ILE:HD12	1:A:518:ILE:N	2.20	0.52
1:A:454:ILE:HG22	2:A:14:HOH:O	2.10	0.51
1:A:491:GLU:O	1:A:497:ARG:HD2	2.10	0.51
1:A:493:GLU:OE1	1:A:495:ARG:HB2	2.11	0.51
1:B:518:ILE:N	1:B:518:ILE:HD12	2.25	0.51
1:A:484:ARG:O	1:B:485:ARG:HD2	2.09	0.51
1:A:515:LEU:CD2	1:A:521:TYR:HA	2.41	0.51
1:B:432:GLU:O	1:B:435:PRO:HD2	2.12	0.50
1:B:516:ARG:HA	2:B:88:HOH:O	2.11	0.50
1:B:555:LEU:HD11	1:B:559:MSE:HE3	1.93	0.50
1:B:454:ILE:H	1:B:454:ILE:CD1	2.03	0.50
1:A:496:LEU:HA	1:A:499:MSE:HE3	1.93	0.50
1:B:463:VAL:O	1:B:467:VAL:HG23	2.11	0.49
1:A:518:ILE:H	1:A:518:ILE:CD1	2.21	0.48
1:A:514:GLU:O	1:A:515:LEU:C	2.51	0.47
1:B:499:MSE:HE3	2:B:31:HOH:O	2.14	0.47
1:B:441:HIS:CE1	1:B:547:LEU:HD11	2.50	0.47

Continued on next page...



Continued from previous page...

	A. O	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \; (\mathring{\rm A})$	overlap (Å)
1:B:453:LEU:HD12	1:B:454:ILE:H	1.79	0.47
1:B:499:MSE:HE2	1:B:505:ILE:HG13	1.95	0.47
1:A:516:ARG:CZ	1:A:517:GLY:H	2.29	0.47
1:B:518:ILE:HD12	1:B:519:MSE:H	1.81	0.46
1:B:454:ILE:CD1	1:B:454:ILE:N	2.72	0.45
1:A:453:LEU:HA	2:A:14:HOH:O	2.17	0.44
1:B:555:LEU:HD11	1:B:559:MSE:HE2	1.99	0.44
1:B:499:MSE:HE2	1:B:505:ILE:CG1	2.48	0.44
1:B:453:LEU:HD12	1:B:454:ILE:N	2.33	0.44
1:B:428:ARG:HG3	1:B:428:ARG:HH11	1.83	0.43
1:B:474:ILE:HD11	1:B:559:MSE:CE	2.40	0.43
1:A:440:LYS:NZ	2:A:43:HOH:O	2.50	0.43
1:A:439:GLU:HG2	1:A:444:TYR:O	2.19	0.43
1:B:411:TYR:C	1:B:411:TYR:CD2	2.92	0.43
1:B:502:THR:O	1:B:502:THR:HG22	2.19	0.42
1:A:514:GLU:O	1:A:514:GLU:HG3	2.20	0.42
1:A:423:THR:HG22	1:A:424:GLY:N	2.35	0.42
1:B:432:GLU:C	1:B:435:PRO:HD2	2.41	0.42
1:B:518:ILE:CD1	1:B:519:MSE:H	2.33	0.41
1:B:534:LEU:HD22	1:B:535:THR:N	2.36	0.41
1:B:506:LYS:HB3	1:B:506:LYS:HE2	1.85	0.41
1:B:552:TRP:CZ3	1:B:555:LEU:HD23	2.56	0.41
1:A:492:LEU:HD11	1:B:527:LEU:HA	2.02	0.41
1:A:485:ARG:HD3	1:B:488:SER:HB2	2.03	0.41
1:B:519:MSE:O	1:B:523:GLU:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	148/159 (93%)	137 (93%)	7 (5%)	4 (3%)	5 3

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	В	147/159 (92%)	141 (96%)	6 (4%)	0	100	100
All	All	295/318 (93%)	278 (94%)	13 (4%)	4 (1%)	11	11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	515	LEU
1	A	547	LEU
1	A	548	ASN
1	A	560	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	141/143 (99%)	137 (97%)	4 (3%)	43 60
1	В	140/143 (98%)	134 (96%)	6 (4%)	29 40
All	All	$281/286 \ (98\%)$	271 (96%)	10 (4%)	35 49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	415	ASP
1	A	427	GLU
1	A	516	ARG
1	A	548	ASN
1	В	409	LEU
1	В	411	TYR
1	В	454	ILE
1	В	504	GLU
1	В	518	ILE
1	В	538	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	480	ASN
1	A	522	GLN
1	A	548	ASN
1	В	441	HIS
1	В	469	GLN
1	В	522	GLN
1	В	548	ASN
1	В	556	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

## 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

