



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2023 – 09:56 PM EDT

PDB ID : 5T78
Title : Crystal structure of therapeutic mAB AR20.5 in complex with MUC1 peptide
Authors : Brooks, C.L.; Movahedin, M.
Deposited on : 2016-09-02
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	94% 6%
1	C	216	91% 9%
2	B	215	91% 5% . .
2	D	215	% 91% 6% .
3	E	8	62% 25% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	8	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NGA	E	101	X	-	-	-
4	NGA	F	101	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13233 atoms, of which 6506 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab fragment AR20.5 - Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	216	3324	1059	1640	286	333	6	0	0	0
1	C	216	3324	1059	1640	286	333	6	0	0	0


- Molecule 2 is a protein called Fab Fragment - AR20.5 - Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	209	3149	1018	1557	257	310	7	0	0	0
2	D	209	3129	1013	1543	256	310	7	0	0	0

- Molecule 3 is a protein called MUC1 Glycopeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	F	7	101	32	49	10	10	0	0	0
3	E	7	101	32	49	10	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-galactopyranose (three-letter code: NGA) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	H	N			O
4	F	1	28	8	14	1	5	0	0
4	E	1	28	8	14	1	5	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	13	13	13	0	0
5	B	11	11	11	0	0
5	C	14	14	14	0	0
5	D	11	11	11	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fab fragment AR20.5 - Light Chain

Chain A:  94% 6%




- Molecule 1: Fab fragment AR20.5 - Light Chain

Chain C:  91% 9%




- Molecule 2: Fab Fragment - AR20.5 - Heavy chain

Chain B:  91% 5% . .





- Molecule 2: Fab Fragment - AR20.5 - Heavy chain

Chain D:  % 91% 6% .



- Molecule 3: MUC1 Glycopeptide

Chain F:  12% 75% 12% 12%



- Molecule 3: MUC1 Glycopeptide



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	97.25Å 100.46Å 235.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.12 – 2.20 49.12 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.12-2.20) 99.9 (49.12-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.20Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.194 , 0.221 0.194 , 0.221	Depositor DCC
R_{free} test set	2945 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	50.3	Xtrriage
Anisotropy	0.502	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.084 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13233	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NGA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/1724	0.62	1/2338 (0.0%)
1	C	0.47	0/1724	0.64	0/2338
2	B	0.44	0/1637	0.60	0/2239
2	D	0.44	0/1631	0.60	0/2231
3	E	0.47	0/54	0.74	0/74
3	F	0.50	0/54	0.85	0/74
All	All	0.46	0/6824	0.62	1/9294 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	LEU	CB-CG-CD2	-5.16	102.22	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1684	1640	1640	5	0
1	C	1684	1640	1640	10	0
2	B	1592	1557	1557	7	0
2	D	1586	1543	1541	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	52	49	49	2	0
3	F	52	49	49	3	0
4	E	14	14	13	0	0
4	F	14	14	13	2	0
5	A	13	0	0	0	0
5	B	11	0	0	0	0
5	C	14	0	0	0	0
5	D	11	0	0	0	0
All	All	6727	6506	6502	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:156:THR:HG22	2:B:199:ASN:HB2	1.83	0.59
2:D:156:THR:HG22	2:D:199:ASN:HB2	1.86	0.58
2:B:100:TYR:HE1	4:F:101:NGA:H61	1.68	0.58
1:C:66:ARG:NH1	1:C:87:ASP:OD2	2.35	0.57
2:B:100:TYR:CE1	4:F:101:NGA:H61	2.39	0.57
2:B:118:LYS:HG3	2:B:119:THR:N	2.20	0.55
1:A:96:GLY:HA3	3:F:5:ARG:HD2	1.89	0.54
1:C:165:LEU:HD11	2:D:172:VAL:HG11	1.90	0.53
1:C:96:GLY:HA3	3:E:5:ARG:HD2	1.89	0.53
1:A:74:THR:OG1	2:D:174:GLN:HG2	2.11	0.51
1:C:215:ASN:O	1:C:216:ARG:HB2	2.11	0.49
1:A:215:ASN:O	1:A:216:ARG:HB2	2.12	0.49
2:D:67:ARG:NH2	2:D:90:ASP:OD2	2.48	0.47
2:B:100:TYR:HB3	2:B:104:ASP:OD2	2.14	0.47
1:A:39:GLU:OE2	3:F:5:ARG:NH1	2.48	0.46
1:C:154:LYS:HB2	1:C:198:THR:HB	1.99	0.45
1:C:13:VAL:HG11	1:C:109:LEU:HD11	1.98	0.45
2:B:2:VAL:O	2:B:3:LYS:HD2	2.17	0.44
2:D:1:GLU:O	2:D:2:VAL:HB	2.16	0.44
2:B:103:PHE:CE1	3:F:5:ARG:NH2	2.86	0.43
1:C:37:TYR:CZ	3:E:6:PRO:HD3	2.54	0.43
1:C:39:GLU:HG2	1:C:54:TYR:HA	2.00	0.42
1:C:113:ARG:HG3	1:C:114:ALA:O	2.19	0.42
1:C:94:PHE:CZ	1:C:101:TRP:HB3	2.56	0.41
1:A:13:VAL:HG11	1:A:109:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/216 (99%)	208 (97%)	6 (3%)	0	100	100
1	C	214/216 (99%)	207 (97%)	7 (3%)	0	100	100
2	B	205/215 (95%)	200 (98%)	4 (2%)	1 (0%)	29	31
2	D	205/215 (95%)	201 (98%)	3 (2%)	1 (0%)	29	31
3	E	5/8 (62%)	5 (100%)	0	0	100	100
3	F	5/8 (62%)	5 (100%)	0	0	100	100
All	All	848/878 (97%)	826 (97%)	20 (2%)	2 (0%)	47	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	2	VAL
2	B	2	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	189 (98%)	4 (2%)	53	67
1	C	193/193 (100%)	191 (99%)	2 (1%)	76	86
2	B	181/184 (98%)	175 (97%)	6 (3%)	38	49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	180/184 (98%)	174 (97%)	6 (3%)	38	49
3	E	6/6 (100%)	6 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	759/766 (99%)	741 (98%)	18 (2%)	49	62

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	SER
1	A	38	LEU
1	A	127	SER
1	A	132	SER
2	B	3	LYS
2	B	59	TYR
2	B	118	LYS
2	B	143	CYS
2	B	181	SER
2	B	193	SER
1	C	10	SER
1	C	38	LEU
2	D	3	LYS
2	D	59	TYR
2	D	143	CYS
2	D	181	SER
2	D	198	CYS
2	D	211	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	195	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NGA	E	101	3	14,14,15	0.57	0	17,19,21	1.50	3 (17%)
4	NGA	F	101	3	14,14,15	0.58	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '–' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NGA	E	101	3	1/1/7/7	2/6/23/26	0/1/1/1
4	NGA	F	101	3	1/1/7/7	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	101	NGA	C1-O5-C5	4.54	118.34	112.19
4	E	101	NGA	C3-C4-C5	2.37	114.47	110.24
4	E	101	NGA	O3-C3-C2	-2.24	104.83	109.47

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	F	101	NGA	C1
4	E	101	NGA	C1

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	101	NGA	C4-C5-C6-O6
4	F	101	NGA	O5-C5-C6-O6
4	E	101	NGA	O5-C5-C6-O6
4	E	101	NGA	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	101	NGA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/216 (100%)	0.16	0 100 100	41, 62, 87, 99	0
1	C	216/216 (100%)	0.17	0 100 100	41, 61, 81, 97	0
2	B	209/215 (97%)	0.07	1 (0%) 91 90	46, 65, 92, 135	0
2	D	209/215 (97%)	0.07	3 (1%) 75 73	43, 62, 88, 129	0
3	E	7/8 (87%)	0.37	0 100 100	58, 60, 100, 116	0
3	F	7/8 (87%)	0.44	1 (14%) 2 2	66, 68, 96, 100	0
All	All	864/878 (98%)	0.12	5 (0%) 89 88	41, 63, 88, 135	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	GLU	2.9
3	F	2	PRO	2.3
2	D	1	GLU	2.2
2	D	194	GLU	2.2
2	D	214	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NGA	F	101	14/15	0.93	0.12	51,64,83,87	0
4	NGA	E	101	14/15	0.96	0.12	51,65,82,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.