

wwPDB X-ray Structure Validation Summary Report (i)

Aug 8, 2020 – 10:27 PM BST

PDB ID : 6TBB

Title : Crystal structure of S. aureus Fabl in complex with NADPH and kalimantacin

A (batumin)

Authors : Fage, C.D.; Masschelein, J.

Deposited on : 2019-11-01

Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{resolution range}(ext{Å}))$
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	Α.	001	7%	
I	A	261	93%	6% •
1	В	261	10%	
I	D	201	95%	5%
-		001	15%	
1	С	261	94%	5% •
	ъ	201	5%	
1	D	261	92%	7% •
	_		7%	
1	E	261	94%	5% •
			9%	
1	F	261	95%	• •





Continued from previous page...

0 0 1000	$J = I \cup I \cup I$	J. o F. oo to do	F ## 3 0	
Mol	Chain	ain Length	Quality of chain	
1	G	G 261	95%	•
1	Н	H 261	91%	8% •



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 16633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Enoyl-[acyl-carrier-protein] reductase [NADPH].

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	259	Total	С	N	О	S	0	0	0
1	A	259	1992	1255	345	387	5	0	U	
1	В	261	Total	С	N	О	S	0	0	0
1	Б	201	2002	1260	347	390	5	0	U	
1	С	259	Total	С	N	О	S	0	0	0
1		209	1992	1255	345	387	5	0	0	U
1	D	258	Total	С	N	О	S	0	0	0
1	D	200	1982	1249	342	386	5	0	U	U
1	Е	258	Total	С	N	О	S	0	0	0
1	L'	200	1982	1249	342	386	5	U	U	
1	F	258	Total	С	N	О	S	0	0	0
1	L	200	1982	1249	342	386	5	U	U	0
1	G	260	Total	С	N	О	S	0	0	0
1	G	200	1998	1258	346	389	5	0	0	
1	Н	259	Total	С	N	О	S	0	0	0
1	11	209	1992	1255	345	387	5		U	

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP A0A0J9X1X7
A	-3	SER	-	expression tag	UNP A0A0J9X1X7
A	-2	HIS	-	expression tag	UNP A0A0J9X1X7
A	-1	MET	_	expression tag	UNP A0A0J9X1X7
A	0	ALA	-	expression tag	UNP A0A0J9X1X7
A	1	SER	-	expression tag	UNP A0A0J9X1X7
A	2	LEU	-	expression tag	UNP A0A0J9X1X7
В	-4	GLY	-	expression tag	UNP A0A0J9X1X7
В	-3	SER	-	expression tag	UNP A0A0J9X1X7
В	-2	HIS	-	expression tag	UNP A0A0J9X1X7
В	-1	MET	-	expression tag	UNP A0A0J9X1X7
В	0	ALA	-	expression tag	UNP A0A0J9X1X7
В	1	SER	-	expression tag	UNP A0A0J9X1X7



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	2	LEU	_	expression tag	UNP A0A0J9X1X7
С	-4	GLY	-	expression tag	UNP A0A0J9X1X7
С	-3	SER	_	expression tag	UNP A0A0J9X1X7
С	-2	HIS	_	expression tag	UNP A0A0J9X1X7
С	-1	MET	_	expression tag	UNP A0A0J9X1X7
С	0	ALA	_	expression tag	UNP A0A0J9X1X7
С	1	SER	_	expression tag	UNP A0A0J9X1X7
С	2	LEU	-	expression tag	UNP A0A0J9X1X7
D	-4	GLY	-	expression tag	UNP A0A0J9X1X7
D	-3	SER	-	expression tag	UNP A0A0J9X1X7
D	-2	HIS	-	expression tag	UNP A0A0J9X1X7
D	-1	MET	_	expression tag	UNP A0A0J9X1X7
D	0	ALA	-	expression tag	UNP A0A0J9X1X7
D	1	SER	-	expression tag	UNP A0A0J9X1X7
D	2	LEU	-	expression tag	UNP A0A0J9X1X7
Е	-4	GLY	-	expression tag	UNP A0A0J9X1X7
Е	-3	SER	_	expression tag	UNP A0A0J9X1X7
Е	-2	HIS	-	expression tag	UNP A0A0J9X1X7
Е	-1	MET	-	expression tag	UNP A0A0J9X1X7
Е	0	ALA	=	expression tag	UNP A0A0J9X1X7
Е	1	SER	_	expression tag	UNP A0A0J9X1X7
Е	2	LEU	-	expression tag	UNP A0A0J9X1X7
F	-4	GLY	_	expression tag	UNP A0A0J9X1X7
F	-3	SER	_	expression tag	UNP A0A0J9X1X7
F	-2	HIS	_	expression tag	UNP A0A0J9X1X7
F	-1	MET	_	expression tag	UNP A0A0J9X1X7
F	0	ALA	_	expression tag	UNP A0A0J9X1X7
F	1	SER	_	expression tag	UNP A0A0J9X1X7
F	2	LEU	_	expression tag	UNP A0A0J9X1X7
G	-4	GLY	_	expression tag	UNP A0A0J9X1X7
G	-3	SER	_	expression tag	UNP A0A0J9X1X7
G	-2	HIS	_	expression tag	UNP A0A0J9X1X7
G	-1	MET	_	expression tag	UNP A0A0J9X1X7
G	0	ALA	_	expression tag	UNP A0A0J9X1X7
G	1	SER	_	expression tag	UNP A0A0J9X1X7
G	2	LEU	-	expression tag	UNP A0A0J9X1X7
Н	-4	GLY	-	expression tag	UNP A0A0J9X1X7
Н	-3	SER	-	expression tag	UNP A0A0J9X1X7
Н	-2	HIS		expression tag	UNP A0A0J9X1X7
Н	-1	MET	-	expression tag	UNP A0A0J9X1X7
Н	0	ALA	-	expression tag	UNP A0A0J9X1X7
Н	1	SER	-	expression tag	UNP A0A0J9X1X7



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Н	2	LEU	-	expression tag	UNP A0A0J9X1X7


• Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
2	A	1	48	21	7	17	3	U	0
2	В	1	Total	С	N	О	Р	0	0
	Б	1	48	21	7	17	3	0	0
2	С	1	Total	С	N	О	Р	0	0
		1	48	21	7	17	3	0	0
2	D	1	Total	С	N	О	Р	0	0
	ע	1	48	21	7	17	3	U	
2	Е	1	Total	С	N	О	Р	0	0
	ت ا	1	48	21	7	17	3	0	0
2	F	1	Total	С	N	О	Р	0	0
	I'	1	48	21	7	17	3	0	0
2	G	1	Total	С	N	О	Р	0	0
	G	1	48	21	7	17	3	0	
2	Н	1	Total	С	N	О	Р	0	0
	11	I.	48	21	7	17	3	U	U

• Molecule 3 is Kalimantacin (three-letter code: KAL) (formula: $C_{30}H_{48}N_2O_7$) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
3	A	1	Total	С	N	О	0	0
3	A	1	39	30	2	7	U	U
3	В	1	Total	С	N	О	0	0
)	Б	1	39	30	2	7	U	0
3	С	1	Total	С	N	О	0	0
)		1	39	30	2	7	U	0
3	D	1	Total	С	N	О	0	0
)	ש	1	39	30	2	7	U	U
3	Е	1	Total	С	N	О	0	0
)	خ1 ا	1	39	30	2	7	U	U
3	F	1	Total	С	N	О	0	0
)	1'	1	39	30	2	7	0	
3	G	1	Total	С	N	О	0	0
0	G	1	39	30	2	7	U	U
3	Н	1	Total	С	N	О	0	0
3	11	1	39	30	2	7	0	

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	С	1	Total O 1 1	0	0
4	D	1	Total O 1 1	0	0
4	E	3	Total O 3 3	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	F	1	Total O 1 1	0	0
4	G	2	Total O 2 2	0	0
4	Н	3	Total O 3 3	0	0




Chain D:

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADPH]




92%

• Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADPH]



7% •





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	63.91Å 107.92Å 294.23Å	Danagitan
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 - 2.45	Depositor
Resolution (A)	29.86 - 2.45	EDS
% Data completeness	99.8 (29.88-2.45)	Depositor
(in resolution range)	99.9 (29.86-2.45)	EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.25 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
D D.	0.196 , 0.247	Depositor
R, R_{free}	0.195 , 0.241	DCC
R_{free} test set	3746 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	81.8	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 55.0	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	16633	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, KAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.70	0/2020	0.80	0/2722	
1	В	0.70	0/2030	0.80	0/2735	
1	С	0.72	0/2020	0.79	0/2722	
1	D	0.72	0/2009	0.79	0/2707	
1	Е	0.70	0/2009	0.82	0/2707	
1	F	0.73	0/2009	0.80	0/2707	
1	G	0.71	0/2026	0.82	0/2730	
1	Н	0.70	0/2020	0.80	0/2722	
All	All	0.71	0/16143	0.80	0/21752	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Α	1992	0	2008	13	0
1	В	2002	0	2016	11	1
1	С	1992	0	2008	10	0
1	D	1982	0	2001	14	0
1	Ε	1982	0	2001	9	1



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1982	0	2001	8	1
1	G	1998	0	2013	10	0
1	Н	1992	0	2008	15	1
2	A	48	0	26	1	0
2	В	48	0	26	2	0
2	С	48	0	26	4	0
2	D	48	0	26	2	0
2	Ε	48	0	26	2	0
2	F	48	0	26	1	0
2	G	48	0	26	3	0
2	Н	48	0	26	4	0
3	A	39	0	0	0	0
3	В	39	0	0	0	0
3	С	39	0	0	2	0
3	D	39	0	0	1	0
3	Ε	39	0	0	0	0
3	F	39	0	0	0	0
3	G	39	0	0	0	0
3	Н	39	0	0	0	0
4	A	4	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
4	Е	3	0	0	0	0
4	F	1	0	0	0	0
4	G	2	0	0	0	0
4	Н	3	0	0	0	0
All	All	16633	0	16264	77	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	Clash overlap (Å)	
1:G:197:SER:HB2	2:G:301:NDP:O1A	1.75	0.87	
1:H:197:SER:HB2	2:H:301:NDP:O1A	1.81	0.79	
1:D:197:SER:HB2	2:D:301:NDP:O2A	1.82	0.79	
1:B:197:SER:HB2	2:B:301:NDP:O2A	1.84	0.77	
1:H:16:ASN:HA	1:H:47:GLU:HG2	1.70	0.73	

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-



metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
1:B:1:SER:OG	1:E:5:GLU:OE1[2_455]	1.98	0.22
1:F:41:LYS:NZ	1:H:213:GLU:OE2[4_554]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	${f ntiles}$
1	A	257/261~(98%)	246 (96%)	11 (4%)	0	100	100
1	В	259/261~(99%)	247 (95%)	12 (5%)	0	100	100
1	С	257/261 (98%)	247 (96%)	10 (4%)	0	100	100
1	D	256/261~(98%)	244 (95%)	12 (5%)	0	100	100
1	E	256/261~(98%)	244 (95%)	12 (5%)	0	100	100
1	F	256/261~(98%)	243 (95%)	13 (5%)	0	100	100
1	G	$258/261 \; (99\%)$	247 (96%)	11 (4%)	0	100	100
1	Н	257/261~(98%)	245 (95%)	12 (5%)	0	100	100
All	All	$2056/2088 \; (98\%)$	1963 (96%)	93 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	A	$212/213 \; (100\%)$	211 (100%)	1 (0%)	88	93
1	В	$213/213 \; (100\%)$	213 (100%)	0	100	100
1	С	$212/213 \; (100\%)$	211 (100%)	1 (0%)	88	93
1	D	$211/213 \ (99\%)$	211 (100%)	0	100	100
1	E	$211/213\ (99\%)$	211 (100%)	0	100	100
1	F	$211/213 \ (99\%)$	211 (100%)	0	100	100
1	G	$213/213\ (100\%)$	213 (100%)	0	100	100
1	Н	$212/213 \; (100\%)$	211 (100%)	1 (0%)	88	93
All	All	1695/1704~(100%)	1692 (100%)	3 (0%)	93	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	MET
1	С	135	LEU
1	Н	43	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
1	A	30	GLN
1	F	155	GLN
1	G	30	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	ond leng	$_{ m ths}$	В	ond ang	gles
WIOI	Type	Chain	rtes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	E	301	_	45,52,52	1.51	6 (13%)	53,80,80	1.55	11 (20%)
3	KAL	Н	302	_	35,38,38	1.94	6 (17%)	38,48,48	1.68	6 (15%)
2	NDP	С	301	-	45,52,52	1.40	4 (8%)	53,80,80	1.61	10 (18%)
2	NDP	A	301	-	45,52,52	1.34	4 (8%)	53,80,80	1.70	16 (30%)
2	NDP	Н	301	-	45,52,52	1.35	5 (11%)	53,80,80	1.70	15 (28%)
2	NDP	F	301	-	45,52,52	1.47	5 (11%)	53,80,80	1.51	8 (15%)
2	NDP	D	301	-	45,52,52	1.46	4 (8%)	53,80,80	1.55	12 (22%)
2	NDP	В	301	-	45,52,52	1.34	4 (8%)	53,80,80	1.68	12 (22%)
3	KAL	В	302	-	35,38,38	2.04	9 (25%)	38,48,48	1.68	7 (18%)
3	KAL	A	302	-	35,38,38	2.15	7 (20%)	38,48,48	1.94	11 (28%)
3	KAL	G	302	-	35,38,38	1.89	6 (17%)	38,48,48	1.44	4 (10%)
3	KAL	F	302	-	35,38,38	1.91	6 (17%)	38,48,48	1.69	8 (21%)
3	KAL	Е	302	-	35,38,38	1.80	7 (20%)	38,48,48	1.65	7 (18%)
3	KAL	D	302	-	35,38,38	1.86	5 (14%)	38,48,48	1.56	8 (21%)
2	NDP	G	301	-	45,52,52	1.50	6 (13%)	53,80,80	1.83	14 (26%)
3	KAL	С	302	-	35,38,38	2.06	6 (17%)	38,48,48	1.64	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	\mathbf{Type}	Chain	Res	Link	Chirals	${f Torsions}$	Rings
2	NDP	Ε	301	-	-	5/30/77/77	0/5/5/5
3	KAL	Н	302	-	-	7/45/47/47	-
2	NDP	С	301	-	-	10/30/77/77	0/5/5/5
2	NDP	A	301	-	-	13/30/77/77	0/5/5/5



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	Н	301	-	-	7/30/77/77	0/5/5/5
2	NDP	F	301	-	-	9/30/77/77	0/5/5/5
2	NDP	D	301	-	-	2/30/77/77	0/5/5/5
2	NDP	В	301	-	-	5/30/77/77	0/5/5/5
3	KAL	В	302	-	-	7/45/47/47	-
3	KAL	A	302	-	-	5/45/47/47	-
3	KAL	G	302	_	-	1/45/47/47	-
3	KAL	F	302	_	-	2/45/47/47	-
3	KAL	E	302	-	-	1/45/47/47	-
3	KAL	D	302	-	-	2/45/47/47	-
2	NDP	G	301	-	-	8/30/77/77	0/5/5/5
3	KAL	С	302	_	-	2/45/47/47	-

The worst 5 of 90 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	${ m Observed(\AA)}$	$\operatorname{Ideal}(ext{\AA})$
3	D	302	KAL	C27-N28	7.07	1.46	1.33
3	Н	302	KAL	C27-N28	6.73	1.46	1.33
3	В	302	KAL	C27-N28	6.64	1.45	1.33
3	G	302	KAL	C27-N28	6.48	1.45	1.33
3	A	302	KAL	C27-N28	6.45	1.45	1.33

The worst 5 of 156 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	Н	302	KAL	O26-C27-N28	5.47	119.09	110.58
3	A	302	KAL	O26-C27-N28	5.38	118.94	110.58
3	Е	302	KAL	O26-C27-N28	5.24	118.73	110.58
3	В	302	KAL	O26-C27-N28	4.94	118.26	110.58
3	С	302	KAL	O26-C27-N28	4.81	118.06	110.58

There are no chirality outliers.

5 of 86 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	301	NDP	C4B-C5B-O5B-PA
2	E	301	NDP	C2B-O2B-P2B-O1X
3	Н	302	KAL	C22-C24-O26-C27
3	Н	302	KAL	N28-C27-O26-C24



Continued from previous page...

Mol	Chain	Res	Type	${f Atoms}$
2	С	301	NDP	C4B-C5B-O5B-PA


There are no ring outliers.


10 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	301	NDP	2	0
2	С	301	NDP	4	0
2	A	301	NDP	1	0
2	Н	301	NDP	4	0
2	F	301	NDP	1	0
2	D	301	NDP	2	0
2	В	301	NDP	2	0
3	D	302	KAL	1	0
2	G	301	NDP	3	0
3	С	302	KAL	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	259/261~(99%)	0.24	19 (7%) 15 1	11	61, 85, 134, 160	0
1	В	$261/261 \; (100\%)$	0.25	25 (9%) 8 5	5	63, 88, 134, 191	1 (0%)
1	С	259/261~(99%)	0.35	38 (14%) 2	1	65, 86, 141, 172	0
1	D	258/261 (98%)	0.06	12 (4%) 31 2	29	59, 83, 131, 151	0
1	E	258/261~(98%)	0.20	19 (7%) 14 1	11	54, 80, 130, 146	0
1	F	258/261~(98%)	0.20	23 (8%) 9	5	55, 82, 127, 152	1 (0%)
1	G	260/261~(99%)	0.36	25 (9%) 8 5	5	57, 83, 128, 176	0
1	Н	$259/261 \ (99\%)$	0.54	34 (13%) 3	2	61, 92, 147, 176	0
All	All	$2072/2088 \; (99\%)$	0.27	195 (9%) 8	5	54, 85, 135, 191	2 (0%)

The worst 5 of 195 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	93	SER	6.1
1	G	12	MET	6.1
1	Н	12	MET	5.5
1	G	92	HIS	5.4
1	G	94	ILE	5.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	KAL	G	302	39/39	0.87	0.19	57,72,93,98	0
3	KAL	A	302	39/39	0.92	0.19	74,88,94,95	0
3	KAL	В	302	39/39	0.93	0.18	73,86,101,104	0
3	KAL	Н	302	39/39	0.93	0.17	68,92,107,108	0
2	NDP	Н	301	48/48	0.93	0.14	61,92,123,143	0
3	KAL	F	302	39/39	0.93	0.16	69,82,99,103	0
3	KAL	С	302	39/39	0.93	0.15	71,81,102,111	0
3	KAL	D	302	39/39	0.94	0.15	59,74,92,100	0
3	KAL	E	302	39/39	0.94	0.15	53,73,87,94	0
2	NDP	В	301	48/48	0.95	0.16	63,82,98,104	0
2	NDP	С	301	48/48	0.96	0.14	64,79,101,104	0
2	NDP	A	301	48/48	0.96	0.15	65,78,97,107	0
2	NDP	E	301	48/48	0.96	0.15	59,72,87,93	0
2	NDP	G	301	48/48	0.96	0.16	63,73,85,97	0
2	NDP	F	301	48/48	0.96	0.14	55,82,98,104	0
2	NDP	D	301	48/48	0.98	0.14	68,78,94,110	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around KAL G 302: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around KAL A 302: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)














Electron density around NDP H 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around KAL F 302:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)















Electron density around KAL E 302: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around NDP B 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)











Electron density around NDP G 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around NDP F 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c \ (\mathrm{at}\ 0.7\ \mathrm{rmsd}) \ \mathrm{in}\ \mathrm{gray}$

 ${
m mF}_o{
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NDP D 301:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

