



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 23, 2023 – 03:59 PM EDT

PDB ID : 5TER
Title : Crystal Structure of HIV-1 Reverse Transcriptase in Complex with 5-chloro-7-(2-(2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy)phenoxy)-8-methyl-2-naphthonitrile (JLJ651), a Non-nucleoside Inhibitor
Authors : Chan, A.H.; Anderson, K.S.
Deposited on : 2016-09-22
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	
2	B	428	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7788 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 REVERSE TRANSCRIPTASE, P66 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	547	4415	2861	728	818	8	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	initiating methionine	UNP P03366
A	0	VAL	-	expression tag	UNP P03366
A	172	ALA	LYS	engineered mutation	UNP P03366
A	173	ALA	LYS	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 2 is a protein called HIV-1 REVERSE TRANSCRIPTASE, P51 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	401	3319	2160	550	603	6	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is 5-chloro-7-(2-(2-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)ethoxy)phenoxy)-8-methyl-2-naphthonitrile (three-letter code: 7AY) (formula: C₂₄H₁₈ClN₃O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
3	A	1	32	24	1	3	4	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	10	Total O 10 10	0	0
6	B	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HIV-1 REVERSE TRANSCRIPTASE, P66 SUBUNIT



- Molecule 2: HIV-1 REVERSE TRANSCRIPTASE, P51 SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.62Å 74.20Å 108.60Å 90.00° 99.98° 90.00°	Depositor
Resolution (Å)	41.29 – 2.70 41.29 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.29-2.70) 99.8 (41.29-2.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.230 , 0.267 0.231 , 0.267	Depositor DCC
R_{free} test set	1759 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	74.5	Xtrriage
Anisotropy	0.328	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7788	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, 7AY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/4532	0.39	0/6172
2	B	0.22	0/3413	0.39	0/4635
All	All	0.22	0/7945	0.39	0/10807

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4415	0	4419	29	0
2	B	3319	0	3341	26	0
3	A	32	0	0	0	0
4	A	5	0	0	1	0
5	A	1	0	0	0	0
6	A	10	0	0	0	0
6	B	6	0	0	0	0
All	All	7788	0	7760	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:544:GLY:HA2	2:B:286:THR:HG22	1.76	0.66
1:A:91:GLN:O	1:A:91:GLN:NE2	2.29	0.66
1:A:28:GLU:HG3	1:A:135:ILE:HD12	1.80	0.64
2:B:308:GLU:HA	2:B:311:LYS:HE2	1.80	0.64
2:B:107:THR:HG1	2:B:198:HIS:HE2	1.50	0.60
1:A:490:GLY:O	1:A:528:LYS:NZ	2.31	0.55
1:A:363:ASN:HD21	1:A:366:LYS:HE3	1.72	0.54
1:A:80:LEU:O	1:A:84:THR:OG1	2.23	0.54
1:A:405:TYR:CE2	1:A:407:GLN:HB2	2.43	0.54
1:A:512:LYS:NZ	4:A:602:SO4:O3	2.40	0.53
1:A:369:THR:O	1:A:373:GLN:HG2	2.08	0.53
2:B:275:LYS:HD3	2:B:276:VAL:N	2.23	0.53
2:B:101:LYS:HD3	2:B:382:ILE:HG23	1.90	0.53
2:B:169:GLU:OE1	2:B:173:LYS:NZ	2.35	0.53
2:B:20:LYS:HE2	2:B:55:PRO:HB2	1.90	0.51
2:B:107:THR:OG1	2:B:198:HIS:NE2	2.39	0.51
1:A:543:GLY:HA3	2:B:284:ARG:HA	1.93	0.51
2:B:72:ARG:NH2	2:B:151:GLN:OE1	2.43	0.51
2:B:393:ILE:HD13	2:B:398:TRP:HB2	1.92	0.50
1:A:178:ILE:HG22	1:A:191:SER:HB3	1.93	0.50
2:B:73:LYS:NZ	2:B:146:TYR:OH	2.44	0.50
2:B:168:LEU:HD13	2:B:180:ILE:HG21	1.93	0.50
2:B:275:LYS:HD3	2:B:276:VAL:H	1.76	0.50
2:B:21:VAL:HB	2:B:59:PRO:HD3	1.93	0.49
2:B:425:LEU:HD23	2:B:428:GLN:HB3	1.95	0.48
1:A:191:SER:OG	1:A:198:HIS:ND1	2.38	0.47
1:A:88:TRP:CD1	2:B:143:ARG:HD2	2.50	0.47
1:A:171:PHE:O	1:A:175:ASN:ND2	2.37	0.46
2:B:271:TYR:HA	2:B:272:PRO:HD3	1.79	0.46
1:A:443:ASP:HB3	1:A:550:LYS:HD3	1.98	0.46
2:B:241:VAL:HG12	2:B:242:GLN:H	1.80	0.46
1:A:342:TYR:HB3	1:A:348:ASN:HA	1.98	0.46
1:A:19:PRO:HG3	1:A:80:LEU:HB2	1.99	0.45
2:B:203:GLU:HA	2:B:206:ARG:HG2	1.98	0.45
1:A:454:LYS:HZ2	1:A:552:VAL:HB	1.82	0.44
1:A:317:VAL:HG23	1:A:349:LEU:HD23	1.99	0.44
1:A:503:LEU:HD22	1:A:535:TRP:HB2	1.99	0.43
1:A:341:ILE:O	1:A:349:LEU:N	2.52	0.43
2:B:275:LYS:HD2	2:B:277:ARG:HG2	2.01	0.43
2:B:163:SER:O	2:B:167:ILE:HG13	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:LYS:HD3	1:A:223:LYS:HA	1.72	0.42
1:A:475:GLN:HB3	1:A:501:TYR:CE2	2.54	0.42
2:B:423:VAL:HA	2:B:426:TRP:CE3	2.54	0.42
1:A:96:HIS:HA	1:A:97:PRO:HD3	1.89	0.42
1:A:54:ASN:HB3	1:A:143:ARG:HH21	1.84	0.41
2:B:167:ILE:HG12	2:B:212:TRP:CE3	2.55	0.41
1:A:38:CYS:SG	1:A:132:ILE:HD11	2.61	0.41
1:A:312:GLU:HA	1:A:313:PRO:HD3	1.83	0.41
2:B:281:LYS:HG3	2:B:284:ARG:NH1	2.36	0.41
1:A:40:GLU:HG3	1:A:41:MET:N	2.35	0.40
2:B:136:ASN:HB3	2:B:138:GLU:HG3	2.03	0.40
1:A:363:ASN:ND2	1:A:366:LYS:HE3	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	543/557 (98%)	524 (96%)	19 (4%)	0	100	100
2	B	395/428 (92%)	384 (97%)	11 (3%)	0	100	100
All	All	938/985 (95%)	908 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	479/495 (97%)	473 (99%)	6 (1%)	69	87
2	B	364/390 (93%)	358 (98%)	6 (2%)	62	85
All	All	843/885 (95%)	831 (99%)	12 (1%)	67	86

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	GLU
1	A	91	GLN
1	A	286	THR
1	A	295	LEU
1	A	546	GLU
1	A	551	LEU
2	B	8	VAL
2	B	174	GLN
2	B	232	TYR
2	B	275	LYS
2	B	417	VAL
2	B	428	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	602	-	4,4,4	0.13	0	6,6,6	0.06	0
3	7AY	A	601	-	35,35,35	1.46	5 (14%)	49,49,49	1.44	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7AY	A	601	-	-	1/12/12/12	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	7AY	C0P-N0H	3.75	1.42	1.37
3	A	601	7AY	CBA-CBC	3.41	1.52	1.44
3	A	601	7AY	O0A-CAL	2.59	1.45	1.39
3	A	601	7AY	CAJ-CAI	-2.42	1.39	1.43
3	A	601	7AY	CAZ-CBA	2.40	1.44	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	7AY	C0E-N0H-C0K	5.75	124.60	118.54
3	A	601	7AY	C0E-N0H-C0P	-3.49	114.43	119.75
3	A	601	7AY	C0D-O0B-C05	3.08	125.23	117.69
3	A	601	7AY	CAM-CAL-CAK	-2.29	119.77	122.45
3	A	601	7AY	C0O-C0P-N0H	-2.11	120.47	122.44
3	A	601	7AY	CAY-CAI-CAH	-2.02	121.47	125.06

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	7AY	C05-C04-O0A-CAL

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	602	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	547/557 (98%)	0.68	71 (12%) 3 2	39, 88, 140, 157	0
2	B	401/428 (93%)	0.49	30 (7%) 14 12	44, 78, 130, 148	0
All	All	948/985 (96%)	0.60	101 (10%) 6 4	39, 83, 137, 157	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	292	VAL	10.7
1	A	254	VAL	7.4
2	B	232	TYR	6.2
2	B	356	ARG	6.2
1	A	257	ILE	5.4
1	A	294	PRO	5.4
1	A	255	ASN	5.3
1	A	291	GLU	5.1
1	A	293	ILE	5.1
2	B	238	LYS	5.1
1	A	300	GLU	5.0
1	A	47	ILE	5.0
1	A	287	LYS	5.0
1	A	2	ILE	4.7
1	A	303	LEU	4.6
1	A	288	ALA	4.5
1	A	286	THR	4.3
1	A	297	GLU	4.3
1	A	214	LEU	4.3
1	A	311	LYS	4.2
1	A	295	LEU	4.2
1	A	290	THR	4.2
1	A	302	GLU	4.2
1	A	252	TRP	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	26	LEU	3.9
1	A	251	SER	3.8
2	B	14	PRO	3.8
2	B	67	ASP	3.8
1	A	246	LEU	3.8
1	A	218	ASP	3.7
2	B	168	LEU	3.7
2	B	237	ASP	3.7
1	A	38	CYS	3.7
2	B	358	ARG	3.6
1	A	259	LYS	3.6
1	A	130	PHE	3.5
1	A	279	LEU	3.4
1	A	74	LEU	3.3
1	A	289	LEU	3.3
1	A	24	TRP	3.3
1	A	144	TYR	3.3
1	A	247	PRO	3.3
1	A	548	VAL	3.3
1	A	253	THR	3.2
2	B	196	GLY	3.1
2	B	207	GLN	3.1
1	A	73	LYS	3.0
2	B	178	ILE	3.0
2	B	240	THR	3.0
1	A	261	VAL	3.0
1	A	552	VAL	3.0
1	A	221	HIS	2.9
2	B	198	HIS	2.9
2	B	239	TRP	2.8
1	A	18	GLY	2.8
2	B	202	ILE	2.8
1	A	299	ALA	2.8
1	A	551	LEU	2.8
1	A	283	LEU	2.8
1	A	19	PRO	2.8
2	B	209	LEU	2.8
1	A	151	GLN	2.7
1	A	547	GLN	2.7
2	B	210	LEU	2.7
1	A	244	ILE	2.7
1	A	132	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	52	PRO	2.6
2	B	359	GLY	2.6
2	B	95	PRO	2.6
2	B	241	VAL	2.6
2	B	199	ARG	2.6
2	B	15	GLY	2.6
1	A	37	ILE	2.6
1	A	263	LYS	2.5
1	A	304	ALA	2.5
2	B	206	ARG	2.5
2	B	85	GLN	2.4
1	A	40	GLU	2.4
1	A	90	VAL	2.4
2	B	88	TRP	2.4
1	A	245	VAL	2.4
1	A	209	LEU	2.4
2	B	195	ILE	2.4
1	A	61	PHE	2.3
1	A	301	LEU	2.3
1	A	15	GLY	2.3
2	B	236	PRO	2.3
2	B	428	GLN	2.3
2	B	211	ARG	2.3
1	A	298	GLU	2.2
1	A	359	GLY	2.2
1	A	127	TYR	2.2
1	A	72	ARG	2.2
1	A	256	ASP	2.1
1	A	285	GLY	2.1
1	A	360	ALA	2.1
1	A	549	ASP	2.1
1	A	29	GLU	2.1
2	B	124	PHE	2.1
1	A	71	TRP	2.0
1	A	550	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	602	5/5	0.89	0.14	103,104,116,119	0
5	MG	A	603	1/1	0.92	0.23	74,74,74,74	0
3	7AY	A	601	32/32	0.96	0.19	60,73,85,90	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.