

Full wwPDB X-ray Structure Validation Report (i)

Aug 15, 2023 – 10:23 PM EDT

PDB ID : 1TK2

Title : Crystal Structure of the Complex formed between Alkaline Proteinase Savinase

and Gramicidin S at 1.5A Resolution

Authors: Bhatt, V.S.; Kaur, P.; Klupsch, S.; Betzel, C.; Brenner, S.; Singh, T.P.

Deposited on : 2004-06-08

Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	A	269		91%	8% •	
2	В	10	40%	50%	10%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ORN	В	2	-	-	X	-
2	ORN	В	7	-	-	X	-
2	DPN	В	9	-	-	X	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called SUBTILISIN SAVINASE.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	269	Total 1879	C 1150	N 347	O 379	S 3	0	0	0

• Molecule 2 is a protein called GRAMICIDIN S.

\mathbf{Mol}	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace			
2	В	10	Total 82	C 60	N 12	O 10	0	0	0	

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ca 2 2	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	190	Total O 190 190	0	0
4	В	2	Total O 2 2	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: SUBTILISIN SAVINASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	76.25Å 73.34Å 40.89Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.32 - 1.54	Depositor
% Data completeness	100.0 (27.32-1.54)	Depositor
(in resolution range)	100.0 (21.02 1.04)	Берозног
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.169 , 0.189	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2155	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ORN, CA, DPN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.74	0/1913	0.76	1/2614 (0.0%)	
2	В	1.06	0/41	2.34	4/49 (8.2%)	
All	All	0.75	0/1954	0.82	5/2663 (0.2%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	3	LEU	CB-CA-C	6.88	123.28	110.20
2	В	3	LEU	CA-CB-CG	-6.32	100.77	115.30
2	В	8	LEU	N-CA-C	-5.76	95.46	111.00
2	В	5	PRO	N-CA-C	5.68	126.87	112.10
1	A	222	MET	CG-SD-CE	-5.17	91.92	100.20

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	В	1	VAL	Peptide
2	В	7	ORN	Peptide
2	В	9	DPN	Mainchain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1879	0	1837	48	0
2	В	82	0	88	83	0
3	A	2	0	0	0	0
4	A	190	0	0	1	0
4	В	2	0	0	0	0
All	All	2155	0	1925	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \mathring{A}) \end{array}$	Clash overlap (Å)
2:B:9:DPN:CD2	2:B:10:PRO:HD3	1.33	1.53
2:B:1:VAL:CG1	2:B:2:ORN:HG3	1.37	1.46
2:B:9:DPN:CE2	2:B:10:PRO:HD3	1.69	1.23
2:B:9:DPN:CD2	2:B:10:PRO:CD	2.19	1.20
2:B:1:VAL:HG12	2:B:2:ORN:CG	1.74	1.16
2:B:1:VAL:CG1	2:B:2:ORN:CG	2.25	1.12
2:B:9:DPN:HD2	2:B:10:PRO:HD3	1.32	1.08
2:B:1:VAL:HG12	2:B:2:ORN:N	1.69	1.06
2:B:2:ORN:C	2:B:3:LEU:HD12	1.86	1.05
1:A:107:ILE:HD11	2:B:8:LEU:HD22	1.45	0.96
2:B:1:VAL:HG12	2:B:2:ORN:HG3	0.98	0.92
1:A:107:ILE:HD11	2:B:8:LEU:CD2	1.99	0.91
2:B:1:VAL:HG11	2:B:2:ORN:HG3	1.50	0.90
2:B:1:VAL:O	2:B:7:ORN:HA	1.72	0.90
1:A:107:ILE:CD1	2:B:8:LEU:HD22	2.01	0.90
1:A:154:GLY:HA3	2:B:3:LEU:CD2	2.06	0.86
2:B:7:ORN:C	2:B:8:LEU:HG	2.06	0.86
1:A:154:GLY:CA	2:B:3:LEU:HD22	2.06	0.85
2:B:3:LEU:HD12	2:B:3:LEU:N	1.92	0.84
1:A:154:GLY:HA3	2:B:3:LEU:HD22	1.58	0.83
1:A:96:LEU:HD13	2:B:6:VAL:CG2	2.10	0.81
2:B:9:DPN:CE2	2:B:10:PRO:CD	2.50	0.81

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		${\rm distance}({\rm \AA})$	overlap(Å)
2:B:1:VAL:CG1	2:B:2:ORN:N	2.43	0.81
1:A:154:GLY:CA	2:B:3:LEU:CD2	2.59	0.78
1:A:126:LEU:HB2	2:B:6:VAL:HG11	1.65	0.77
1:A:191:GLN:NE2	2:B:3:LEU:HD23	2.01	0.76
1:A:96:LEU:HD13	2:B:6:VAL:HG22	1.67	0.76
1:A:191:GLN:NE2	2:B:3:LEU:CD2	2.50	0.74
1:A:155:ASN:HD21	1:A:220:THR:H	1.35	0.74
1:A:126:LEU:HB2	2:B:6:VAL:CG1	2.17	0.74
1:A:156:SER:H	1:A:191:GLN:HE21	1.36	0.74
2:B:7:ORN:O	2:B:8:LEU:HG	1.88	0.74
1:A:191:GLN:HE22	2:B:3:LEU:HD21	1.51	0.73
1:A:126:LEU:CB	2:B:6:VAL:HG11	2.18	0.73
1:A:191:GLN:HE22	2:B:3:LEU:CD2	2.02	0.72
2:B:2:ORN:C	2:B:3:LEU:CD1	2.66	0.72
1:A:96:LEU:CD1	2:B:6:VAL:HG21	2.19	0.72
2:B:1:VAL:HG12	2:B:2:ORN:CB	2.20	0.71
1:A:155:ASN:H	2:B:3:LEU:HD23	1.58	0.68
1:A:125:SER:HB3	1:A:221:SER:OG	1.93	0.68
2:B:1:VAL:H2	2:B:9:DPN:C	2.07	0.67
2:B:5:PRO:O	2:B:6:VAL:HG23	1.94	0.67
1:A:126:LEU:HD13	2:B:8:LEU:HD11	1.76	0.67
2:B:2:ORN:HB2	2:B:7:ORN:HB3	1.74	0.67
1:A:155:ASN:N	2:B:3:LEU:HD23	2.11	0.65
2:B:7:ORN:C	2:B:8:LEU:CG	2.71	0.65
2:B:3:LEU:N	2:B:3:LEU:CD1	2.60	0.65
2:B:5:PRO:O	2:B:6:VAL:CG2	2.46	0.63
1:A:154:GLY:HA2	2:B:3:LEU:HD22	1.78	0.63
2:B:1:VAL:HG23	2:B:10:PRO:C	2.19	0.63
2:B:7:ORN:O	2:B:8:LEU:CG	2.46	0.63
1:A:166:SER:OG	2:B:3:LEU:CD2	2.48	0.61
1:A:166:SER:OG	2:B:3:LEU:HD21	2.00	0.61
1:A:154:GLY:CA	2:B:3:LEU:HD23	2.30	0.61
1:A:155:ASN:ND2	1:A:220:THR:H	1.99	0.60
2:B:2:ORN:O	2:B:2:ORN:HG2	2.01	0.60
2:B:1:VAL:HG23	2:B:10:PRO:O	2.04	0.56
2:B:8:LEU:O	2:B:9:DPN:C	2.51	0.56
2:B:1:VAL:HG13	2:B:2:ORN:HG3	1.68	0.56
2:B:1:VAL:HG13	2:B:2:ORN:CG	2.27	0.56
1:A:96:LEU:HD13	2:B:6:VAL:HG21	1.81	0.55
1:A:130:SER:OG	2:B:10:PRO:HG2	2.06	0.55
1:A:38:THR:HG23	1:A:59:GLN:HE22	1.71	0.54

Continued on next page...



Continued from previous page...

A Lange 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:2:GLN:HE22	1:A:76:ASN:HD22	1.56	0.54
1:A:107:ILE:HD12	2:B:8:LEU:HD22	1.87	0.53
1:A:107:ILE:HD11	2:B:8:LEU:HD21	1.85	0.53
1:A:126:LEU:HD13	2:B:8:LEU:CD1	2.40	0.52
1:A:96:LEU:CD1	2:B:6:VAL:CG2	2.78	0.52
2:B:7:ORN:O	2:B:8:LEU:HD23	2.10	0.52
2:B:7:ORN:O	2:B:8:LEU:CD2	2.58	0.51
1:A:126:LEU:HB3	2:B:6:VAL:HG11	1.93	0.51
2:B:1:VAL:CG1	2:B:2:ORN:CB	2.84	0.51
1:A:156:SER:H	1:A:191:GLN:NE2	2.08	0.49
1:A:126:LEU:HB2	2:B:6:VAL:HG12	1.95	0.48
2:B:1:VAL:N	2:B:9:DPN:C	2.75	0.47
1:A:166:SER:OG	2:B:3:LEU:HD22	2.15	0.46
2:B:9:DPN:HD2	2:B:10:PRO:CD	2.13	0.46
1:A:154:GLY:HA3	2:B:3:LEU:HD23	1.90	0.46
1:A:154:GLY:HA2	2:B:3:LEU:CD2	2.38	0.45
2:B:3:LEU:O	2:B:4:DPN:C	2.65	0.45
1:A:126:LEU:CB	2:B:6:VAL:CG1	2.88	0.44
2:B:2:ORN:C	2:B:3:LEU:CG	2.95	0.44
1:A:129:PRO:HD2	2:B:1:VAL:N	2.33	0.44
1:A:129:PRO:HD3	2:B:2:ORN:N	2.33	0.43
2:B:1:VAL:CB	2:B:10:PRO:O	2.66	0.43
1:A:100:GLY:O	2:B:6:VAL:HG22	2.18	0.43
2:B:1:VAL:HG12	2:B:2:ORN:CA	2.45	0.43
2:B:1:VAL:CG2	2:B:10:PRO:O	2.68	0.42
1:A:242:SER:OG	1:A:245:GLN:HG3	2.20	0.41
1:A:173:ASN:ND2	4:A:2121:HOH:O	2.52	0.41
2:B:5:PRO:C	2:B:6:VAL:HG23	2.41	0.40
2:B:9:DPN:CG	2:B:10:PRO:CD	2.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$267/269 \ (99\%)$	259 (97%)	8 (3%)	0	100	100
2	В	4/10 (40%)	2 (50%)	1 (25%)	1 (25%)	0	0
All	All	271/279 (97%)	261 (96%)	9 (3%)	1 (0%)	34	13

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	В	8	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	194/194 (100%)	190 (98%)	4 (2%)	53	22	
2	В	6/6 (100%)	6 (100%)	0	100	100	
All	All	200/200 (100%)	196 (98%)	4 (2%)	55	24	

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	170	ARG
1	A	221	SER
1	A	222	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN
1	A	109	GLN
1	A	155	ASN
1	A	173	ASN
1	A	182	GLN
1	A	191	GLN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	204	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Chain	Res	Link	B	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	ORN	В	2	2	6,7,8	0.76	0	2,7,9	0.21	0	
2	ORN	В	7	2	6,7,8	0.89	0	2,7,9	0.28	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ORN	В	2	2	-	0/5/6/8	-
2	ORN	В	7	2	-	2/5/6/8	_

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	7	ORN	N-CA-CB-CG
2	В	7	ORN	C-CA-CB-CG



There are no ring outliers.

2 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	2	ORN	18	0
2	В	7	ORN	8	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

