



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2023 – 03:13 AM EDT

PDB ID : 4TQO  
Title : The crystal structure of methanol dehydrogenase from *Methylococcus capsulatus* (Bath)  
Authors : Culpepper, M.A.; Rosenzweig, A.C.  
Deposited on : 2014-06-11  
Resolution : 2.57 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	573	90% 9% .
1	B	573	89% 9% ..
1	C	573	90% 10%
1	D	573	92% 7% .
1	E	573	92% 8% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	573	 90% 10%
1	G	573	 89% 11%
1	H	573	 91% 9%
2	I	72	 90% 8% .
2	J	72	 93% 6% .
2	K	72	 3% 96% . .
2	L	72	 94% . .
2	M	72	 82% 18%
2	N	72	 89% 10% .
2	O	72	 90% 10%
2	P	72	 90% 10%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 43059 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methanol dehydrogenase protein, large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	573	4491	2871	765	832	23	0	0	0
1	A	573	4491	2871	765	832	23	0	0	0
1	C	573	4491	2871	765	832	23	0	0	0
1	D	573	4490	2871	765	831	23	0	0	0
1	E	573	4491	2871	765	832	23	0	0	0
1	F	573	4491	2871	765	832	23	0	0	0
1	G	573	4491	2871	765	832	23	0	0	0
1	H	573	4491	2871	765	832	23	0	0	0

- Molecule 2 is a protein called Methanol dehydrogenase, small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	I	71	568	356	100	109	3	0	0	0
2	J	71	568	356	100	109	3	0	0	0
2	K	71	568	356	100	109	3	0	0	0
2	L	71	568	356	100	109	3	0	0	0
2	M	72	577	362	102	110	3	0	0	0
2	N	72	577	362	102	110	3	0	0	0

*Continued on next page...*



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			24	14	2	8		
4	A	1	Total	C	N	O	0	0
			24	14	2	8		
4	C	1	Total	C	N	O	0	0
			24	14	2	8		
4	D	1	Total	C	N	O	0	0
			24	14	2	8		
4	E	1	Total	C	N	O	0	0
			24	14	2	8		
4	F	1	Total	C	N	O	0	0
			24	14	2	8		
4	G	1	Total	C	N	O	0	0
			24	14	2	8		
4	H	1	Total	C	N	O	0	0
			24	14	2	8		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	276	Total	O	0	0
			276	276		
5	I	57	Total	O	0	0
			57	57		
5	A	288	Total	O	0	0
			288	288		
5	J	45	Total	O	0	0
			45	45		
5	C	237	Total	O	0	0
			237	237		
5	K	29	Total	O	0	0
			29	29		
5	D	247	Total	O	0	0
			247	247		
5	L	46	Total	O	0	0
			46	46		
5	E	241	Total	O	0	0
			241	241		
5	M	33	Total	O	0	0
			33	33		
5	F	251	Total	O	0	0
			251	251		
5	N	39	Total	O	0	0
			39	39		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	G	244	Total 244	O 244	0	0
5	O	35	Total 35	O 35	0	0
5	H	254	Total 254	O 254	0	0
5	P	30	Total 30	O 30	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methanol dehydrogenase protein, large subunit




- Molecule 1: Methanol dehydrogenase protein, large subunit



- Molecule 1: Methanol dehydrogenase protein, large subunit



- Molecule 1: Methanol dehydrogenase protein, large subunit









- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain E: 92% 8%




- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain F: 90% 10%




- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain G: 89% 11%



- Molecule 1: Methanol dehydrogenase protein, large subunit

Chain H: 91% 9%



- Molecule 2: Methanol dehydrogenase, small subunit

Chain I: 90% 8%



- Molecule 2: Methanol dehydrogenase, small subunit




- Molecule 2: Methanol dehydrogenase, small subunit




- Molecule 2: Methanol dehydrogenase, small subunit



- Molecule 2: Methanol dehydrogenase, small subunit




- Molecule 2: Methanol dehydrogenase, small subunit



- Molecule 2: Methanol dehydrogenase, small subunit



- Molecule 2: Methanol dehydrogenase, small subunit





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.49Å 210.29Å 231.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	105.15 – 2.57 105.14 – 2.57	Depositor EDS
% Data completeness (in resolution range)	99.9 (105.15-2.57) 100.0 (105.14-2.57)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.59 (at 2.58Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.4_1496)	Depositor
R, $R_{free}$	0.158 , 0.208 0.163 , 0.209	Depositor DCC
$R_{free}$ test set	9931 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	43059	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PQQ, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	2/4622 (0.0%)	0.49	0/6281
1	B	0.42	3/4622 (0.1%)	0.52	4/6281 (0.1%)
1	C	0.30	0/4622	0.47	0/6281
1	D	0.34	0/4621	0.49	0/6281
1	E	0.29	0/4622	0.48	0/6281
1	F	0.37	2/4622 (0.0%)	0.48	0/6281
1	G	0.27	0/4622	0.48	0/6281
1	H	0.28	0/4622	0.48	0/6281
2	I	0.34	0/583	0.42	0/785
2	J	0.28	0/583	0.43	0/785
2	K	0.25	0/583	0.41	0/785
2	L	0.27	0/583	0.42	0/785
2	M	0.31	0/592	0.54	0/796
2	N	0.33	0/592	0.53	1/796 (0.1%)
2	O	0.25	0/592	0.41	0/796
2	P	0.26	0/592	0.41	0/796
All	All	0.33	7/41675 (0.0%)	0.48	5/56572 (0.0%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	141	TYR	CD1-CE1	-6.59	1.29	1.39
1	B	141	TYR	CD2-CE2	-5.91	1.30	1.39
1	A	401	TYR	CD2-CE2	-5.77	1.30	1.39
1	A	401	TYR	CD1-CE1	-5.66	1.30	1.39
1	B	148	GLY	C-O	-5.50	1.14	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	523	ARG	NE-CZ-NH2	-7.38	116.61	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	523	ARG	NE-CZ-NH1	7.28	123.94	120.30
2	N	74	GLN	CA-CB-CG	6.28	127.21	113.40
1	B	144	ALA	C-N-CA	-5.72	110.28	122.30
1	B	145	GLY	N-CA-C	5.25	126.21	113.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4491	0	4320	47	0
1	B	4491	0	4320	40	0
1	C	4491	0	4320	38	0
1	D	4490	0	4320	31	0
1	E	4491	0	4320	31	0
1	F	4491	0	4320	37	0
1	G	4491	0	4320	42	0
1	H	4491	0	4320	36	0
2	I	568	0	545	6	0
2	J	568	0	545	3	0
2	K	568	0	545	2	0
2	L	568	0	545	3	0
2	M	577	0	558	12	0
2	N	577	0	558	9	0
2	O	577	0	558	5	0
2	P	577	0	558	5	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	24	0	3	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	24	0	3	3	0
4	C	24	0	3	1	0
4	D	24	0	3	1	0
4	E	24	0	3	0	0
4	F	24	0	3	2	0
4	G	24	0	3	2	0
4	H	24	0	3	2	0
5	A	288	0	0	3	0
5	B	276	0	0	5	1
5	C	237	0	0	9	0
5	D	247	0	0	7	1
5	E	241	0	0	3	0
5	F	251	0	0	8	0
5	G	244	0	0	9	0
5	H	254	0	0	9	1
5	I	57	0	0	3	1
5	J	45	0	0	1	1
5	K	29	0	0	1	0
5	L	46	0	0	2	0
5	M	33	0	0	3	0
5	N	39	0	0	2	1
5	O	35	0	0	0	0
5	P	30	0	0	0	0
All	All	43059	0	38996	337	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 337 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:400:GLU:OE1	1:A:401:TYR:CE2	1.71	1.41
1:A:400:GLU:OE1	1:A:401:TYR:CZ	1.79	1.33
1:A:400:GLU:OE1	1:A:401:TYR:CD2	2.09	1.03
2:M:35:TRP:O	5:M:706:HOH:O	1.80	0.98
1:A:400:GLU:OE1	1:A:401:TYR:CE1	2.26	0.89

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:705:HOH:O	5:J:702:HOH:O[4_444]	2.00	0.20
5:B:811:HOH:O	5:H:810:HOH:O[3_554]	2.04	0.16
5:D:810:HOH:O	5:N:707:HOH:O[3_654]	2.10	0.10

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	571/573 (100%)	543 (95%)	25 (4%)	3 (0%)	29	50
1	B	571/573 (100%)	539 (94%)	28 (5%)	4 (1%)	22	41
1	C	571/573 (100%)	537 (94%)	32 (6%)	2 (0%)	34	55
1	D	571/573 (100%)	536 (94%)	33 (6%)	2 (0%)	34	55
1	E	571/573 (100%)	541 (95%)	27 (5%)	3 (0%)	29	50
1	F	571/573 (100%)	533 (93%)	35 (6%)	3 (0%)	29	50
1	G	571/573 (100%)	538 (94%)	29 (5%)	4 (1%)	22	41
1	H	571/573 (100%)	536 (94%)	31 (5%)	4 (1%)	22	41
2	I	69/72 (96%)	69 (100%)	0	0	100	100
2	J	69/72 (96%)	69 (100%)	0	0	100	100
2	K	69/72 (96%)	69 (100%)	0	0	100	100
2	L	69/72 (96%)	68 (99%)	1 (1%)	0	100	100
2	M	70/72 (97%)	70 (100%)	0	0	100	100
2	N	70/72 (97%)	68 (97%)	2 (3%)	0	100	100
2	O	70/72 (97%)	70 (100%)	0	0	100	100
2	P	70/72 (97%)	70 (100%)	0	0	100	100
All	All	5124/5160 (99%)	4856 (95%)	243 (5%)	25 (0%)	29	50

5 of 25 Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	B	145	GLY
1	D	294	ASN
1	G	90	ASP
1	H	294	ASN
1	F	294	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	464/464 (100%)	461 (99%)	3 (1%)	86	94
1	B	464/464 (100%)	458 (99%)	6 (1%)	69	85
1	C	464/464 (100%)	459 (99%)	5 (1%)	73	88
1	D	464/464 (100%)	460 (99%)	4 (1%)	78	90
1	E	464/464 (100%)	459 (99%)	5 (1%)	73	88
1	F	464/464 (100%)	463 (100%)	1 (0%)	93	98
1	G	464/464 (100%)	463 (100%)	1 (0%)	93	98
1	H	464/464 (100%)	461 (99%)	3 (1%)	86	94
2	I	60/61 (98%)	60 (100%)	0	100	100
2	J	60/61 (98%)	60 (100%)	0	100	100
2	K	60/61 (98%)	60 (100%)	0	100	100
2	L	60/61 (98%)	60 (100%)	0	100	100
2	M	61/61 (100%)	61 (100%)	0	100	100
2	N	61/61 (100%)	61 (100%)	0	100	100
2	O	61/61 (100%)	61 (100%)	0	100	100
2	P	61/61 (100%)	61 (100%)	0	100	100
All	All	4196/4200 (100%)	4168 (99%)	28 (1%)	84	93

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	53	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	H	453	ARG
1	D	337	TYR
1	G	90	ASP
1	D	132	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	M	74	GLN
2	N	74	GLN
1	G	63	HIS
2	N	79	ASN
1	A	588	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PQQ	B	702	-	23,26,26	1.45	3 (13%)	29,40,40	1.90	10 (34%)
4	PQQ	E	702	-	23,26,26	1.40	3 (13%)	29,40,40	1.55	6 (20%)
4	PQQ	D	702	-	23,26,26	1.49	3 (13%)	29,40,40	1.54	7 (24%)
4	PQQ	C	702	-	23,26,26	1.42	3 (13%)	29,40,40	1.53	6 (20%)
4	PQQ	A	702	-	23,26,26	1.38	3 (13%)	29,40,40	1.70	7 (24%)
4	PQQ	G	702	-	23,26,26	1.41	3 (13%)	29,40,40	1.62	8 (27%)
4	PQQ	H	702	-	23,26,26	1.45	3 (13%)	29,40,40	1.55	6 (20%)
4	PQQ	F	702	-	23,26,26	1.44	3 (13%)	29,40,40	1.79	10 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PQQ	B	702	-	-	4/10/28/28	0/3/3/3
4	PQQ	E	702	-	-	0/10/28/28	0/3/3/3
4	PQQ	D	702	-	-	5/10/28/28	0/3/3/3
4	PQQ	C	702	-	-	0/10/28/28	0/3/3/3
4	PQQ	A	702	-	-	6/10/28/28	0/3/3/3
4	PQQ	G	702	-	-	1/10/28/28	0/3/3/3
4	PQQ	H	702	-	-	1/10/28/28	0/3/3/3
4	PQQ	F	702	-	-	4/10/28/28	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	702	PQQ	C9A-C1A	3.91	1.51	1.42
4	D	702	PQQ	C9A-C1A	3.90	1.51	1.42
4	E	702	PQQ	C9A-C1A	3.82	1.51	1.42
4	H	702	PQQ	C9A-C1A	3.80	1.51	1.42
4	F	702	PQQ	C9A-C1A	3.80	1.50	1.42

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	702	PQQ	O7B-C7X-C7	3.29	122.17	114.69
4	B	702	PQQ	C2-N1-C1A	3.14	110.36	103.90
4	F	702	PQQ	C9-C9A-C6A	-3.14	117.48	121.68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	702	PQQ	C9-C9A-C6A	-3.08	117.56	121.68
4	C	702	PQQ	C2-N1-C1A	3.07	110.20	103.90

There are no chirality outliers.

5 of 21 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	702	PQQ	N1-C2-C2X-O2A
4	A	702	PQQ	C9A-C9-C9X-O9B
4	D	702	PQQ	N1-C2-C2X-O2A
4	D	702	PQQ	N6-C7-C7X-O7A
4	D	702	PQQ	N6-C7-C7X-O7B


There are no ring outliers.


7 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	702	PQQ	3	0
4	D	702	PQQ	1	0
4	C	702	PQQ	1	0
4	A	702	PQQ	3	0
4	G	702	PQQ	2	0
4	H	702	PQQ	2	0
4	F	702	PQQ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	573/573 (100%)	-0.34	1 (0%) 95   95	15, 21, 32, 46	0
1	B	573/573 (100%)	-0.40	0 100   100	15, 22, 33, 49	0
1	C	573/573 (100%)	-0.19	3 (0%) 91   90	15, 25, 36, 53	0
1	D	573/573 (100%)	-0.38	0 100   100	16, 23, 34, 50	0
1	E	573/573 (100%)	-0.37	0 100   100	16, 23, 35, 48	0
1	F	573/573 (100%)	-0.40	0 100   100	15, 24, 34, 45	0
1	G	573/573 (100%)	-0.36	0 100   100	15, 24, 35, 50	0
1	H	573/573 (100%)	-0.22	2 (0%) 94   94	16, 24, 36, 52	0
2	I	71/72 (98%)	-0.17	0 100   100	20, 27, 45, 53	0
2	J	71/72 (98%)	-0.26	0 100   100	22, 27, 41, 48	0
2	K	71/72 (98%)	0.05	2 (2%) 53   49	28, 36, 46, 51	0
2	L	71/72 (98%)	-0.34	0 100   100	19, 31, 44, 55	0
2	M	72/72 (100%)	-0.15	0 100   100	24, 31, 48, 64	0
2	N	72/72 (100%)	-0.31	0 100   100	25, 32, 44, 48	0
2	O	72/72 (100%)	-0.18	0 100   100	25, 36, 48, 59	0
2	P	72/72 (100%)	0.16	0 100   100	23, 32, 47, 59	0
All	All	5156/5160 (99%)	-0.31	8 (0%) 95   95	15, 24, 38, 64	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	293	TRP	2.6
1	A	400	GLU	2.3
1	C	330	TRP	2.2
1	C	445	TRP	2.1
2	K	23	TYR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	H	701	1/1	0.67	0.16	97,97,97,97	0
4	PQQ	H	702	24/24	0.90	0.33	30,42,48,50	0
4	PQQ	E	702	24/24	0.91	0.28	27,40,43,45	0
4	PQQ	F	702	24/24	0.91	0.25	28,38,42,45	0
4	PQQ	B	702	24/24	0.91	0.29	27,36,40,43	0
4	PQQ	D	702	24/24	0.92	0.27	29,39,45,47	0
4	PQQ	C	702	24/24	0.92	0.27	26,36,42,44	0
4	PQQ	G	702	24/24	0.93	0.27	25,38,43,44	0
4	PQQ	A	702	24/24	0.94	0.25	25,33,36,38	0
3	CA	F	701	1/1	0.95	0.06	49,49,49,49	0
3	CA	D	701	1/1	0.95	0.07	44,44,44,44	0
3	CA	G	701	1/1	0.97	0.14	43,43,43,43	0
3	CA	C	701	1/1	0.97	0.14	46,46,46,46	0
3	CA	E	701	1/1	0.98	0.10	39,39,39,39	0
3	CA	A	701	1/1	0.98	0.15	40,40,40,40	0
3	CA	B	701	1/1	0.99	0.08	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around PQQ H 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)











**Electron density around PQQ F 702:**



$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around PQQ D 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around PQQ C 702:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around PQQ G 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.