



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 08:30 am BST

PDB ID : 5TRO
Title : 1.8 Angstrom Resolution Crystal Structure of Dimerization and Transpeptidase domains (residues 39-608) of Penicillin-Binding Protein 1 from *Staphylococcus aureus*.
Authors : Minasov, G.; Shuvalova, L.; Kiryukhina, O.; Dubrovskaya, I.; Grimshaw, S.; Kwon, K.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2016-10-26
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	573	
1	B	573	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8690 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Penicillin-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	505	4049	2565	699	770	15	0	9	0
1	B	506	4006	2539	690	762	15	0	3	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	36	SER	-	expression tag	UNP A0A0H2WVW5
A	37	ASN	-	expression tag	UNP A0A0H2WVW5
A	38	ALA	-	expression tag	UNP A0A0H2WVW5
B	36	SER	-	expression tag	UNP A0A0H2WVW5
B	37	ASN	-	expression tag	UNP A0A0H2WVW5
B	38	ALA	-	expression tag	UNP A0A0H2WVW5

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Cl 1	0	0
2	A	1	Total 1	Cl 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	364	Total 376	O 376	0	14
3	B	251	Total 257	O 257	0	6


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Penicillin-binding protein 1



- Molecule 1: Penicillin-binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	178.84Å 72.20Å 86.36Å 90.00° 103.18° 90.00°	Depositor
Resolution (Å)	29.59 – 1.80 29.02 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.59-1.80) 99.3 (29.02-1.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.179 , 0.209 0.193 , 0.223	Depositor DCC
R_{free} test set	4950 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtrriage
Anisotropy	0.276	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8690	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/4121	0.68	0/5511
1	B	0.44	0/4075	0.65	0/5453
All	All	0.46	0/8196	0.66	0/10964

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4049	0	4037	12	0
1	B	4006	0	3993	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	376	0	0	3	0
3	B	257	0	0	0	0
All	All	8690	0	8030	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261[B]:GLU:CD	3:A:801:HOH:O	2.14	0.85
1:A:261[B]:GLU:OE2	3:A:801:HOH:O	1.97	0.82
1:A:261[B]:GLU:OE1	3:A:801:HOH:O	2.12	0.65
1:A:119:MSE:HE1	1:A:127:ARG:NE	2.13	0.62
1:B:197:LEU:HD23	1:B:198:GLY:N	2.17	0.60
1:A:150:LYS:HA	1:A:163:LEU:HD13	1.85	0.58
1:A:352:ASN:O	1:A:353:ARG:HB2	2.07	0.53
1:B:378:ASP:OD1	1:B:419:LYS:NZ	2.40	0.53
1:A:235:LYS:HG2	1:A:236:ARG:H	1.74	0.51
1:A:143:THR:HG22	1:A:144:ASN:H	1.76	0.50
1:B:325:ILE:HD13	1:B:484:LYS:HB3	1.95	0.48
1:B:502:ASN:O	1:B:575:LYS:HD2	2.15	0.47
1:B:271:VAL:HG11	1:B:435:LEU:HD11	2.01	0.42
1:A:63:PRO:HB2	1:A:209:GLY:HA3	2.00	0.42
1:A:542:ALA:HA	1:A:543:PRO:C	2.40	0.42
1:A:343:ILE:HD11	1:A:415:GLU:C	2.41	0.42
1:A:243:THR:OG1	1:A:452:PHE:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/573 (89%)	496 (97%)	14 (3%)	0	100	100
1	B	505/573 (88%)	493 (98%)	12 (2%)	0	100	100
All	All	1015/1146 (89%)	989 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/467 (93%)	432 (100%)	2 (0%)	88	87
1	B	429/467 (92%)	425 (99%)	4 (1%)	78	75
All	All	863/934 (92%)	857 (99%)	6 (1%)	84	81

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	LYS
1	A	140	ARG
1	B	70	ASP
1	B	92	LYS
1	B	98	LYS
1	B	358	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/573 (85%)	0.36	46 (9%) 8 6	21, 34, 77, 101	0
1	B	491/573 (85%)	0.51	54 (10%) 5 4	25, 44, 78, 94	0
All	All	981/1146 (85%)	0.44	100 (10%) 6 5	21, 39, 78, 101	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	ALA	6.1
1	B	235	LYS	5.7
1	B	347	ARG	5.6
1	B	210	SER	5.1
1	A	346	SER	4.7
1	B	340	HIS	4.5
1	A	525	GLY	4.5
1	A	345	GLY	4.4
1	B	353	ARG	4.4
1	B	96	ASN	4.4
1	A	116	VAL	4.3
1	B	148	GLN	4.2
1	B	348	ILE	4.2
1	A	96	ASN	4.0
1	B	526	GLY	4.0
1	A	98	LYS	3.9
1	A	506	ASP	3.9
1	B	338	SER	3.9
1	B	342	ASP	3.8
1	A	354	VAL	3.7
1	A	235	LYS	3.7
1	B	319	TYR	3.7
1	B	354	VAL	3.6
1	B	315	THR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	335	LYS	3.5
1	A	118	ASN	3.4
1	A	61	GLN	3.4
1	B	525	GLY	3.4
1	A	590	SER	3.4
1	A	140	ARG	3.3
1	A	133	ALA	3.3
1	B	98	LYS	3.3
1	A	91	LYS	3.2
1	B	61	GLN	3.2
1	B	546	ASN	3.2
1	A	552	TYR	3.2
1	A	120	LYS	3.2
1	A	151	LEU	3.2
1	B	337	LYS	3.2
1	B	155	LYS	3.1
1	B	435	LEU	3.1
1	B	358	GLU	3.0
1	A	92	LYS	3.0
1	B	545	LYS	3.0
1	A	132	LYS	2.9
1	A	104	VAL	2.9
1	A	99	LYS	2.9
1	A	340	HIS	2.9
1	B	91	LYS	2.9
1	A	347	ARG	2.9
1	B	367	TYR	2.9
1	A	342	ASP	2.8
1	B	349	SER	2.8
1	B	316	PHE	2.8
1	A	148	GLN	2.8
1	B	118	ASN	2.8
1	A	101	ARG	2.8
1	B	333	ASP	2.8
1	B	345	GLY	2.7
1	B	589	LYS	2.7
1	A	319	TYR	2.7
1	B	115	THR	2.7
1	A	149	ASP	2.6
1	A	589	LYS	2.6
1	A	526	GLY	2.6
1	B	149	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	346	SER	2.6
1	B	416	LEU	2.5
1	A	97	SER	2.5
1	B	99	LYS	2.5
1	A	270	ALA	2.5
1	A	529	LYS	2.4
1	A	111	LYS	2.4
1	B	312	PRO	2.4
1	A	435	LEU	2.4
1	B	552	TYR	2.3
1	A	353	ARG	2.3
1	B	90	ASP	2.3
1	B	180	LEU	2.3
1	A	343	ILE	2.3
1	A	107	LYS	2.3
1	B	271	VAL	2.2
1	B	343	ILE	2.2
1	B	317	LYS	2.2
1	B	508	TYR	2.2
1	B	524	GLY	2.2
1	B	95	ALA	2.2
1	A	523	ASN	2.2
1	A	112	LYS	2.2
1	B	132	LYS	2.2
1	A	524	GLY	2.1
1	B	92	LYS	2.1
1	A	115	THR	2.1
1	A	123	GLU	2.1
1	B	437	ALA	2.1
1	B	153	ILE	2.1
1	A	139	GLY	2.1
1	B	341	ARG	2.0
1	B	133	ALA	2.0
1	B	515	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	701	1/1	0.96	0.11	40,40,40,40	0
2	CL	B	701	1/1	0.98	0.06	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.