



Full wwPDB X-ray Structure Validation Report i

Nov 13, 2023 – 05:54 PM EST

PDB ID : 8TSA
Title : Human PI3K p85alpha/p110alpha H1047R bound to compound 2
Authors : Holliday, M.; Tang, Y.; Bulku, A.; Wilbur, J.; Fraser, J.; Valverde, R.
Deposited on : 2023-08-11
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 21101 atoms, of which 10548 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	1005	16447	5251	8237	1407	1484	68	63	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	expression tag	UNP P42336
A	-8	SER	-	expression tag	UNP P42336
A	-7	PRO	-	expression tag	UNP P42336
A	-6	GLY	-	expression tag	UNP P42336
A	-5	ILE	-	expression tag	UNP P42336
A	-4	SER	-	expression tag	UNP P42336
A	-3	GLY	-	expression tag	UNP P42336
A	-2	GLY	-	expression tag	UNP P42336
A	-1	GLY	-	expression tag	UNP P42336
A	0	GLY	-	expression tag	UNP P42336
A	1	GLY	-	expression tag	UNP P42336
A	1047	ARG	HIS	engineered mutation	UNP P42336


- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	272	4607	1451	2293	412	444	7	127	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	316	GLY	-	expression tag	UNP P27986
B	317	PRO	-	expression tag	UNP P27986

- Molecule 3 is 5-(3-bromo-5-fluorobenzamido)-N-methyl-6-(2-methylanilino)pyridine-3-carboxamide (three-letter code: UIW) (formula: $C_{21}H_{18}BrFN_4O_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	Br	C	F	H	N	O	0	0
			47	1	21	1	18	4	2		


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



- Molecule 2: Phosphatidylinositol 3-kinase regulatory subunit alpha



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.33Å 120.38Å 190.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	95.49 – 2.51 95.49 – 2.51	Depositor EDS
% Data completeness (in resolution range)	99.7 (95.49-2.51) 99.7 (95.49-2.51)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.06 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R , R_{free}	0.241 , 0.276 0.236 , 0.273	Depositor DCC
R_{free} test set	1767 reflections (2.58%)	wwPDB-VP
Wilson B-factor (Å ²)	88.4	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 69.7	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	21101	wwPDB-VP
Average B, all atoms (Å ²)	123.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UIW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/8392	0.47	0/11339
2	B	0.24	0/2354	0.48	0/3153
All	All	0.24	0/10746	0.47	0/14492

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8210	8237	8247	16	0
2	B	2314	2293	2292	5	0
3	A	29	18	0	1	0
All	All	10553	10548	10539	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:992:ARG:NH1	1:A:1027:ALA:O	2.26	0.67
2:B:356:LEU:HD23	2:B:357:VAL:N	2.12	0.65
1:A:913:ILE:HG22	1:A:913:ILE:O	2.04	0.58
1:A:398:ARG:NH1	1:A:434:ASP:OD1	2.37	0.55
1:A:198:VAL:HG22	1:A:200:PRO:HD2	1.90	0.52
1:A:480:PHE:O	1:A:481:SER:OG	2.18	0.52
1:A:543:ILE:HG23	1:A:543:ILE:O	2.11	0.51
1:A:199:SER:O	1:A:201:ASN:N	2.46	0.49
2:B:390:TYR:N	2:B:398:PHE:O	2.45	0.49
2:B:354:THR:HG22	2:B:426:TYR:HB2	1.97	0.46
1:A:807:LEU:HD12	1:A:846:GLY:HA3	1.99	0.45
1:A:787:ASP:OD1	1:A:788:ILE:N	2.49	0.45
3:A:1101:UIW:N18	3:A:1101:UIW:C10	2.81	0.43
1:A:142:ASN:O	1:A:145:ASN:ND2	2.51	0.43
1:A:121:ILE:HG22	1:A:688:LEU:HB3	2.01	0.42
1:A:208:THR:C	1:A:209:LEU:HD12	2.40	0.42
2:B:562:ARG:O	2:B:566:ILE:HD13	2.20	0.41
1:A:861:GLN:HA	1:A:876:THR:HG21	2.03	0.41
2:B:371:THR:HG22	2:B:380:LEU:HG	2.03	0.41
1:A:873:ASN:O	1:A:876:THR:HG22	2.22	0.40
1:A:139:PHE:CZ	1:A:143:ILE:HD13	2.57	0.40
1:A:628:LEU:C	1:A:628:LEU:HD13	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	995/1060 (94%)	957 (96%)	38 (4%)	0	100 100
2	B	270/300 (90%)	257 (95%)	13 (5%)	0	100 100
All	All	1265/1360 (93%)	1214 (96%)	51 (4%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	922/960 (96%)	915 (99%)	7 (1%)	81 93
2	B	253/277 (91%)	251 (99%)	2 (1%)	81 93
All	All	1175/1237 (95%)	1166 (99%)	9 (1%)	81 93

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	ARG
1	A	115	ARG
1	A	187	LYS
1	A	282	MET
1	A	342	THR
1	A	523	LEU
1	A	1047	ARG
2	B	396	LEU
2	B	413	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	UIW	A	1101	-	30,31,31	2.36	6 (20%)	41,43,43	1.36	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UIW	A	1101	-	-	4/18/18/18	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	UIW	C03-N02	9.47	1.45	1.33
3	A	1101	UIW	C08-N09	5.48	1.46	1.36
3	A	1101	UIW	C19-N18	3.64	1.45	1.35
3	A	1101	UIW	C10-N09	2.39	1.46	1.39
3	A	1101	UIW	O04-C03	-2.22	1.18	1.23
3	A	1101	UIW	O20-C19	-2.19	1.18	1.23

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1101	UIW	C10-N09-C08	-2.98	122.18	129.63
3	A	1101	UIW	C17-N18-C19	-2.87	119.06	126.93
3	A	1101	UIW	C25-C23-C22	-2.65	120.17	123.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	1101	UIW	C14-C15-C10	2.56	119.85	117.44
3	A	1101	UIW	C05-C03-N02	2.51	120.23	116.89
3	A	1101	UIW	C06-N07-C08	2.32	120.70	116.05
3	A	1101	UIW	C21-C19-N18	2.07	120.47	115.92

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1101	UIW	C17-C08-N09-C10
3	A	1101	UIW	N07-C08-N09-C10
3	A	1101	UIW	C29-C17-N18-C19
3	A	1101	UIW	C08-C17-N18-C19

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1101	UIW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1005/1060 (94%)	0.79	86 (8%) 10 10	67, 101, 145, 186	5 (0%)
2	B	272/300 (90%)	1.03	41 (15%) 2 1	82, 136, 190, 214	9 (3%)
All	All	1277/1360 (93%)	0.84	127 (9%) 7 7	67, 105, 170, 214	14 (1%)

All (127) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	500	VAL	6.4
1	A	367	LEU	5.3
1	A	45	ILE	5.2
2	B	370	LEU	5.2
2	B	392	PHE	4.6
2	B	353	GLY	4.6
2	B	355	PHE	4.4
2	B	333	TRP	4.3
2	B	589	VAL	4.3
1	A	181	ILE	3.9
1	A	302	PHE	3.8
2	B	447	LYS	3.6
1	A	34	LEU	3.6
2	B	381	ILE	3.5
2	B	449	LEU	3.5
1	A	989	LEU	3.5
1	A	777	ARG	3.4
1	A	37	LEU	3.4
2	B	334	TYR	3.3
1	A	880	TRP	3.3
1	A	738	GLN	3.2
1	A	429	LEU	3.2
1	A	1016	PHE	3.1
2	B	338	ILE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	179	LYS	3.1
2	B	357	VAL	3.1
2	B	379	LYS	3.1
1	A	734	PHE	3.1
2	B	427	PRO	3.0
1	A	11	TRP	3.0
1	A	1019	ILE	3.0
2	B	586	GLN	3.0
1	A	415	ALA	2.9
2	B	363	LYS	2.9
1	A	731	GLN	2.9
1	A	534	ILE	2.9
1	A	479	TRP	2.9
1	A	881	LEU	2.8
2	B	347	LEU	2.8
1	A	5	PRO	2.8
1	A	715	LEU	2.8
2	B	588	GLY	2.8
1	A	230	ARG	2.8
1	A	49	LEU	2.8
1	A	531	LEU	2.8
2	B	475	GLN	2.8
1	A	913	ILE	2.7
2	B	430	LYS	2.7
2	B	383	ILE	2.7
2	B	401	VAL	2.7
2	B	422	VAL	2.7
1	A	310	ARG	2.7
2	B	425	LEU	2.7
1	A	278	MET	2.7
1	A	281	ARG	2.7
1	A	528	LYS	2.7
1	A	477	PHE	2.7
1	A	132	LYS	2.7
1	A	443	LEU	2.6
1	A	153	LEU	2.6
1	A	540	LEU	2.6
1	A	473	LEU	2.6
1	A	403	CYS	2.6
2	B	424	LEU	2.6
2	B	437	VAL	2.6
1	A	3	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	663	ILE	2.5
1	A	548	LYS	2.5
1	A	404	LEU	2.5
1	A	445	LEU	2.5
2	B	356	LEU	2.5
1	A	309	ARG	2.5
2	B	420	LEU	2.5
1	A	325	LYS	2.5
2	B	538	ILE	2.5
2	B	394	ASP	2.4
1	A	339	LEU	2.4
2	B	549	LEU	2.4
1	A	475	LEU	2.4
2	B	416	TYR	2.4
1	A	498	TRP	2.4
1	A	728	GLN	2.4
1	A	568	LEU	2.4
1	A	866	LEU	2.3
2	B	433	GLN	2.3
1	A	409	VAL	2.3
1	A	564	ILE	2.3
1	A	781	LEU	2.3
1	A	386	TRP	2.3
1	A	718	ILE	2.3
1	A	328	TRP	2.3
1	A	771	ILE	2.3
1	A	725	ASP	2.3
2	B	571	ILE	2.2
1	A	455	LEU	2.2
1	A	1010	MET	2.2
1	A	190	ILE	2.2
1	A	200	PRO	2.2
1	A	551	LEU	2.2
1	A	501	SER	2.2
1	A	58	LEU	2.2
1	A	396	LEU	2.2
2	B	328	LEU	2.2
1	A	50	PHE	2.2
2	B	587	LYS	2.2
1	A	436	LEU	2.2
1	A	452	LEU	2.2
2	B	518	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1004	MET	2.1
2	B	428	VAL	2.1
1	A	449	PRO	2.1
1	A	1022	ILE	2.1
2	B	415	GLN	2.1
1	A	829	LEU	2.1
2	B	408	TYR	2.1
1	A	277	ILE	2.1
1	A	422	LEU	2.1
1	A	723	LYS	2.1
1	A	243	VAL	2.1
1	A	346	VAL	2.1
1	A	117	ILE	2.1
1	A	1039	PHE	2.0
1	A	407	CYS	2.0
1	A	735	LEU	2.0
2	B	404	LEU	2.0
1	A	929	LEU	2.0
1	A	621	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	UIW	A	1101	29/29	0.96	0.25	83,111,158,165	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.