



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2023 – 12:01 AM EDT

PDB ID : 3TV8  
Title : Pharmacological Chaperoning in Human alpha-Galactosidase  
Authors : Rogich, J.J.; Guce, A.I.; Clark, N.E.; Garman, S.C.  
Deposited on : 2011-09-19  
Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	
1	B	404	
2	C	3	
3	D	4	

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-galactosidase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	390	3119	1987	534	572	26	0	0	0
1	B	391	3136	1997	537	575	27	0	1	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	170	ALA	ASP	engineered mutation	UNP P06280
A	430	HIS	-	expression tag	UNP P06280
A	431	HIS	-	expression tag	UNP P06280
A	432	HIS	-	expression tag	UNP P06280
A	433	HIS	-	expression tag	UNP P06280
A	434	HIS	-	expression tag	UNP P06280
A	435	HIS	-	expression tag	UNP P06280
B	170	ALA	ASP	engineered mutation	UNP P06280
B	430	HIS	-	expression tag	UNP P06280
B	431	HIS	-	expression tag	UNP P06280
B	432	HIS	-	expression tag	UNP P06280
B	433	HIS	-	expression tag	UNP P06280
B	434	HIS	-	expression tag	UNP P06280
B	435	HIS	-	expression tag	UNP P06280

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	4	50	28	2	20	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is NONAETHYLENE GLYCOL (three-letter code: 2PE) (formula: C<sub>18</sub>H<sub>38</sub>O<sub>10</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is (2R,3S,4R,5S)-2-(hydroxymethyl)piperidine-3,4,5-triol (three-letter code: DGJ) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			11	6	1	4		
7	B	1	Total	C	N	O	0	0
			11	6	1	4		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	36	Total	O	0	0
			36	36		
8	B	28	Total	O	0	0
			28	28		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-galactosidase A



- Molecule 1: Alpha-galactosidase A



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.47Å 90.47Å 216.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.54 – 2.64 44.54 – 2.64	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.54-2.64) 99.7 (44.54-2.64)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 2.65Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.201 , 0.239 0.199 , 0.233	Depositor DCC
$R_{free}$ test set	1582 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 36.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6516	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, 2PE, NAG, SO4, DGJ, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.43	4/3206 (0.1%)	0.59	3/4354 (0.1%)
1	B	0.47	5/3223 (0.2%)	0.58	3/4376 (0.1%)
All	All	0.45	9/6429 (0.1%)	0.58	6/8730 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	418	GLU	CD-OE1	9.23	1.35	1.25
1	B	241	SER	CB-OG	8.22	1.52	1.42
1	B	418	GLU	CD-OE2	6.58	1.32	1.25
1	B	218	GLU	CD-OE1	-5.77	1.19	1.25
1	A	218	GLU	CD-OE1	-5.58	1.19	1.25
1	A	218	GLU	CD-OE2	-5.50	1.19	1.25
1	A	341	GLU	CD-OE1	-5.30	1.19	1.25
1	B	218	GLU	CD-OE2	-5.22	1.20	1.25
1	A	241	SER	CB-OG	5.17	1.49	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	GLU	OE1-CD-OE2	-7.09	114.80	123.30
1	B	218	GLU	OE1-CD-OE2	-7.04	114.85	123.30
1	A	341	GLU	OE1-CD-OE2	-6.81	115.13	123.30
1	B	341	GLU	OE1-CD-OE2	-6.20	115.86	123.30
1	A	136	ASP	CB-CG-OD1	5.37	123.13	118.30
1	B	136	ASP	CB-CG-OD1	5.24	123.01	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3119	0	2982	26	0
1	B	3136	0	2998	31	0
2	C	39	0	34	0	0
3	D	50	0	43	0	0
4	A	28	0	26	0	0
4	B	28	0	26	0	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
6	A	11	0	12	0	0
6	B	4	0	4	0	0
7	A	11	0	13	0	0
7	B	11	0	13	0	0
8	A	36	0	0	0	0
8	B	28	0	0	0	0
All	All	6516	0	6151	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:ARG:HH11	1:A:158:THR:HG23	1.10	1.14
1:A:403:LEU:HD11	1:A:415:LEU:HD13	1.57	0.86
1:A:105:ARG:NH1	1:A:158:THR:HG23	1.90	0.86
1:A:99:GLN:HE21	1:A:99:GLN:HA	1.41	0.86
1:B:99:GLN:HE21	1:B:99:GLN:HA	1.42	0.82
1:A:105:ARG:HH11	1:A:158:THR:CG2	1.92	0.81
1:A:369:VAL:HA	1:A:372:LEU:HD12	1.65	0.76
1:A:351:VAL:HG21	1:A:372:LEU:HD22	1.68	0.75
1:A:366:THR:HG22	1:A:404:ARG:HD3	1.71	0.72
1:B:403:LEU:HD21	1:B:415:LEU:HD13	1.75	0.68
1:A:99:GLN:HE21	1:A:99:GLN:CA	2.07	0.67
1:A:403:LEU:HD11	1:A:415:LEU:CD1	2.25	0.67
1:B:99:GLN:HE21	1:B:99:GLN:CA	2.09	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:THR:CG2	1:A:404:ARG:HD3	2.27	0.65
1:B:351:VAL:HG21	1:B:372:LEU:CD2	2.26	0.65
1:B:228:ASN:HB3	1:B:245:TRP:CH2	2.32	0.64
1:B:351:VAL:HG21	1:B:372:LEU:HD23	1.80	0.64
1:A:228:ASN:HB3	1:A:245:TRP:CH2	2.33	0.63
1:B:236:TRP:CE2	1:B:240:LYS:HD2	2.39	0.57
1:B:177:LEU:HD12	1:B:211:PHE:HB3	1.89	0.55
1:B:369:VAL:HA	1:B:372:LEU:HD13	1.89	0.55
1:B:236:TRP:NE1	1:B:240:LYS:HD2	2.23	0.54
1:A:420:THR:O	1:A:421:MET:HG2	2.07	0.53
1:B:369:VAL:HA	1:B:372:LEU:CD1	2.39	0.53
1:B:137:VAL:HG12	1:B:171:GLY:HA2	1.90	0.52
1:B:145:PHE:HB3	1:B:146:PRO:HD2	1.93	0.50
1:A:145:PHE:HB3	1:A:146:PRO:HD2	1.93	0.50
1:A:101:ASP:HB3	1:A:107:GLN:OE1	2.13	0.48
1:A:386:GLN:HA	1:A:415:LEU:HD23	1.94	0.48
1:B:177:LEU:HD11	1:B:212:GLN:HG2	1.96	0.47
1:A:137:VAL:HG12	1:A:171:GLY:HA2	1.97	0.47
1:B:418:GLU:OE2	1:B:418:GLU:HA	2.15	0.47
1:B:99:GLN:CA	1:B:99:GLN:NE2	2.78	0.46
1:A:99:GLN:CA	1:A:99:GLN:NE2	2.77	0.46
1:B:351:VAL:CG2	1:B:372:LEU:HD23	2.45	0.45
1:B:75:LEU:HD11	1:B:301:ARG:HG2	1.97	0.45
1:B:101:ASP:HB3	1:B:107:GLN:OE1	2.16	0.45
1:A:418:GLU:OE2	1:A:418:GLU:HA	2.18	0.44
1:A:69:PHE:CZ	1:A:91:ILE:HG23	2.53	0.43
1:B:180:LEU:HD23	1:B:208:MET:HG3	1.98	0.43
1:B:335:ASP:O	1:B:335:ASP:OD1	2.37	0.43
1:B:324:LEU:HD13	1:B:326:LYS:HG3	2.01	0.43
1:A:75:LEU:HD11	1:A:301:ARG:HG2	2.00	0.42
1:A:214:PRO:O	1:A:252:ARG:NH2	2.53	0.42
1:B:351:VAL:HG21	1:B:372:LEU:HD22	2.00	0.42
1:A:124:VAL:HG13	1:A:129:LEU:HB2	2.02	0.42
1:B:369:VAL:HG22	1:B:401:SER:O	2.19	0.42
1:B:366:THR:CG2	1:B:404:ARG:HG2	2.49	0.42
1:B:124:VAL:HG13	1:B:129:LEU:HB2	2.02	0.42
1:B:69:PHE:CZ	1:B:91:ILE:HG23	2.54	0.42
1:B:403:LEU:HD21	1:B:415:LEU:CD1	2.48	0.42
1:B:89:LEU:O	1:B:131:LEU:HD12	2.21	0.41
1:B:366:THR:HG22	1:B:404:ARG:HG2	2.03	0.40
1:A:204:TRP:HB3	1:A:205:PRO:HD3	2.03	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:386:GLN:HA	1:B:415:LEU:HD23	2.03	0.40
1:A:89:LEU:O	1:A:131:LEU:HD12	2.22	0.40
1:A:335:ASP:OD1	1:A:363:ARG:NH1	2.48	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	388/404 (96%)	379 (98%)	8 (2%)	1 (0%)	41 56
1	B	390/404 (96%)	379 (97%)	10 (3%)	1 (0%)	41 56
All	All	778/808 (96%)	758 (97%)	18 (2%)	2 (0%)	41 56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	ASP
1	B	266	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	330/344 (96%)	318 (96%)	12 (4%)	35 52

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	332/344 (96%)	322 (97%)	10 (3%)	41	59
All	All	662/688 (96%)	640 (97%)	22 (3%)	38	55

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	A	107	GLN
1	A	154	ILE
1	A	158	THR
1	A	241	SER
1	A	256	VAL
1	A	336	ASN
1	A	345	SER
1	A	403	LEU
1	A	404	ARG
1	A	418	GLU
1	A	421	MET
1	B	99	GLN
1	B	107	GLN
1	B	154	ILE
1	B	158	THR
1	B	196	ARG
1	B	213	LYS
1	B	256	VAL
1	B	324	LEU
1	B	345	SER
1	B	418	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	A	228	ASN
1	A	336	ASN
1	B	99	GLN
1	B	179	ASN
1	B	212	GLN
1	B	228	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	1	2,1	14,14,15	0.48	0	17,19,21	0.73	0
2	NAG	C	2	2	14,14,15	0.47	0	17,19,21	1.18	2 (11%)
2	BMA	C	3	2	11,11,12	0.43	0	15,15,17	0.93	1 (6%)
3	NAG	D	1	3,1	14,14,15	0.52	0	17,19,21	1.38	3 (17%)
3	NAG	D	2	3	14,14,15	0.50	0	17,19,21	0.79	1 (5%)
3	BMA	D	3	3	11,11,12	0.48	0	15,15,17	1.35	2 (13%)
3	MAN	D	4	3	11,11,12	0.60	0	15,15,17	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	D	4	3	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3	BMA	O3-C3-C2	3.41	116.52	109.99
3	D	1	NAG	C4-C3-C2	-3.09	106.49	111.02
3	D	1	NAG	O5-C1-C2	-3.00	106.55	111.29
2	C	3	BMA	O5-C5-C6	2.55	111.20	107.20
3	D	1	NAG	C1-C2-N2	2.52	114.78	110.49
3	D	2	NAG	C1-O5-C5	2.14	115.10	112.19
3	D	3	BMA	O5-C5-C6	2.10	110.49	107.20
2	C	2	NAG	C3-C4-C5	-2.09	106.51	110.24
2	C	2	NAG	C1-O5-C5	2.08	115.02	112.19

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	BMA	O5-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
2	C	3	BMA	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
3	D	4	MAN	O5-C5-C6-O6
3	D	1	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
7	DGJ	B	802	-	11,11,11	0.66	0	13,15,15	1.46	1 (7%)
5	SO4	B	7	-	4,4,4	0.15	0	6,6,6	0.08	0
6	2PE	B	12	-	3,3,27	0.45	0	2,2,26	0.14	0
4	NAG	A	715	1	14,14,15	0.58	0	17,19,21	1.17	2 (11%)
6	2PE	A	9	-	6,6,27	0.48	0	5,5,26	0.19	0
6	2PE	A	11	-	3,3,27	0.45	0	2,2,26	0.20	0
7	DGJ	A	801	-	11,11,11	0.75	0	13,15,15	1.56	1 (7%)
4	NAG	B	715	1	14,14,15	0.55	0	17,19,21	0.94	1 (5%)
4	NAG	A	639	1	14,14,15	0.49	0	17,19,21	1.04	2 (11%)
5	SO4	A	8	-	4,4,4	0.14	0	6,6,6	0.09	0
4	NAG	B	639	1	14,14,15	0.52	0	17,19,21	0.91	0
5	SO4	B	6	-	4,4,4	0.17	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	DGJ	B	802	-	-	0/2/19/19	0/1/1/1
4	NAG	A	715	1	-	2/6/23/26	0/1/1/1
6	2PE	A	9	-	-	2/4/4/25	-
6	2PE	A	11	-	-	0/1/1/25	-
7	DGJ	A	801	-	-	0/2/19/19	0/1/1/1
4	NAG	B	715	1	-	2/6/23/26	0/1/1/1
4	NAG	A	639	1	-	0/6/23/26	0/1/1/1
4	NAG	B	639	1	-	0/6/23/26	0/1/1/1
6	2PE	B	12	-	-	1/1/1/25	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	801	DGJ	C1-N5-C5	5.30	121.07	109.61
7	B	802	DGJ	C1-N5-C5	4.77	119.93	109.61
4	A	715	NAG	C1-O5-C5	2.48	115.55	112.19
4	B	715	NAG	C1-O5-C5	2.18	115.14	112.19
4	A	639	NAG	O5-C5-C6	2.12	110.53	107.20
4	A	639	NAG	C1-O5-C5	2.10	115.04	112.19
4	A	715	NAG	O5-C5-C6	2.06	110.43	107.20

There are no chirality outliers.


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	715	NAG	C4-C5-C6-O6
4	A	715	NAG	O5-C5-C6-O6
6	A	9	2PE	O13-C14-C15-O16
6	B	12	2PE	O13-C14-C15-O16
4	B	715	NAG	C4-C5-C6-O6
6	A	9	2PE	C15-C14-O13-C12
4	B	715	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	390/404 (96%)	-0.14	4 (1%) 82 81	17, 31, 55, 73	0
1	B	391/404 (96%)	-0.04	2 (0%) 91 90	18, 36, 60, 79	0
All	All	781/808 (96%)	-0.09	6 (0%) 86 85	17, 33, 58, 79	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	102	SER	3.7
1	A	175	ASP	3.0
1	A	182	ASP	2.3
1	A	211	PHE	2.3
1	B	161	ASP	2.2
1	A	183	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BMA	C	3	11/12	0.66	0.26	62,73,80,83	0
3	MAN	D	4	11/12	0.76	0.24	84,87,89,91	0
3	BMA	D	3	11/12	0.80	0.34	67,73,80,84	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	D	1	14/15	0.90	0.26	39,53,65,65	0
3	NAG	D	2	14/15	0.91	0.30	42,62,66,68	0
2	NAG	C	2	14/15	0.94	0.14	43,51,64,65	0
2	NAG	C	1	14/15	0.95	0.15	8,31,36,38	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	B	639	14/15	0.63	0.28	83,89,91,92	0
6	2PE	B	12	4/28	0.77	0.30	75,76,78,80	0
6	2PE	A	9	7/28	0.78	0.31	49,59,64,66	0
4	NAG	B	715	14/15	0.78	0.25	81,87,88,88	0
4	NAG	A	639	14/15	0.83	0.37	77,81,82,82	0
6	2PE	A	11	4/28	0.90	0.26	53,54,61,63	0
4	NAG	A	715	14/15	0.90	0.19	42,53,57,58	0
5	SO4	B	7	5/5	0.92	0.36	88,89,92,93	0
7	DGJ	A	801	11/11	0.94	0.25	22,33,40,43	0
7	DGJ	B	802	11/11	0.96	0.15	23,33,36,37	0
5	SO4	A	8	5/5	0.98	0.15	38,40,44,46	0
5	SO4	B	6	5/5	0.98	0.23	53,56,63,66	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.