



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 24, 2023 – 12:12 AM EDT

PDB ID : 5TVG  
Title : Crystal structure of an alpha,alpha-trehalose-phosphate synthase (UDP-forming) from Burkholderia vietnamiensis  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2016-11-08  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	 89% 8% 3% 2%
1	B	481	 88% 6% 6% 2%
1	C	481	 87% 8% 5% 5%
1	D	481	 86% 9% 5% 0%
1	E	481	 87% 5% 7% 6%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	481	<p>%</p>
1	G	481	<p>11%</p>
1	H	481	<p>11%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	503	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 28914 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha,alpha-trehalose-phosphate synthase (UDP-forming).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	468	3687	2338	676	657	16	0	5	0
1	B	454	3505	2232	627	630	16	0	0	0
1	C	458	3542	2248	639	639	16	0	1	0
1	D	456	3547	2252	645	634	16	0	0	0
1	E	447	3423	2176	616	615	16	0	1	0
1	F	456	3537	2250	636	635	16	0	1	0
1	G	439	3291	2089	591	595	16	0	1	0
1	H	443	3311	2101	590	607	13	0	1	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP A4JGS8
A	-6	ALA	-	expression tag	UNP A4JGS8
A	-5	HIS	-	expression tag	UNP A4JGS8
A	-4	HIS	-	expression tag	UNP A4JGS8
A	-3	HIS	-	expression tag	UNP A4JGS8
A	-2	HIS	-	expression tag	UNP A4JGS8
A	-1	HIS	-	expression tag	UNP A4JGS8
A	0	HIS	-	expression tag	UNP A4JGS8
B	-7	MET	-	initiating methionine	UNP A4JGS8
B	-6	ALA	-	expression tag	UNP A4JGS8
B	-5	HIS	-	expression tag	UNP A4JGS8
B	-4	HIS	-	expression tag	UNP A4JGS8
B	-3	HIS	-	expression tag	UNP A4JGS8

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	expression tag	UNP A4JGS8
B	-1	HIS	-	expression tag	UNP A4JGS8
B	0	HIS	-	expression tag	UNP A4JGS8
C	-7	MET	-	initiating methionine	UNP A4JGS8
C	-6	ALA	-	expression tag	UNP A4JGS8
C	-5	HIS	-	expression tag	UNP A4JGS8
C	-4	HIS	-	expression tag	UNP A4JGS8
C	-3	HIS	-	expression tag	UNP A4JGS8
C	-2	HIS	-	expression tag	UNP A4JGS8
C	-1	HIS	-	expression tag	UNP A4JGS8
C	0	HIS	-	expression tag	UNP A4JGS8
D	-7	MET	-	initiating methionine	UNP A4JGS8
D	-6	ALA	-	expression tag	UNP A4JGS8
D	-5	HIS	-	expression tag	UNP A4JGS8
D	-4	HIS	-	expression tag	UNP A4JGS8
D	-3	HIS	-	expression tag	UNP A4JGS8
D	-2	HIS	-	expression tag	UNP A4JGS8
D	-1	HIS	-	expression tag	UNP A4JGS8
D	0	HIS	-	expression tag	UNP A4JGS8
E	-7	MET	-	initiating methionine	UNP A4JGS8
E	-6	ALA	-	expression tag	UNP A4JGS8
E	-5	HIS	-	expression tag	UNP A4JGS8
E	-4	HIS	-	expression tag	UNP A4JGS8
E	-3	HIS	-	expression tag	UNP A4JGS8
E	-2	HIS	-	expression tag	UNP A4JGS8
E	-1	HIS	-	expression tag	UNP A4JGS8
E	0	HIS	-	expression tag	UNP A4JGS8
F	-7	MET	-	initiating methionine	UNP A4JGS8
F	-6	ALA	-	expression tag	UNP A4JGS8
F	-5	HIS	-	expression tag	UNP A4JGS8
F	-4	HIS	-	expression tag	UNP A4JGS8
F	-3	HIS	-	expression tag	UNP A4JGS8
F	-2	HIS	-	expression tag	UNP A4JGS8
F	-1	HIS	-	expression tag	UNP A4JGS8
F	0	HIS	-	expression tag	UNP A4JGS8
G	-7	MET	-	initiating methionine	UNP A4JGS8
G	-6	ALA	-	expression tag	UNP A4JGS8
G	-5	HIS	-	expression tag	UNP A4JGS8
G	-4	HIS	-	expression tag	UNP A4JGS8
G	-3	HIS	-	expression tag	UNP A4JGS8
G	-2	HIS	-	expression tag	UNP A4JGS8
G	-1	HIS	-	expression tag	UNP A4JGS8

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP A4JGS8
H	-7	MET	-	initiating methionine	UNP A4JGS8
H	-6	ALA	-	expression tag	UNP A4JGS8
H	-5	HIS	-	expression tag	UNP A4JGS8
H	-4	HIS	-	expression tag	UNP A4JGS8
H	-3	HIS	-	expression tag	UNP A4JGS8
H	-2	HIS	-	expression tag	UNP A4JGS8
H	-1	HIS	-	expression tag	UNP A4JGS8
H	0	HIS	-	expression tag	UNP A4JGS8

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>O<sub>12</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	C	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	D	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	E	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	F	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
2	G	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	H	1	25	9	2	12	2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	C	O	0	0
			6	3	3		
3	G	1	Total	C	O	0	0
			6	3	3		
3	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	276	Total	O	0	0
			276	276		
5	B	154	Total	O	0	0
			154	154		
5	C	72	Total	O	0	0
			72	72		
5	D	65	Total	O	0	1
			66	66		
5	E	59	Total	O	0	0
			59	59		
5	F	101	Total	O	0	0
			101	101		
5	G	35	Total	O	0	0
			35	35		
5	H	27	Total	O	0	0
			27	27		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)





- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)





- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)





- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)





- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)





- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)



- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)



- Molecule 1: Alpha,alpha-trehalose-phosphate synthase (UDP-forming)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.17Å 103.02Å 218.47Å 90.00° 96.42° 90.00°	Depositor
Resolution (Å)	46.54 – 2.30 48.02 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.54-2.30) 98.9 (48.02-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575)	Depositor
R, $R_{free}$	0.178 , 0.225 0.179 , 0.227	Depositor DCC
$R_{free}$ test set	2000 reflections (1.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.3	Xtrriage
Anisotropy	0.678	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	28914	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, UDP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/3793	0.66	1/5154 (0.0%)
1	B	0.44	1/3588 (0.0%)	0.61	2/4880 (0.0%)
1	C	0.38	0/3629	0.54	0/4935
1	D	0.36	0/3635	0.53	0/4945
1	E	0.36	0/3510	0.53	0/4777
1	F	0.40	0/3625	0.56	0/4930
1	G	0.34	0/3372	0.53	0/4589
1	H	0.34	0/3398	0.51	0/4637
All	All	0.40	1/28550 (0.0%)	0.56	3/38847 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	103	GLU	CB-CG	5.13	1.61	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	453	LEU	CA-CB-CG	5.58	128.14	115.30
1	A	421	MET	CG-SD-CE	5.37	108.79	100.20
1	B	67	LEU	CA-CB-CG	5.32	127.55	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3687	0	3609	33	0
1	B	3505	0	3401	18	0
1	C	3542	0	3418	22	1
1	D	3547	0	3426	21	0
1	E	3423	0	3233	18	0
1	F	3537	0	3444	24	0
1	G	3291	0	3036	16	0
1	H	3311	0	3022	16	1
2	A	25	0	11	0	0
2	B	25	0	11	1	0
2	C	25	0	11	1	0
2	D	25	0	11	0	0
2	E	25	0	11	0	0
2	F	25	0	11	0	0
2	G	25	0	11	0	0
2	H	25	0	11	0	0
3	A	30	0	40	6	0
3	B	6	0	8	1	0
3	C	6	0	8	1	0
3	D	12	0	16	0	0
3	E	6	0	8	0	0
3	F	6	0	8	0	0
3	G	6	0	8	0	0
3	H	6	0	8	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	276	0	0	6	0
5	B	154	0	0	2	0
5	C	72	0	0	2	0
5	D	66	0	0	1	0
5	E	59	0	0	3	0
5	F	101	0	0	3	0
5	G	35	0	0	0	0
5	H	27	0	0	0	0
All	All	28914	0	26781	160	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:ARG:HH12	3:A:503:GOL:H11	1.38	0.88
1:B:43:ASN:HB3	1:B:66:THR:HB	1.66	0.76
1:B:76:GLN:OE1	1:B:109:ARG:NH2	2.18	0.75
1:A:109:ARG:NH1	5:A:602:HOH:O	2.25	0.69
1:A:344:ARG:HH22	3:A:503:GOL:H32	1.59	0.67

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2:SER:OG	1:H:209:ASP:OD1[2_556]	2.13	0.07

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/481 (98%)	456 (97%)	13 (3%)	2 (0%)	34	42
1	B	448/481 (93%)	437 (98%)	11 (2%)	0	100	100
1	C	453/481 (94%)	442 (98%)	10 (2%)	1 (0%)	47	58
1	D	452/481 (94%)	440 (97%)	11 (2%)	1 (0%)	47	58
1	E	442/481 (92%)	431 (98%)	11 (2%)	0	100	100
1	F	453/481 (94%)	442 (98%)	11 (2%)	0	100	100
1	G	432/481 (90%)	420 (97%)	12 (3%)	0	100	100
1	H	440/481 (92%)	428 (97%)	12 (3%)	0	100	100
All	All	3591/3848 (93%)	3496 (97%)	91 (2%)	4 (0%)	51	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	209	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	210	ASP
1	C	22	GLY
1	D	22	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/383 (97%)	367 (99%)	3 (1%)	81	91
1	B	346/383 (90%)	342 (99%)	4 (1%)	71	84
1	C	348/383 (91%)	340 (98%)	8 (2%)	50	67
1	D	351/383 (92%)	346 (99%)	5 (1%)	67	81
1	E	326/383 (85%)	320 (98%)	6 (2%)	59	75
1	F	350/383 (91%)	346 (99%)	4 (1%)	73	86
1	G	301/383 (79%)	299 (99%)	2 (1%)	84	92
1	H	302/383 (79%)	297 (98%)	5 (2%)	60	76
All	All	2694/3064 (88%)	2657 (99%)	37 (1%)	67	81

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	266	LEU
1	H	358	TYR
1	F	364	ASP
1	H	132	ASP
1	C	288	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	C	9	ASN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	F	140	HIS
1	G	9	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 3 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UDP	G	501	-	24,26,26	0.33	0	37,40,40	0.65	1 (2%)
2	UDP	E	501	-	24,26,26	0.51	0	37,40,40	0.65	1 (2%)
2	UDP	B	501	-	24,26,26	0.28	0	37,40,40	0.64	1 (2%)
3	GOL	A	503	-	5,5,5	0.43	0	5,5,5	0.51	0
3	GOL	A	505	-	5,5,5	0.33	0	5,5,5	0.28	0
3	GOL	E	502	-	5,5,5	0.24	0	5,5,5	0.57	0
3	GOL	G	502	-	5,5,5	0.33	0	5,5,5	0.47	0
2	UDP	A	501	-	24,26,26	0.49	0	37,40,40	0.89	1 (2%)
2	UDP	H	501	-	24,26,26	0.53	0	37,40,40	0.74	2 (5%)
2	UDP	C	501	-	24,26,26	0.43	0	37,40,40	0.64	1 (2%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	A	506	-	5,5,5	0.41	0	5,5,5	0.68	0
2	UDP	F	501	-	24,26,26	0.36	0	37,40,40	0.61	1 (2%)
3	GOL	B	502	-	5,5,5	0.30	0	5,5,5	0.79	0
3	GOL	F	502	-	5,5,5	0.35	0	5,5,5	0.47	0
3	GOL	A	504	-	5,5,5	0.43	0	5,5,5	1.02	0
2	UDP	D	501	-	24,26,26	0.42	0	37,40,40	0.46	0
3	GOL	C	502	-	5,5,5	0.39	0	5,5,5	0.27	0
3	GOL	H	502	-	5,5,5	0.32	0	5,5,5	0.53	0
3	GOL	D	503	-	5,5,5	0.33	0	5,5,5	0.34	0
3	GOL	A	502	-	5,5,5	0.45	0	5,5,5	0.34	0
3	GOL	D	502	-	5,5,5	0.27	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. <sup>1,2</sup> means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	G	501	-	-	4/16/32/32	0/2/2/2
2	UDP	E	501	-	-	4/16/32/32	0/2/2/2
2	UDP	B	501	-	-	6/16/32/32	0/2/2/2
3	GOL	A	503	-	-	2/4/4/4	-
3	GOL	A	505	-	-	2/4/4/4	-
3	GOL	E	502	-	-	4/4/4/4	-
3	GOL	G	502	-	-	0/4/4/4	-
2	UDP	A	501	-	-	3/16/32/32	0/2/2/2
2	UDP	H	501	-	-	6/16/32/32	0/2/2/2
2	UDP	C	501	-	-	4/16/32/32	0/2/2/2
3	GOL	A	506	-	-	0/4/4/4	-
2	UDP	F	501	-	-	3/16/32/32	0/2/2/2
3	GOL	B	502	-	-	2/4/4/4	-
3	GOL	F	502	-	-	2/4/4/4	-
3	GOL	A	504	-	-	2/4/4/4	-
2	UDP	D	501	-	-	4/16/32/32	0/2/2/2
3	GOL	C	502	-	-	2/4/4/4	-
3	GOL	H	502	-	-	2/4/4/4	-
3	GOL	D	503	-	-	2/4/4/4	-
3	GOL	A	502	-	-	0/4/4/4	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	502	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	UDP	PA-O3A-PB	-2.77	123.31	132.83
2	C	501	UDP	O3B-PB-O3A	2.38	112.61	104.64
2	H	501	UDP	O3B-PB-O3A	2.37	112.58	104.64
2	H	501	UDP	PA-O3A-PB	-2.34	124.79	132.83
2	G	501	UDP	O3B-PB-O3A	2.29	112.30	104.64

There are no chirality outliers.

5 of 56 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	UDP	O4'-C4'-C5'-O5'
2	B	501	UDP	C5'-O5'-PA-O2A
3	A	504	GOL	O1-C1-C2-C3
3	A	505	GOL	O1-C1-C2-O2
3	A	505	GOL	O1-C1-C2-C3


There are no ring outliers.



5 monomers are involved in 8 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	UDP	1	0
3	A	503	GOL	6	0
2	C	501	UDP	1	0
3	B	502	GOL	1	0
3	C	502	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	468/481 (97%)	0.05	2 (0%) 92 95	28, 38, 60, 109	0
1	B	454/481 (94%)	0.11	10 (2%) 62 69	28, 47, 72, 97	0
1	C	458/481 (95%)	0.33	25 (5%) 25 31	37, 60, 84, 109	0
1	D	456/481 (94%)	0.00	5 (1%) 80 85	39, 57, 82, 97	0
1	E	447/481 (92%)	0.39	30 (6%) 17 23	32, 66, 108, 151	0
1	F	456/481 (94%)	0.11	5 (1%) 80 85	39, 56, 77, 97	0
1	G	439/481 (91%)	0.44	54 (12%) 4 6	40, 67, 111, 131	0
1	H	443/481 (92%)	0.66	54 (12%) 4 6	54, 81, 111, 143	0
All	All	3621/3848 (94%)	0.26	185 (5%) 28 35	28, 58, 99, 151	0

The worst 5 of 185 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	55	ALA	6.8
1	G	68	GLY	6.5
1	G	42	TRP	6.4
1	E	20	SER	6.1
1	G	77	TYR	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	H	502	6/6	0.85	0.21	59,64,68,68	0
3	GOL	A	505	6/6	0.86	0.38	66,79,90,99	0
3	GOL	A	506	6/6	0.90	0.21	36,63,71,78	0
3	GOL	D	503	6/6	0.91	0.29	66,73,76,77	0
2	UDP	H	501	25/25	0.91	0.16	59,86,106,108	0
3	GOL	A	503	6/6	0.93	0.17	39,64,68,69	0
3	GOL	A	502	6/6	0.95	0.16	32,37,41,44	0
2	UDP	E	501	25/25	0.96	0.14	40,60,83,85	0
2	UDP	F	501	25/25	0.96	0.14	41,53,71,73	0
2	UDP	G	501	25/25	0.96	0.11	53,61,78,82	0
3	GOL	C	502	6/6	0.96	0.14	42,45,48,52	0
2	UDP	C	501	25/25	0.96	0.15	38,56,71,73	0
3	GOL	G	502	6/6	0.96	0.13	59,62,63,64	0
2	UDP	D	501	25/25	0.96	0.13	41,51,61,68	0
3	GOL	E	502	6/6	0.97	0.15	45,48,52,53	0
3	GOL	F	502	6/6	0.97	0.17	39,42,48,49	0
2	UDP	A	501	25/25	0.97	0.16	26,37,57,66	0
2	UDP	B	501	25/25	0.97	0.13	37,56,76,81	0
4	MG	D	504	1/1	0.97	0.08	61,61,61,61	0
3	GOL	B	502	6/6	0.98	0.17	37,42,47,49	0
3	GOL	A	504	6/6	0.98	0.18	27,32,39,44	0
4	MG	A	507	1/1	0.98	0.09	38,38,38,38	0
4	MG	B	503	1/1	0.98	0.07	50,50,50,50	0
3	GOL	D	502	6/6	0.98	0.16	41,45,50,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around UDP H 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UDP E 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UDP F 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UDP G 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UDP C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UDP D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UDP A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around UDP B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.