



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 04:59 PM EST

PDB ID : 8TWS
Title : AvrB bound with UDP-rhamnose and RIN4 C-NOI motif
Authors : Peng, W.; Orth, K.
Deposited on : 2023-08-21
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	
2	B	30	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4972 atoms, of which 2422 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Avirulence protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	294	4541	1441	2225	419	449	7	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	expression tag	UNP P13835
A	0	HIS	-	expression tag	UNP P13835

- Molecule 2 is a protein called RPM1-interacting protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	25	374	126	175	33	40	0	0	0

- Molecule 3 is [(2 {R},3 {S},4 {R},5 {R})-5-[2,4-bis(oxidanylidene)pyrimidin-1-yl]-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl [(2 {R},3 {R},4 {R},5 {R},6 {S})-6-methyl-3,4,5-tris(oxidanyl)oxan-2-yl] hydrogen phosphate (three-letter code: AWU) (formula: C₁₅H₂₄N₂O₁₆P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
3	A	1	57	15	22	2	16	2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Avirulence protein B



- Molecule 2: RPM1-interacting protein 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	58.62Å 123.58Å 45.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.97 – 2.85 52.96 – 2.85	Depositor EDS
% Data completeness (in resolution range)	94.9 (52.97-2.85) 94.9 (52.96-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 2.86Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.280 , 0.336 0.283 , 0.339	Depositor DCC
R_{free} test set	777 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å ²)	60.9	Xtrriage
Anisotropy	0.299	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.34$, $\langle L^2 \rangle = 0.17$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	4972	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AWU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/2366	0.55	0/3213
2	B	0.29	0/206	0.44	0/280
All	All	0.28	0/2572	0.55	0/3493

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2316	2225	2226	56	1
2	B	199	175	175	5	0
3	A	35	22	0	1	0
All	All	2550	2422	2401	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:THR:OG1	1:A:62:ASN:ND2	1.87	1.05
1:A:59:THR:HG1	1:A:62:ASN:HD22	1.25	0.82
1:A:197:GLU:N	1:A:197:GLU:OE1	2.16	0.78
1:A:93:GLY:O	1:A:97:GLU:HG3	1.87	0.74
1:A:167:TYR:CE1	1:A:172:VAL:HG23	2.33	0.64
1:A:153:LEU:HD22	1:A:206:HIS:CD2	2.35	0.62
1:A:62:ASN:O	1:A:65:TYR:N	2.34	0.61
1:A:305:GLU:O	1:A:309:VAL:HG23	2.02	0.59
1:A:117:ARG:NE	1:A:262:MET:O	2.35	0.58
1:A:253:ARG:NH1	1:A:309:VAL:HG21	2.21	0.56
1:A:167:TYR:HE1	1:A:172:VAL:HG23	1.73	0.54
1:A:159:VAL:HG23	1:A:161:VAL:HG23	1.89	0.53
1:A:239:SER:OG	1:A:240:SER:N	2.40	0.53
1:A:183:ASP:O	1:A:187:LEU:HD23	2.10	0.52
1:A:206:HIS:O	1:A:210:TYR:N	2.43	0.51
1:A:37:LEU:HD22	1:A:286:GLU:HG3	1.93	0.51
1:A:130:PRO:HD2	1:A:131:TYR:CD2	2.45	0.51
1:A:88:PHE:N	1:A:231:ASN:OD1	2.44	0.50
1:A:37:LEU:CD2	1:A:286:GLU:HG3	2.42	0.50
2:B:156:GLU:O	2:B:157:ASN:OD1	2.28	0.50
1:A:208:GLN:HE22	2:B:163:ASP:H	1.60	0.49
1:A:91:LEU:HD12	1:A:91:LEU:O	2.12	0.49
1:A:107:GLU:HB2	1:A:108:PRO:HD3	1.95	0.48
1:A:191:VAL:HG13	1:A:192:ALA:N	2.29	0.48
1:A:61:GLU:O	1:A:64:ALA:HB3	2.15	0.47
1:A:177:THR:HG21	1:A:179:ARG:NH1	2.31	0.45
1:A:188:ARG:NH1	1:A:189:ASP:OD1	2.49	0.45
1:A:288:PRO:HG2	1:A:316:LEU:HB3	1.99	0.45
1:A:291:ARG:NH1	1:A:319:ASP:OD2	2.43	0.45
1:A:237:ALA:O	1:A:283:HIS:NE2	2.36	0.45
1:A:52:ASP:O	1:A:53:ASP:HB2	2.17	0.44
1:A:181:ALA:HB2	1:A:212:PRO:HB3	1.99	0.44
1:A:188:ARG:HA	1:A:191:VAL:HG12	1.99	0.44
1:A:207:LEU:HD12	2:B:160:SER:O	2.17	0.44
1:A:164:GLN:NE2	1:A:300:ALA:O	2.50	0.44
1:A:37:LEU:HD22	1:A:286:GLU:CG	2.47	0.44
1:A:36:GLN:HB3	1:A:70:TYR:OH	2.18	0.44
1:A:222:TYR:O	1:A:226:ILE:HG13	2.18	0.43
1:A:266:ARG:NH2	3:A:1001:AWU:O2B	2.51	0.43
1:A:153:LEU:HD22	1:A:206:HIS:CG	2.54	0.43
1:A:109:LEU:HD23	1:A:109:LEU:HA	1.91	0.43
2:B:165:TYR:O	2:B:167:HIS:CD2	2.72	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:LEU:HD23	1:A:180:VAL:HG22	2.00	0.42
1:A:91:LEU:HD12	1:A:91:LEU:C	2.39	0.42
1:A:223:LEU:N	1:A:224:PRO:CD	2.83	0.42
1:A:91:LEU:O	1:A:94:ARG:N	2.53	0.42
1:A:298:ILE:HG22	1:A:302:LEU:HD12	2.02	0.41
1:A:85:ILE:HD11	1:A:94:ARG:HD3	2.01	0.41
1:A:246:GLU:O	1:A:249:SER:N	2.53	0.41
1:A:198:LEU:HB3	1:A:202:ALA:HB3	2.02	0.41
1:A:204:LYS:HE2	1:A:208:GLN:NE2	2.35	0.41
1:A:204:LYS:HE2	1:A:208:GLN:HE21	1.86	0.41
1:A:188:ARG:HG2	1:A:188:ARG:HH11	1.86	0.41
1:A:109:LEU:O	1:A:113:PHE:HD1	2.04	0.41
1:A:49:ARG:HG2	1:A:57:SER:OG	2.21	0.40
1:A:208:GLN:OE1	1:A:208:GLN:HA	2.21	0.40
1:A:123:ALA:HB3	1:A:217:HIS:HB2	2.03	0.40
1:A:184:THR:OG1	2:B:149:PRO:HD2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:GLU:OE2	1:A:94:ARG:NH2[2_655]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/323 (90%)	254 (87%)	35 (12%)	3 (1%)	15	40
2	B	23/30 (77%)	19 (83%)	3 (13%)	1 (4%)	2	8
All	All	315/353 (89%)	273 (87%)	38 (12%)	4 (1%)	12	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	ASP
1	A	109	LEU
2	B	157	ASN
1	A	128	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/274 (89%)	234 (96%)	10 (4%)	30	61
2	B	21/26 (81%)	20 (95%)	1 (5%)	25	55
All	All	265/300 (88%)	254 (96%)	11 (4%)	30	60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	ASP
1	A	81	SER
1	A	94	ARG
1	A	152	ASN
1	A	185	GLU
1	A	186	TYR
1	A	221	SER
1	A	235	ARG
1	A	249	SER
1	A	254	THR
2	B	158	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	145	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	206	HIS
1	A	208	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	AWU	A	1001	-	34,37,37	3.36	14 (41%)	52,57,57	1.95	15 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AWU	A	1001	-	-	5/21/57/57	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	AWU	O4C-C1C	7.68	1.60	1.42
3	A	1001	AWU	C2-N1	7.20	1.50	1.38
3	A	1001	AWU	C2-N3	6.88	1.50	1.38
3	A	1001	AWU	C2C-C1C	-6.54	1.32	1.53
3	A	1001	AWU	C3C-C4C	-6.44	1.36	1.53
3	A	1001	AWU	C6-C5	5.78	1.48	1.35
3	A	1001	AWU	C4-N3	4.04	1.45	1.38
3	A	1001	AWU	O4C-C4C	3.94	1.53	1.45
3	A	1001	AWU	C3C-C2C	3.72	1.63	1.53
3	A	1001	AWU	C6-N1	3.20	1.45	1.38
3	A	1001	AWU	O4-C4	-2.93	1.18	1.24
3	A	1001	AWU	O3'-C3'	2.59	1.49	1.43
3	A	1001	AWU	C5-C4	2.37	1.48	1.43
3	A	1001	AWU	O2-C2	-2.25	1.18	1.23

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	AWU	C4-N3-C2	-5.12	119.82	126.58
3	A	1001	AWU	C1'-C2'-C3'	4.35	119.05	110.00
3	A	1001	AWU	O3A-PB-O3B	4.21	110.97	102.48
3	A	1001	AWU	N3-C2-N1	3.96	120.15	114.89
3	A	1001	AWU	C2'-C3'-C4'	3.73	117.33	110.82
3	A	1001	AWU	C5-C4-N3	3.36	119.87	114.84
3	A	1001	AWU	C3C-C2C-C1C	3.36	107.81	101.43
3	A	1001	AWU	O5'-C1'-C2'	3.33	117.41	110.35
3	A	1001	AWU	O4-C4-C5	-2.82	120.19	125.16
3	A	1001	AWU	C2C-C3C-C4C	2.76	108.01	102.64
3	A	1001	AWU	PB-O3A-PA	-2.53	124.16	132.83
3	A	1001	AWU	O5'-C1'-O3B	-2.45	108.17	111.36
3	A	1001	AWU	O5'-C5'-C6'	2.43	111.95	106.70
3	A	1001	AWU	C3'-C4'-C5'	2.31	113.36	109.77
3	A	1001	AWU	C6'-C5'-C4'	-2.07	109.24	113.07

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1001	AWU	C5C-O5C-PA-O1A
3	A	1001	AWU	C5C-O5C-PA-O2A
3	A	1001	AWU	PA-O3A-PB-O3B
3	A	1001	AWU	C5C-O5C-PA-O3A
3	A	1001	AWU	PB-O3A-PA-O2A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	AWU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/323 (91%)	1.03	48 (16%) 1 1	38, 58, 114, 145	0
2	B	25/30 (83%)	1.93	7 (28%) 0 0	50, 94, 129, 183	0
All	All	319/353 (90%)	1.10	55 (17%) 1 1	38, 58, 118, 183	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	GLN	7.9
2	B	151	PHE	6.9
1	A	149	MET	6.6
1	A	27	ARG	6.3
2	B	149	PRO	5.6
2	B	152	GLY	5.3
1	A	187	LEU	5.3
1	A	107	GLU	5.2
1	A	238	ILE	4.9
1	A	191	VAL	4.8
2	B	155	ASP	4.3
1	A	153	LEU	4.3
1	A	156	ASP	4.1
1	A	130	PRO	4.0
1	A	151	ASP	3.9
1	A	285	ILE	3.7
1	A	84	GLY	3.7
1	A	108	PRO	3.7
1	A	242	LEU	3.5
1	A	40	TYR	3.5
1	A	80	ILE	3.1
1	A	194	LEU	3.1
1	A	247	LEU	3.1
1	A	131	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	154	TRP	3.0
1	A	160	LYS	3.0
1	A	184	THR	3.0
1	A	206	HIS	2.9
1	A	157	LEU	2.9
1	A	150	GLN	2.8
1	A	182	THR	2.8
1	A	303	SER	2.8
1	A	222	TYR	2.8
1	A	105	ARG	2.7
1	A	122	ASP	2.6
1	A	239	SER	2.5
1	A	126	PRO	2.5
1	A	109	LEU	2.5
1	A	251	ILE	2.5
1	A	103	LEU	2.4
1	A	54	SER	2.4
1	A	78	ASP	2.4
1	A	53	ASP	2.3
1	A	198	LEU	2.3
1	A	318	SER	2.2
2	B	153	ASP	2.2
1	A	154	PHE	2.2
1	A	110	TYR	2.2
1	A	199	GLY	2.1
2	B	1	ALA	2.1
1	A	145	HIS	2.1
1	A	129	LYS	2.0
1	A	192	ALA	2.0
1	A	188	ARG	2.0
1	A	85	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	AWU	A	1001	35/35	0.86	0.18	54,64,78,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.