



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:35 pm BST

PDB ID : 4TZS  
Title : Structure of *C. elegans* HTP-2 bound to HIM-3 closure motif, P212121 form  
Authors : Rosenberg, S.C.; Corbett, K.D.  
Deposited on : 2014-07-10  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	 4% 75% 15% 9%
1	B	253	 6% 74% 16% 9%
2	C	20	 20% 80% 5% 15%
2	D	20	 5% 70% 15% 15%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein HTP-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	229	Total 1825	C 1162	N 312	O 342	S 9	0	0	0
1	B	230	Total 1831	C 1165	N 313	O 344	S 9	0	0	0

- Molecule 2 is a protein called C. elegans HIM-3 closure motif.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	17	Total 129	C 78	N 25	O 26	0	0	0
2	D	17	Total 129	C 78	N 25	O 26	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	272	SER	-	expression tag	UNP G5EBG0
C	273	ASN	-	expression tag	UNP G5EBG0
C	274	ALA	-	expression tag	UNP G5EBG0
D	272	SER	-	expression tag	UNP G5EBG0
D	273	ASN	-	expression tag	UNP G5EBG0
D	274	ALA	-	expression tag	UNP G5EBG0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total 12	O 12	0	0
3	B	5	Total 5	O 5	0	0
3	C	1	Total 1	O 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein HTP-2




- Molecule 1: Protein HTP-2



- Molecule 2: *C. elegans* HIM-3 closure motif



- Molecule 2: *C. elegans* HIM-3 closure motif



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.24Å 82.68Å 118.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.04 – 2.55 39.04 – 2.55	Depositor EDS
% Data completeness (in resolution range)	98.4 (39.04-2.55) 98.7 (39.04-2.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.66 (at 2.54Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.229 , 0.264 0.235 , 0.267	Depositor DCC
$R_{free}$ test set	1086 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.7	Xtrriage
Anisotropy	0.670	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3932	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.4071e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1865	0.44	0/2521
1	B	0.27	0/1871	0.45	0/2529
2	C	0.25	0/130	0.43	0/173
2	D	0.23	0/130	0.45	0/173
All	All	0.27	0/3996	0.44	0/5396

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1825	0	1788	22	0
1	B	1831	0	1793	27	0
2	C	129	0	128	1	0
2	D	129	0	128	3	0
3	A	12	0	0	1	0
3	B	5	0	0	0	0
3	C	1	0	0	0	0
All	All	3932	0	3837	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:ARG:NH1	1:A:60:LEU:O	2.23	0.71
1:B:55:ARG:NH1	1:B:60:LEU:O	2.25	0.69
1:A:143:THR:HG22	1:A:144:ASP:H	1.58	0.69
1:B:210:ARG:NH2	2:D:274:ALA:O	2.29	0.66
1:B:96:ALA:HA	1:B:211:ILE:HD13	1.79	0.64
1:B:136:GLY:O	1:B:236:ARG:NH2	2.31	0.63
1:A:40:ASN:OD1	1:A:101:LYS:NZ	2.33	0.62
1:A:114:GLU:HG3	1:A:122:ILE:HD11	1.83	0.60
1:A:201:TYR:HE1	1:A:211:ILE:HG13	1.68	0.58
1:B:156:LEU:HD21	1:B:243:HIS:HB2	1.87	0.56
1:A:67:LYS:NZ	1:A:69:TYR:OH	2.29	0.55
1:B:143:THR:HG22	1:B:144:ASP:H	1.74	0.53
1:B:143:THR:HG22	1:B:144:ASP:N	2.24	0.53
1:A:103:GLY:HA2	1:A:131:TYR:CE1	2.44	0.52
1:B:141:LEU:HA	1:B:142:SER:HB3	1.91	0.51
1:B:103:GLY:HA2	1:B:131:TYR:CE1	2.45	0.51
1:B:156:LEU:HD11	1:B:243:HIS:HB2	1.90	0.51
1:A:64:TYR:OH	1:A:85:ARG:NH1	2.44	0.51
1:A:57:ARG:NH2	1:A:123:GLU:OE1	2.44	0.51
1:B:114:GLU:HG3	1:B:122:ILE:HD11	1.93	0.50
1:A:186:LEU:HD22	1:A:250:SER:HB2	1.93	0.49
1:B:74:LEU:HD22	1:B:172:VAL:HG13	1.95	0.48
1:B:116:ASP:HA	1:B:117:GLY:HA2	1.60	0.48
1:B:186:LEU:HD22	1:B:250:SER:HB2	1.96	0.47
1:A:140:ARG:HG2	1:A:141:LEU:H	1.78	0.47
1:B:49:VAL:O	1:B:53:VAL:HG23	2.15	0.47
1:A:154:ALA:O	1:A:158:GLN:HG2	2.14	0.47
1:B:167:VAL:HA	1:B:170:GLN:NE2	2.30	0.47
1:B:60:LEU:HD21	2:D:285:ILE:HD13	1.97	0.47
1:B:207:SER:O	2:D:275:ARG:HB2	2.16	0.46
1:A:142:SER:N	1:A:204:ASP:OD1	2.47	0.46
1:A:116:ASP:HA	1:A:117:GLY:HA2	1.59	0.44
1:B:116:ASP:HB2	1:B:194:THR:HG21	1.99	0.44
1:A:207:SER:O	2:C:275:ARG:HB2	2.18	0.43
1:B:113:THR:HG22	1:B:121:ALA:HA	2.00	0.43
1:A:49:VAL:HG11	1:A:244:MET:SD	2.59	0.42
1:B:38:SER:OG	1:B:240:HIS:ND1	2.33	0.42
1:A:174:ILE:HG12	1:A:233:GLY:HA3	2.02	0.42
1:B:186:LEU:HD23	1:B:251:MET:HB3	2.02	0.42
1:B:57:ARG:HD2	1:B:57:ARG:HA	1.90	0.42
1:A:57:ARG:HD2	1:A:57:ARG:HA	1.79	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:VAL:HA	1:B:170:GLN:HE21	1.86	0.41
1:B:170:GLN:HE21	1:B:170:GLN:HB2	1.69	0.41
1:B:228:GLN:HB2	1:B:251:MET:HG3	2.03	0.41
1:A:39:SER:OG	1:A:101:LYS:HD2	2.20	0.41
1:A:168:ARG:NH1	3:A:303:HOH:O	2.53	0.41
1:A:140:ARG:HG2	1:A:141:LEU:N	2.37	0.40
1:A:56:GLN:HG2	1:A:179:GLN:HA	2.04	0.40
1:B:47:TYR:OH	1:B:79:LEU:O	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/253 (89%)	219 (97%)	5 (2%)	1 (0%)	34	46
1	B	226/253 (89%)	216 (96%)	10 (4%)	0	100	100
2	C	15/20 (75%)	14 (93%)	1 (7%)	0	100	100
2	D	15/20 (75%)	14 (93%)	1 (7%)	0	100	100
All	All	481/546 (88%)	463 (96%)	17 (4%)	1 (0%)	47	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/224 (89%)	198 (99%)	2 (1%)	76	84
1	B	201/224 (90%)	196 (98%)	5 (2%)	47	62
2	C	14/17 (82%)	14 (100%)	0	100	100
2	D	14/17 (82%)	14 (100%)	0	100	100
All	All	429/482 (89%)	422 (98%)	7 (2%)	62	77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	ARG
1	A	220	PHE
1	B	55	ARG
1	B	73	ASN
1	B	133	GLU
1	B	162	GLU
1	B	220	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	HIS
1	B	87	HIS
1	B	170	GLN
1	B	234	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	229/253 (90%)	0.36	11 (4%) 30 37	50, 80, 115, 133	0
1	B	230/253 (90%)	0.52	15 (6%) 18 22	51, 80, 115, 129	0
2	C	17/20 (85%)	0.97	4 (23%) 0 0	69, 87, 129, 140	0
2	D	17/20 (85%)	0.64	1 (5%) 22 26	63, 82, 109, 131	0
All	All	493/546 (90%)	0.46	31 (6%) 20 23	50, 80, 116, 140	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	289	ASN	4.7
2	C	289	ASN	4.3
1	B	22	LYS	4.2
1	B	143	THR	4.0
1	B	141	LEU	3.8
1	A	142	SER	3.6
1	A	154	ALA	3.2
1	B	153	PHE	3.2
2	C	288	LYS	3.1
1	B	152	HIS	3.1
1	B	18	ALA	3.0
1	A	152	HIS	2.5
1	A	22	LYS	2.4
1	B	37	ARG	2.4
2	C	273	ASN	2.3
1	A	153	PHE	2.3
1	A	135	GLY	2.3
1	B	64	TYR	2.3
1	A	119	VAL	2.3
1	A	225	ASP	2.3
1	B	114	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	122	ILE	2.2
1	A	18	ALA	2.2
1	A	138	VAL	2.2
1	A	143	THR	2.1
2	C	274	ALA	2.1
1	B	119	VAL	2.0
1	B	218	SER	2.0
1	B	101	LYS	2.0
1	B	160	ARG	2.0
1	B	92	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.