

Full wwPDB X-ray Structure Validation Report (i)

Dec 17, 2023 - 02:52 am GMT

PDB ID	:	4U0R
Title	:	Plasmodium falciparum reticulocyte-binding protein homologue 5 (PfRH5)
		bound to monoclonal antibody 9AD4
Authors	:	Wright, K.E.; Hjerrild, K.A.; Bartlett, J.; Douglas, A.D.; Jin, J.; Brown, R.E.;
		Ashfield, R.; Clemmensen, S.B.; de Jongh, W.A.; Draper, S.J.; Higgins, M.K.
Deposited on	:	2014-07-14
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575(2.30-2.30)
Sidechain outliers	138945	5575(2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	526	3% 42%	,)	11%	•	45%	-	
2	В	258	3%	67%			15%	•	14%
3	С	238	<u>2%</u>	70%			18%		• 9%



4U0R

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6005 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Reticulocyte binding protein 5.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	288	Total 2450	C 1584	N 410	0 441	S 15	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	216	ALA	THR	conflict	UNP B2L3N7

• Molecule 2 is a protein called Monoclonal antibody 9AD4 heavy chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	В	221	Total 1668	C 1060	N 272	0 327	S 9	0	0	0

• Molecule 3 is a protein called Monoclonal antibody 9AD4 light chain.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
3	С	217	Total 1677	C 1046	N 286	O 338	${f S}7$	0	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	45	$\begin{array}{cc} \text{Total} & \text{O} \\ 45 & 45 \end{array}$	0	0
4	В	100	Total O 100 100	0	0
4	С	65	$\begin{array}{cc} \text{Total} & \text{O} \\ 65 & 65 \end{array}$	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Reticulocyte binding protein 5

• Molecule 3: Monoclonal antibody 9AD4 light chain







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	39.79Å 86.66Å 324.04Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution(A)	40.83 - 2.30	Depositor
Resolution (A)	40.83 - 2.30	EDS
% Data completeness	96.1 (40.83-2.30)	Depositor
(in resolution range)	$96.1 \ (40.83 - 2.30)$	EDS
R_{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.75 (at 2.29 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
P. P.	0.221 , 0.258	Depositor
II, II, <i>free</i>	0.236 , 0.253	DCC
R_{free} test set	2462 reflections $(4.99%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	41.1	Xtriage
Anisotropy	0.603	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.31 , 41.9	EDS
L-test for $twinning^2$	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6005	wwPDB-VP
Average B, all atoms $(Å^2)$	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Moi Chain		# Z > 5	RMSZ	# Z > 5	
1	А	0.52	0/2502	0.70	2/3354~(0.1%)	
2	В	0.55	1/1714~(0.1%)	0.82	0/2343	
3	С	0.60	2/1717~(0.1%)	0.73	1/2332~(0.0%)	
All	All	0.55	3/5933~(0.1%)	0.75	3/8029~(0.0%)	

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	С	51	LEU	C-N	-7.71	1.16	1.34
3	С	50	LEU	C-N	-7.60	1.16	1.34
2	В	204	CYS	CB-SG	-6.51	1.71	1.82

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	С	51	LEU	O-C-N	-6.90	111.66	122.70
1	А	501	LEU	N-CA-CB	-5.95	98.51	110.40
1	А	503	MET	N-CA-CB	-5.09	101.43	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	${ m H}({ m model})$	H(added)	Clashes	Symm-Clashes
1	А	2450	0	2481	61	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	1668	0	1612	35	0
3	С	1677	0	1605	24	0
4	А	45	0	0	15	0
4	В	100	0	0	14	0
4	С	65	0	0	5	0
All	All	6005	0	5698	116	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (116) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A + 1	A + 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:238:LEU:CD2	1:A:300:PHE:HE2	1.45	1.29
1:A:238:LEU:CD2	1:A:300:PHE:CE2	2.17	1.27
1:A:496:HIS:O	1:A:501:LEU:HD13	1.32	1.23
2:B:21:SER:HB2	4:B:309:HOH:O	1.35	1.22
1:A:501:LEU:HD12	1:A:501:LEU:N	1.59	1.16
1:A:496:HIS:O	1:A:501:LEU:CD1	1.95	1.14
1:A:238:LEU:HD21	1:A:300:PHE:CE2	1.79	1.13
1:A:238:LEU:HD23	1:A:300:PHE:HE2	1.15	1.10
1:A:300:PHE:CE2	1:A:409:THR:HG22	1.87	1.09
1:A:500:VAL:C	1:A:501:LEU:HD12	1.75	1.06
3:C:217:GLU:HG2	4:C:352:HOH:O	1.62	0.98
2:B:97:THR:HG21	2:B:109:PHE:HB3	1.45	0.98
1:A:501:LEU:N	1:A:501:LEU:CD1	2.30	0.91
3:C:23:CYS:HG	3:C:92:CYS:HG	1.15	0.89
1:A:451:GLN:HE22	1:A:453:ASP:HB3	1.44	0.83
1:A:478:MET:O	1:A:482:THR:HG23	1.78	0.82
1:A:238:LEU:HD23	1:A:300:PHE:CE2	2.02	0.82
3:C:188:ASP:OD2	3:C:192:ARG:NH1	2.12	0.81
2:B:128:PRO:HB3	2:B:154:TYR:HB3	1.66	0.77
2:B:39:GLN:HE22	3:C:42:GLN:HE22	1.33	0.76
1:A:500:VAL:CA	1:A:501:LEU:HD12	2.17	0.75
1:A:476:ARG:NH2	4:A:623:HOH:O	1.86	0.75
2:B:97:THR:CG2	2:B:109:PHE:HB3	2.17	0.73
3:C:154:ILE:HD11	3:C:183:LEU:HD21	1.72	0.71
1:A:366:LYS:HE3	4:A:636:HOH:O	1.91	0.70
1:A:164:LEU:HD22	1:A:478:MET:HB3	1.74	0.70
1:A:503:MET:O	1:A:504:LYS:C	2.30	0.70
1:A:238:LEU:HD21	1:A:300:PHE:CZ	2.26	0.69



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:B:13:GLN:NE2	4:B:301:HOH:O	2.26	0.68
2:B:65:THR:HG21	4:B:397:HOH:O	1.92	0.68
2:B:64:VAL:HG13	2:B:68:PHE:HB2	1.80	0.63
2:B:156:PRO:O	2:B:208:HIS:HE1	1.82	0.62
2:B:7:SER:HB2	4:B:309:HOH:O	1.99	0.61
1:A:381:SER:HB3	4:A:640:HOH:O	2.00	0.61
2:B:193:THR:HG23	4:B:315:HOH:O	2.00	0.61
2:B:222:ARG:NH2	3:C:124:PRO:O	2.34	0.61
1:A:345:CYS:HG	1:A:351:CYS:HG	0.69	0.60
1:A:300:PHE:CD2	1:A:409:THR:HG22	2.36	0.60
2:B:72:ARG:HD3	2:B:74:ASN:OD1	2.02	0.60
2:B:124:LYS:HD3	4:B:301:HOH:O	2.02	0.60
1:A:238:LEU:CD2	1:A:300:PHE:CZ	2.82	0.59
1:A:300:PHE:HE1	1:A:412:PHE:HB2	1.66	0.59
2:B:145:VAL:HG22	2:B:194:SER:HA	1.84	0.59
1:A:498:ILE:O	1:A:502:GLN:HA	2.02	0.59
3:C:54:GLY:O	3:C:55:ALA:HB3	2.02	0.59
2:B:39:GLN:NE2	3:C:42:GLN:HE22	1.99	0.59
1:A:487:GLU:CG	4:A:603:HOH:O	2.51	0.58
1:A:501:LEU:HD12	1:A:501:LEU:H	1.63	0.58
1:A:211:LYS:HD3	4:A:643:HOH:O	2.03	0.58
1:A:427:HIS:ND1	1:A:472:SER:HB2	2.19	0.57
1:A:491:ASN:O	4:A:601:HOH:O	2.17	0.57
1:A:366:LYS:HG3	4:A:636:HOH:O	2.05	0.57
1:A:429:LYS:HG3	4:A:639:HOH:O	2.06	0.56
3:C:83:GLU:OE2	4:C:360:HOH:O	2.18	0.56
3:C:36:LEU:HD12	3:C:96:THR:HA	1.89	0.55
1:A:341:GLU:HG2	2:B:53:ASN:HB2	1.89	0.54
3:C:26:SER:HB3	4:C:326:HOH:O	2.08	0.54
3:C:217:GLU:CG	4:C:352:HOH:O	2.35	0.54
2:B:124:LYS:CE	4:B:301:HOH:O	2.57	0.53
1:A:393:LEU:O	1:A:397:LEU:HB2	2.08	0.53
3:C:127:GLU:HB2	4:C:321:HOH:O	2.09	0.53
1:A:186:LYS:HG3	1:A:210:ILE:HD13	1.91	0.53
3:C:41:GLN:HB2	3:C:51:LEU:HD11	1.91	0.53
1:A:300:PHE:CE2	1:A:409:THR:CG2	2.78	0.52
2:B:207:ALA:HB2	2:B:214:LYS:HD3	1.90	0.52
1:A:361:ASP:O	4:A:636:HOH:O	2.19	0.52
1:A:300:PHE:CE1	1:A:412:PHE:HB2	2.45	0.51
2:B:191:THR:HG23	4:B:347:HOH:O	2.10	0.51
2:B:62:ASP:CB	4:B:397:HOH:O	2.59	0.50



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
2:B:208:HIS:HD2	2:B:211:SER:OG	1.95	0.49
2:B:62:ASP:HB3	4:B:397:HOH:O	2.11	0.49
1:A:419:HIS:HB3	1:A:423:ARG:NH2	2.27	0.49
1:A:381:SER:CB	4:A:640:HOH:O	2.60	0.49
3:C:154:ILE:HD12	3:C:159:ARG:HG3	1.95	0.49
1:A:362:GLU:O	1:A:362:GLU:HG3	2.11	0.49
1:A:186:LYS:CE	4:A:638:HOH:O	2.62	0.48
1:A:501:LEU:C	1:A:502:GLN:NE2	2.66	0.48
3:C:38:GLN:OE1	3:C:53:HIS:O	2.32	0.48
3:C:95:SER:HA	3:C:100:TRP:CD1	2.49	0.48
1:A:500:VAL:CB	1:A:501:LEU:HD12	2.43	0.48
1:A:504:LYS:HB2	1:A:504:LYS:HE3	1.75	0.46
1:A:300:PHE:CD2	1:A:409:THR:CG2	2.97	0.46
3:C:126:SER:O	3:C:130:THR:HG23	2.16	0.46
1:A:186:LYS:NZ	4:A:638:HOH:O	2.25	0.45
2:B:62:ASP:HA	4:B:397:HOH:O	2.16	0.45
1:A:501:LEU:CD1	1:A:501:LEU:H	2.23	0.45
3:C:167:TRP:CD1	3:C:167:TRP:N	2.84	0.45
2:B:12:VAL:CG2	2:B:18:ARG:HG3	2.47	0.44
1:A:419:HIS:HB2	4:A:628:HOH:O	2.18	0.44
2:B:124:LYS:HE2	4:B:301:HOH:O	2.17	0.44
1:A:237:LYS:HB2	1:A:303:MET:HE1	1.99	0.44
1:A:477:GLN:HG2	4:A:611:HOH:O	2.18	0.44
3:C:149:ASN:HD21	3:C:151:LYS:HE3	1.83	0.44
2:B:65:THR:CG2	4:B:397:HOH:O	2.56	0.44
2:B:191:THR:CG2	4:B:347:HOH:O	2.66	0.43
1:A:500:VAL:HB	1:A:501:LEU:CD1	2.48	0.43
3:C:54:GLY:C	3:C:56:SER:H	2.22	0.43
2:B:83:MET:HB3	2:B:86:LEU:HD21	2.00	0.43
2:B:146:THR:HB	2:B:191:THR:HG22	2.01	0.43
2:B:32:TYR:O	2:B:72:ARG:NH2	2.49	0.43
2:B:128:PRO:CB	2:B:154:TYR:HB3	2.43	0.43
1:A:497:LEU:O	1:A:503:MET:HG3	2.19	0.42
1:A:195:HIS:CB	1:A:343:LEU:HD11	2.49	0.42
1:A:419:HIS:HB3	1:A:423:ARG:HH22	1.83	0.42
3:C:24:ARG:HA	3:C:73:THR:O	2.20	0.42
2:B:180:GLN:O	2:B:180:GLN:HG2	2.19	0.42
2:B:146:THR:HB	2:B:191:THR:HB	2.02	0.42
1:A:415:LYS:HE3	1:A:415:LYS:HB3	1.39	0.41
1:A:500:VAL:C	1:A:501:LEU:CD1	2.66	0.41
3:C:12:ALA:HB1	3:C:111:LYS:HG3	2.03	0.41



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:419:HIS:CB	4:A:628:HOH:O	2.68	0.41
3:C:13:VAL:HG11	3:C:82:VAL:HG21	2.03	0.41
1:A:451:GLN:NE2	1:A:453:ASP:HB3	2.25	0.41
2:B:19:LYS:HG3	2:B:82:GLU:HG2	2.03	0.41
1:A:240:HIS:HE2	1:A:489:HIS:HE2	1.69	0.41
1:A:300:PHE:CE1	1:A:412:PHE:CB	3.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	284/526~(54%)	274~(96%)	10 (4%)	0	100	100
2	В	219/258~(85%)	212~(97%)	7(3%)	0	100	100
3	С	215/238~(90%)	208~(97%)	7 (3%)	0	100	100
All	All	718/1022 (70%)	694 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	279/511~(55%)	248 (89%)	31 (11%)	6 7



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	В	185/216~(86%)	162 (88%)	23 (12%)	4 5
3	С	190/209~(91%)	171 (90%)	19 (10%)	7 9
All	All	654/936~(70%)	581 (89%)	73 (11%)	6 6

Continued from previous page...

All (73) residues with a non-rotameric sidechain are listed below:

\mathbf{Mol}	Chain	Res	Type
1	А	162	ASP
1	А	164	LEU
1	А	213	ILE
1	А	221	LYS
1	А	222	SER
1	А	224	CYS
1	А	226	ASP
1	А	230	ASP
1	А	238	LEU
1	А	314	LEU
1	А	316	LYS
1	А	322	GLU
1	А	327	LYS
1	А	330	MET
1	А	347	ASN
1	А	390	SER
1	А	392	LEU
1	А	394	LEU
1	А	397	LEU
1	А	415	LYS
1	А	416	GLU
1	А	418	LYS
1	А	455	LEU
1	А	458	ARG
1	А	465	GLU
1	А	475	LEU
1	A	482	THR
1	А	487	GLU
1	А	501	LEU
1	A	502	GLN
1	А	504	LYS
2	В	4	LEU
2	В	18	ARG
2	В	65	THR



Mol	Chain	Res	Type
2	В	84	SER
2	В	97	THR
2	В	98	ARG
2	В	125	THR
2	В	126	THR
2	В	145	VAL
2	В	146	THR
2	В	159	VAL
2	В	161	LEU
2	В	168	LEU
2	В	180	GLN
2	В	183	LEU
2	В	187	SER
2	В	191	THR
2	В	193	THR
2	В	194	SER
2	В	201	SER
2	В	205	ASN
2	В	214	LYS
2	В	217	LYS
3	С	7	SER
3	С	13	VAL
3	С	14	SER
3	С	24	ARG
3	С	89	MET
3	С	97	LYS
3	С	98	VAL
3	С	107	LYS
3	С	109	GLU
3	С	118	THR
3	C	130	THR
3	C	159	ARG
3	С	160	GLN
3	С	161	ASN
3	С	167	TRP
3	С	174	ASP
3	C	197	THR
3	С	203	LYS
3	C	207	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:



Mol	Chain	Res	Type
1	А	165	GLN
1	А	195	HIS
1	А	342	GLN
1	А	347	ASN
1	А	365	HIS
1	А	439	GLN
1	А	451	GLN
1	А	502	GLN
2	В	39	GLN
2	В	180	GLN
2	В	208	HIS
3	С	17	GLN
3	С	38	GLN
3	С	57	ASN
3	С	149	ASN
3	С	160	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

The following chains have linkage breaks:



Mol	Chain	Number of breaks
3	С	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	С	50:LEU	С	51:LEU	Ν	1.16
1	С	51:LEU	С	52:ILE	N	1.16



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	288/526~(54%)	0.47	15 (5%) 27 34	24, 67, 93, 117	0
2	В	221/258~(85%)	0.46	8 (3%) 42 49	10, 41, 61, 72	8 (3%)
3	С	217/238 (91%)	0.26	4 (1%) 68 74	29, 46, 66, 106	0
All	All	726/1022 (71%)	0.41	27 (3%) 41 48	10, 49, 87, 117	8 (1%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	140	THR	15.2
2	В	138	GLY	12.1
2	В	142	GLY	11.5
2	В	137	CYS	8.9
1	А	300	PHE	7.6
2	В	141	THR	5.9
1	А	301	LYS	5.9
3	С	217	GLU	3.7
1	А	230	ASP	3.5
1	А	446	ILE	3.4
2	В	139	ASP	3.3
1	А	433	ASP	3.2
3	С	24	ARG	3.1
1	А	396	ASN	3.1
2	В	143	SER	3.0
1	А	302	LYS	3.0
2	В	136	VAL	2.7
1	А	236	LYS	2.5
1	А	392	LEU	2.5
1	А	233	ALA	2.4
1	А	227	ILE	2.3
1	A	388	GLN	2.2
1	А	235	ILE	2.2



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	С	21	ILE	2.2
3	С	74	ASP	2.1
1	А	319	LYS	2.1
1	А	402	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

