



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 29, 2024 – 09:09 AM EST

PDB ID : 5U00  
Title : CRYSTAL STRUCTURE OF HUMAN PHOSPHODIESTERASE 2A IN COMPLEX WITH 3,3-difluoro-1-[(4-fluoro-3-iodophenyl)carbonyl]-5-{5-methyl-[1,2,4]triazolo[1,5-a]pyrimidin-7-yl}piperidine  
Authors : Xu, R.; Aertgeerts, K.  
Deposited on : 2016-11-22  
Resolution : 1.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2579 (1.44-1.40)
Clashscore	141614	2696 (1.44-1.40)
Ramachandran outliers	138981	2632 (1.44-1.40)
Sidechain outliers	138945	2631 (1.44-1.40)
RSRZ outliers	127900	2528 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	 94% 5% .
1	B	344	 92% 6% ..
1	C	344	 90% 8% .
1	D	344	 93% 5% .

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called cGMP-dependent 3',5'-cyclic phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2786	1774	477	509	26	0	0	0
1	B	339	2775	1768	475	506	26	0	0	0
1	C	339	2775	1768	475	506	26	0	0	0
1	D	336	2751	1755	472	499	25	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	576	SER	-	expression tag	UNP O00408
A	577	ALA	-	expression tag	UNP O00408
A	578	MET	-	expression tag	UNP O00408
B	576	SER	-	expression tag	UNP O00408
B	577	ALA	-	expression tag	UNP O00408
B	578	MET	-	expression tag	UNP O00408
C	576	SER	-	expression tag	UNP O00408
C	577	ALA	-	expression tag	UNP O00408
C	578	MET	-	expression tag	UNP O00408
D	576	SER	-	expression tag	UNP O00408
D	577	ALA	-	expression tag	UNP O00408
D	578	MET	-	expression tag	UNP O00408

- Molecule 2 is [(5S)-3,3-difluoro-5-(5-methyl[1,2,4]triazolo[1,5-a]pyrimidin-7-yl)piperidin-1-yl][(4-fluoro-3-iodophenyl)methanone (three-letter code: 7OV) (formula: C<sub>18</sub>H<sub>15</sub>F<sub>3</sub>IN<sub>5</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	I	N			O
2	A	1	Total	C	F	I	N	O	0	0
			28	18	3	1	5	1		
2	B	1	Total	C	F	I	N	O	0	0
			28	18	3	1	5	1		
2	C	1	Total	C	F	I	N	O	0	0
			28	18	3	1	5	1		
2	D	1	Total	C	F	I	N	O	0	0
			28	18	3	1	5	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	A	1	Total	Mg	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Mg 1	0	0
4	C	1	Total 1	Mg 1	0	0
4	D	1	Total 1	Mg 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	276	Total 276	O 276	0	0
5	B	247	Total 247	O 247	0	0
5	C	244	Total 244	O 244	0	0
5	D	245	Total 245	O 245	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cGMP-dependent 3',5'-cyclic phosphodiesterase




- Molecule 1: cGMP-dependent 3',5'-cyclic phosphodiesterase



- Molecule 1: cGMP-dependent 3',5'-cyclic phosphodiesterase



- Molecule 1: cGMP-dependent 3',5'-cyclic phosphodiesterase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.99Å 73.25Å 91.21Å 109.38° 90.79° 91.10°	Depositor
Resolution (Å)	65.53 – 1.41 65.52 – 1.33	Depositor EDS
% Data completeness (in resolution range)	93.1 (65.53-1.41) 86.1 (65.52-1.33)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.31 (at 1.33Å)	Xtrriage
Refinement program	PHENIX (1.10.1-2155_1692: ???)	Depositor
R, $R_{free}$	0.199 , 0.228 0.199 , 0.228	Depositor DCC
$R_{free}$ test set	13398 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.5	Xtrriage
Anisotropy	0.255	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 48.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12219	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, 7OV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/2853	0.51	0/3849
1	B	0.34	0/2842	0.51	0/3834
1	C	0.32	0/2842	0.50	1/3834 (0.0%)
1	D	0.33	0/2818	0.50	0/3802
All	All	0.33	0/11355	0.50	1/15319 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	808	ASP	CB-CG-OD1	5.27	123.04	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2786	0	2728	14	0
1	B	2775	0	2718	16	0
1	C	2775	0	2718	12	0
1	D	2751	0	2701	9	0
2	A	28	0	0	1	0
2	B	28	0	0	1	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	28	0	0	0	0
2	D	28	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	276	0	0	2	0
5	B	247	0	0	5	0
5	C	244	0	0	0	0
5	D	245	0	0	2	0
All	All	12219	0	10865	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:824:GLU:HG2	1:A:828:LYS:HE2	1.74	0.69
1:C:892:ARG:NH1	1:C:893:GLU:OE2	2.27	0.68
1:D:593:VAL:HA	1:D:596:ILE:HD12	1.84	0.59
1:A:790:ARG:O	1:A:796:HIS:HE1	1.84	0.59
1:B:856:PRO:HG3	1:B:902:PHE:CD1	2.38	0.59
1:A:794:GLN:OE1	1:A:797:ARG:NH1	2.37	0.58
1:C:751:ARG:CZ	1:C:751:ARG:H	2.16	0.58
1:A:685:GLU:OE1	1:A:796:HIS:HD2	1.85	0.58
1:D:682:GLU:OE1	1:D:756:ARG:NH2	2.38	0.56
1:D:859:GLN:HG2	1:D:895:TRP:CE2	2.41	0.56
1:A:864:GLU:OE2	1:A:892:ARG:NH2	2.30	0.55
1:B:905:ARG:NH2	5:B:1101:HOH:O	2.30	0.55
1:A:892:ARG:HH12	1:A:896:THR:CG2	2.21	0.54
1:C:771:ALA:O	1:C:775:ARG:HG3	2.09	0.53
1:A:582:TYR:CE1	1:A:641:ARG:HG3	2.44	0.52
1:B:814:LYS:O	1:B:887:ARG:NH1	2.43	0.52
1:B:724:SER:N	5:B:1107:HOH:O	2.44	0.50
1:A:778:LYS:HG2	5:A:1105:HOH:O	2.12	0.49
1:C:821:LYS:NZ	1:C:824:GLU:OE2	2.38	0.49

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:828:LYS:NZ	5:A:1103:HOH:O	2.47	0.48
1:B:872:LYS:NZ	1:B:876:ASP:OD1	2.41	0.48
1:B:692:SER:O	1:B:696:HIS:HB3	2.13	0.47
1:B:892:ARG:HH12	1:B:896:THR:HG21	1.80	0.47
1:D:699:ASP:O	1:D:701:ARG:HD2	2.14	0.47
1:B:630:ASN:ND2	5:B:1109:HOH:O	2.47	0.46
1:A:817:LYS:HG3	1:A:820:ARG:HH12	1.81	0.45
1:B:578:MET:SD	1:B:578:MET:N	2.90	0.44
1:C:692:SER:O	1:C:696:HIS:HB3	2.17	0.44
1:B:827:TYR:OH	2:B:1001:7OV:I27	3.06	0.43
1:B:705:ASN:O	1:B:709:VAL:HG23	2.18	0.43
1:D:790:ARG:HG3	5:D:1203:HOH:O	2.17	0.43
1:C:746:PHE:CE2	1:C:757:MET:HG2	2.54	0.43
1:B:892:ARG:HH12	1:B:896:THR:CG2	2.32	0.42
1:B:709:VAL:HG22	1:B:722:GLU:OE2	2.20	0.42
1:A:852:LYS:HA	1:A:852:LYS:HD3	1.92	0.42
1:A:648:LYS:HG3	1:D:790:ARG:HD2	2.02	0.42
1:A:892:ARG:HH12	1:A:896:THR:HG23	1.85	0.42
1:B:617:MET:HG3	5:B:1229:HOH:O	2.20	0.41
1:C:791:ASN:N	1:C:791:ASN:HD22	2.19	0.41
1:C:664:VAL:HG13	1:C:807:CYS:HB3	2.02	0.41
1:C:644:LEU:O	1:C:648:LYS:HG2	2.20	0.41
1:B:797:ARG:HD2	5:B:1268:HOH:O	2.19	0.41
1:C:859:GLN:HG2	1:C:895:TRP:CE2	2.55	0.41
1:A:827:TYR:OH	2:A:1001:7OV:I27	3.09	0.41
1:D:617:MET:HG2	5:D:1145:HOH:O	2.19	0.41
1:D:664:VAL:HG13	1:D:807:CYS:HB3	2.03	0.41
1:B:916:LEU:HD23	1:B:916:LEU:HA	1.90	0.41
1:C:626:MET:HG2	1:C:672:TYR:CD2	2.57	0.40
1:D:582:TYR:OH	1:D:637:PRO:HB3	2.21	0.40
1:C:834:GLY:HA3	1:C:848:MET:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/344 (98%)	336 (99%)	3 (1%)	0	100	100
1	B	337/344 (98%)	332 (98%)	5 (2%)	0	100	100
1	C	337/344 (98%)	332 (98%)	3 (1%)	2 (1%)	25	7
1	D	334/344 (97%)	332 (99%)	2 (1%)	0	100	100
All	All	1347/1376 (98%)	1332 (99%)	13 (1%)	2 (0%)	51	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	589	GLY
1	C	590	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/310 (99%)	305 (99%)	2 (1%)	84	65
1	B	306/310 (99%)	302 (99%)	4 (1%)	69	41
1	C	306/310 (99%)	300 (98%)	6 (2%)	55	22
1	D	303/310 (98%)	299 (99%)	4 (1%)	69	41
All	All	1222/1240 (98%)	1206 (99%)	16 (1%)	69	41

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	752	LYS
1	A	892	ARG
1	B	752	LYS
1	B	782	LYS
1	B	797	ARG
1	B	892	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	578	MET
1	C	617	MET
1	C	624	GLN
1	C	751	ARG
1	C	790	ARG
1	C	852	LYS
1	D	676	GLU
1	D	701	ARG
1	D	797	ARG
1	D	909	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	796	HIS
1	A	859	GLN
1	B	630	ASN
1	B	875	GLN
1	C	599	ASN
1	C	772	HIS
1	C	791	ASN
1	C	859	GLN
1	D	599	ASN
1	D	679	ASN
1	D	842	ASN
1	D	859	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	7OV	A	1001	-	26,31,31	1.47	4 (15%)	24,47,47	1.96	7 (29%)
2	7OV	B	1001	-	26,31,31	1.66	5 (19%)	24,47,47	1.86	8 (33%)
2	7OV	C	1001	-	26,31,31	1.56	4 (15%)	24,47,47	1.91	5 (20%)
2	7OV	D	1001	-	26,31,31	1.59	4 (15%)	24,47,47	2.05	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7OV	A	1001	-	-	0/12/26/26	0/4/4/4
2	7OV	B	1001	-	-	0/12/26/26	0/4/4/4
2	7OV	C	1001	-	-	0/12/26/26	0/4/4/4
2	7OV	D	1001	-	-	0/12/26/26	0/4/4/4

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1001	7OV	C1-C2	4.64	1.47	1.38
2	C	1001	7OV	C1-C2	4.62	1.47	1.38
2	D	1001	7OV	C1-C2	4.55	1.47	1.38
2	A	1001	7OV	C1-C2	4.26	1.46	1.38
2	B	1001	7OV	C1-C6	-4.18	1.33	1.38
2	D	1001	7OV	C1-C6	-3.93	1.34	1.38
2	C	1001	7OV	C1-C6	-3.72	1.34	1.38
2	A	1001	7OV	C1-C6	-3.38	1.34	1.38
2	C	1001	7OV	C4-N7	-2.62	1.31	1.35
2	B	1001	7OV	C16-N15	-2.52	1.43	1.46
2	D	1001	7OV	C4-N7	-2.47	1.32	1.35

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	7OV	C4-N7	-2.32	1.32	1.35
2	B	1001	7OV	C4-N7	-2.23	1.32	1.35
2	D	1001	7OV	C16-N15	-2.18	1.44	1.46
2	C	1001	7OV	C2-N3	-2.15	1.29	1.33
2	A	1001	7OV	C2-N3	-2.08	1.29	1.33
2	B	1001	7OV	C2-N3	-2.06	1.29	1.33

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1001	7OV	C24-C25-I27	-5.68	115.51	120.75
2	B	1001	7OV	F28-C24-C25	4.02	121.63	118.42
2	D	1001	7OV	C24-C25-I27	-3.87	117.19	120.75
2	D	1001	7OV	C11-C2-C1	-3.78	116.03	121.81
2	A	1001	7OV	F28-C24-C25	3.46	121.18	118.42
2	A	1001	7OV	C20-C19-N15	3.45	123.09	118.72
2	D	1001	7OV	F17-C13-C12	-3.39	105.95	109.41
2	B	1001	7OV	C24-C25-I27	-3.38	117.63	120.75
2	D	1001	7OV	C20-C19-N15	3.29	122.89	118.72
2	A	1001	7OV	C11-C2-C1	-3.27	116.81	121.81
2	C	1001	7OV	C26-C25-I27	3.21	124.53	118.61
2	D	1001	7OV	O21-C19-C20	-2.95	114.50	120.23
2	B	1001	7OV	C20-C19-N15	2.94	122.45	118.72
2	A	1001	7OV	C26-C25-I27	2.86	123.88	118.61
2	D	1001	7OV	C26-C25-I27	2.85	123.87	118.61
2	D	1001	7OV	F28-C24-C25	2.81	120.66	118.42
2	C	1001	7OV	F18-C13-C12	2.80	112.27	109.41
2	D	1001	7OV	C12-C10-C6	2.74	115.50	112.06
2	C	1001	7OV	C11-C2-C1	-2.72	117.65	121.81
2	A	1001	7OV	F17-C13-C12	-2.70	106.66	109.41
2	B	1001	7OV	C11-C2-C1	-2.62	117.82	121.81
2	B	1001	7OV	C12-C10-C6	2.55	115.26	112.06
2	A	1001	7OV	O21-C19-C20	-2.53	115.30	120.23
2	A	1001	7OV	C24-C25-I27	-2.39	118.55	120.75
2	B	1001	7OV	O21-C19-C20	-2.32	115.72	120.23
2	B	1001	7OV	C26-C25-I27	2.31	122.87	118.61
2	C	1001	7OV	C20-C19-N15	2.29	121.63	118.72
2	D	1001	7OV	C1-C2-N3	2.27	125.26	122.57
2	B	1001	7OV	C22-C20-C26	2.09	121.71	119.24

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	7OV	1	0
2	B	1001	7OV	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	341/344 (99%)	-0.20	4 (1%) 79 77	9, 21, 37, 50	0
1	B	339/344 (98%)	-0.13	3 (0%) 84 82	10, 22, 37, 49	0
1	C	339/344 (98%)	0.07	15 (4%) 34 34	11, 26, 47, 64	0
1	D	336/344 (97%)	-0.11	9 (2%) 54 53	11, 23, 42, 57	0
All	All	1355/1376 (98%)	-0.09	31 (2%) 60 60	9, 23, 41, 64	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	586	LEU	5.0
1	D	589	GLY	4.5
1	B	723	GLY	4.2
1	D	582	TYR	4.0
1	D	588	ASP	3.9
1	C	680	TYR	3.8
1	C	588	ASP	3.4
1	B	722	GLU	3.1
1	D	590	ILE	2.9
1	C	589	GLY	2.9
1	D	586	LEU	2.8
1	B	721	SER	2.8
1	C	578	MET	2.8
1	D	587	HIS	2.8
1	C	595	ALA	2.7
1	A	598	SER	2.6
1	C	582	TYR	2.6
1	D	584	LYS	2.6
1	C	594	ALA	2.6
1	C	843	ARG	2.6
1	C	590	ILE	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	579	ASP	2.4
1	C	902	PHE	2.3
1	D	914	ASP	2.3
1	A	909	SER	2.2
1	C	593	VAL	2.2
1	C	580	ASP	2.2
1	A	843	ARG	2.1
1	C	909	SER	2.1
1	D	915	PHE	2.1
1	A	839	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	7OV	A	1001	28/28	0.99	0.06	10,15,19,22	0
2	7OV	B	1001	28/28	0.99	0.08	12,16,23,25	0
2	7OV	C	1001	28/28	0.99	0.06	12,17,21,27	0
2	7OV	D	1001	28/28	0.99	0.06	11,15,21,22	0
4	MG	C	1003	1/1	0.99	0.05	12,12,12,12	0
3	ZN	B	1002	1/1	1.00	0.05	13,13,13,13	0
3	ZN	C	1002	1/1	1.00	0.04	13,13,13,13	0
3	ZN	D	1002	1/1	1.00	0.04	13,13,13,13	0
4	MG	A	1003	1/1	1.00	0.06	9,9,9,9	0
4	MG	B	1003	1/1	1.00	0.05	10,10,10,10	0
3	ZN	A	1002	1/1	1.00	0.03	12,12,12,12	0
4	MG	D	1003	1/1	1.00	0.06	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around 7OV B 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 7OV C 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.